

# 1

## Introduction to the Galois Theory of Linear Differential Equations

Michael F. Singer

*Department of Mathematics  
North Carolina State University  
Raleigh, NC 27695-8205  
singer@math.ncsu.edu*

### 1.1 Introduction

This paper is an expanded version of the 10 lectures I gave as the 2006 London Mathematical Society Invited Lecture Series at the Heriot-Watt University, 31 July - 4 August 2006<sup>†</sup>. My goal was to give the audience an introduction to the algebraic, analytic and algorithmic aspects of the Galois theory of linear differential equations by focusing on some of the main ideas and philosophies and on examples. There are several texts ([Beu92, Kap76, Kol76, Mag94, dPS03] to name a few) that give detailed expositions and I hope that the taste offered here will encourage the reader to dine more fully with one of these.

The rest of the paper is organized as follows. In Section 1.2, *What is a Linear Differential Equation?*, I discuss three ways to think about linear differential equations: scalar equations, linear systems and differential modules. Just as it is useful to think of linear maps in terms of linear equations, matrices and associated modules, it will be helpful in future sections to go back and forth between the different ways of presenting linear differential equations.

In Section 1.3, *Basic Galois Theory and Applications*, I will give the basic definitions and describe the Galois correspondence. In addition I will describe the notion of monodromy and its relation to the Galois

<sup>†</sup> I would like to thank the London Mathematical Society for inviting me to present these lectures, the Heriot-Watt University for hosting them and the International Centre for the Mathematical Sciences for sponsoring a mini-programme on the Algebraic Theory of Differential Equations to complement these talks. Thanks also go to the Edinburgh Mathematical Society and the Royal Society of Edinburgh, who provided support for some of the participants and to Chris Eilbeck, Malcolm MacCallum, Alexandre Mikhailov, and Sergey Tsarev for organizing these events. This material is based upon work supported by the National Science Foundation under Grant No. 0096842 and 0634123

theory. I will end by giving several applications and ramifications, one of which will be to answer the question *Although  $y = \cos x$  satisfies  $y'' + y = 0$ , why doesn't  $\sec x$  satisfy a linear differential equation?*

In Section 1.4, *Local Galois Theory*, I will discuss the formal solution of a linear differential equation at a singular point and describe the asymptotics which allow one to associate with it an analytic solution in a small enough sector at that point. This will involve a discussion of Gevrey asymptotics, the Stokes phenomenon and its relation to the Galois theory. A motivating question (which we will answer in this section) is *In what sense does the function*

$$f(x) = \int_0^\infty \frac{1}{1+\zeta} e^{-\frac{\zeta}{x}} d\zeta$$

*represent the divergent series*

$$\sum_{n \geq 0} (-1)^n n! x^{n+1} ?$$

In Section 1.5, *Algorithms*, I turn to a discussion of algorithms that allow us to determine properties of a linear differential equation and its Galois group. I will show how category theory, and in particular the tannakian formalism, points us in a direction that naturally leads to algorithms. I will also discuss algorithms to find “closed form solutions” of linear differential equations.

In Section 1.6, *Inverse Problems*, I will consider the problem of which groups can appear as Galois groups of linear differential equations. I will show how monodromy and the ideas in Section 1.4 play a role as well as ideas from Lie theory.

In Section 1.7, *Families of Linear Differential Equations*, I will look at linear differential equations that contain parameters and ask *How does the Galois group change as we vary the parameters?* This will lead to a discussion of a generalization of the above theory to a Galois theory of parameterized linear differential equations.

## 1.2 What is a Linear Differential Equation?

I will develop an algebraic setting for the study of linear differential equations. Although there are many interesting questions concerning differential equations in characteristic  $p$  [MvdP03a, MvdP03b, dP95, dP96], we will restrict ourselves throughout this paper, without further mention, to fields of characteristic 0. I begin with some basic definitions.

**Definition 1.2.1** 1) A differential ring  $(R, \Delta)$  is a ring  $R$  with a set  $\Delta = \{\partial_1, \dots, \partial_m\}$  of maps (derivations)  $\partial_i : R \rightarrow R$ , such that

- (i)  $\partial_i(a+b) = \partial_i(a) + \partial_i(b)$ ,  $\partial_i(ab) = \partial_i(a)b + a\partial_i(b)$  for all  $a, b \in R$ ,  
and
- (ii)  $\partial_i\partial_j = \partial_j\partial_i$  for all  $i, j$ .

2) The ring  $C_R = \{c \in R \mid \partial(c) = 0 \ \forall \ \partial \in \Delta\}$  is called the ring of constants of  $R$ .

When  $m = 1$ , we say  $R$  is an *ordinary differential ring*  $(R, \partial)$ . We frequently use the notation  $a'$  to denote  $\partial(a)$  for  $a \in R$ . A differential ring that is also a field is called a *differential field*. If  $k$  is a differential field, then  $C_k$  is also a field.

**Examples 1.2.2** 1)  $(C^\infty(\mathbb{R}^m), \Delta = \{\frac{\partial}{\partial x_1}, \dots, \frac{\partial}{\partial x_m}\})$  = infinitely differentiable functions on  $\mathbb{R}^m$ .

2)  $(\mathbb{C}(x_1, \dots, x_m), \Delta = \{\frac{\partial}{\partial x_1}, \dots, \frac{\partial}{\partial x_m}\})$  = field of rational functions

3)  $(\mathbb{C}[[x]], \frac{\partial}{\partial x})$  = ring of formal power series

$$\mathbb{C}((x)) = \text{quotient field of } \mathbb{C}[[x]] = \mathbb{C}[[x]][\frac{1}{x}]$$

4)  $(\mathbb{C}\{\{x\}\}, \frac{\partial}{\partial x})$  = ring of germs of convergent series

$$\mathbb{C}(\{x\}) = \text{quotient field of } \mathbb{C}\{\{x\}\} = \mathbb{C}\{\{x\}\}[\frac{1}{x}]$$

5)  $(\mathcal{M}_{\mathcal{O}}, \Delta = \{\frac{\partial}{\partial x_1}, \dots, \frac{\partial}{\partial x_m}\})$  = field of functions meromorphic on  $\mathcal{O}_{\text{open, connected}} \subset \mathbb{C}^m$

The following result of Seidenberg [Sei58, Sei69] shows that many examples reduce to Example 5) above:

**Theorem 1.2.3** Any differential field  $k$ , finitely generated over  $\mathbb{Q}$ , is isomorphic to a differential subfield of some  $\mathcal{M}_{\mathcal{O}}$ .

We wish to consider and compare three different versions of the notion of a linear differential equation.

**Definition 1.2.4** Let  $(k, \partial)$  be a differential field.

- (i) A scalar linear differential equation is an equation of the form

$$L(y) = a_n y^{(n)} + \dots + a_0 y = 0, \quad a_i \in k.$$

- (ii) A matrix linear differential equation is an equation of the form

$$Y' = AY, \quad A \in \text{gl}_n(k)$$

where  $\text{gl}_n(k)$  denotes the ring of  $n \times n$  matrices with entries in  $k$ .

- (iii) A differential module of dimension  $n$  is an  $n$ -dimensional  $k$ -vector space  $M$  with a map  $\partial : M \rightarrow M$  satisfying

$$\partial(fm) = f'm + f\partial m \text{ for all } f \in k, m \in M.$$

We shall show that these three notions are equivalent and give some further properties.

**From scalar to matrix equations:** Let  $L(y) = y^{(n)} + a_{n-1}y^{(n-1)} + \dots + a_0y = 0$ . If we let  $y_1 = y, y_2 = y', \dots, y_n = y^{(n-1)}$ , then we have

$$\begin{pmatrix} y_1 \\ y_2 \\ \vdots \\ y_{n-1} \\ y_n \end{pmatrix}' = \begin{pmatrix} 0 & 1 & 0 & \dots & 0 \\ 0 & 0 & 1 & \dots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & 0 & \dots & 1 \\ -a_0 & -a_1 & -a_2 & \dots & -a_{n-1} \end{pmatrix} \begin{pmatrix} y_1 \\ y_2 \\ \vdots \\ y_{n-1} \\ y_n \end{pmatrix}$$

We shall write this last equation as  $Y' = A_L Y$  and refer  $A_L$  as the companion matrix of the scalar equation and the matrix equation as the companion equation. Clearly any solution of the scalar equation yields a solution of the companion equation and *vice versa*.

**From matrix equations to differential modules (and back):**

Given  $Y' = AY, A \in \text{gl}_n(k)$ , we construct a differential module in the following way: Let  $M = k^n$ ,  $e_1, \dots, e_n$  the usual basis. Define  $\partial e_i = -\sum_j a_{j,i} e_j$ , i.e.,  $\partial e = -A^t e$ . Note that if  $m = \sum_i f_i e_i$  then  $\partial m = \sum_i (f_i' - \sum_j a_{j,i} f_j) e_i$ . In particular, we have that  $\partial m = 0$  if and only if

$$\begin{pmatrix} f_1 \\ \vdots \\ f_n \end{pmatrix}' = A \begin{pmatrix} f_1 \\ \vdots \\ f_n \end{pmatrix}$$

It is this latter fact that motivates the seemingly strange definition of this differential module, which we denote by  $(M_A, \partial)$ .

Conversely, given a differential module  $(M, \partial)$ , select a basis  $e = (e_1, \dots, e_n)$ . Define  $A_M \in \text{gl}_n(k)$  by  $\partial e_i = \sum_j a_{j,i} e_j$ . This yields a matrix equation

$Y' = AY$ . If  $\bar{e} = (\bar{e}_1, \dots, \bar{e}_n)$  is another basis, we get another equation  $Y' = \bar{A}Y$ . If  $f$  and  $\bar{f}$  are vectors with respect to these two bases and  $\bar{f} = Bf$ ,  $B \in \text{GL}_n(k)$ , then

$$\bar{A} = B^{-1}AB - B^{-1}B'.$$

**Definition 1.2.5** Let  $(M_1, \partial_1)$  and  $(M_2, \partial_2)$  be differential modules.

- 1) A differential module homomorphism  $\phi : M_1 \rightarrow M_2$  is a  $k$ -linear map  $\phi$  such that  $\phi(\partial_1(m)) = \partial_2(\phi(m))$  for all  $m \in M_1$ .
- 2) The differential modules  $(M_1, \partial_1)$  and  $(M_2, \partial_2)$  are isomorphic if there exists a bijective differential homomorphism  $\phi : M_1 \rightarrow M_2$ .
- 3) Two differential equations  $Y' = A_1Y$  and  $Y' = A_2Y$  are equivalent if the differential modules  $M_{A_1}$  and  $M_{A_2}$  are isomorphic

Instead of equivalent, some authors use the term “gauge equivalent” or “of the same type”.

Differential modules offer us an opportunity to study linear differential equations in a basis-free environment. Besides being of theoretical interest, it is important in computations to know that a concept is independent of bases, since this allows one to then select a convenient basis in which to compute.

Before we show how one can recover a scalar equation from a differential module, we show that the standard constructions of linear algebra can be carried over to the context of differential modules. Let  $(M_1, \partial_1)$  and  $(M_2, \partial_2)$  be differential modules and assume that for certain bases these correspond to the equations  $Y' = A_1Y$  and  $Y' = A_2Y$ .

In Table 1.1 we list some standard linear algebra constructions and how they generalize to differential modules. In this table,  $I_{n_i}$  represents the  $n_i \times n_i$  identity matrix and for two matrices  $A = (a_{i,j})$  and  $B$ ,  $A \otimes B$  is the matrix where the  $i, j$ -entry of  $A$  is replaced by  $a_{i,j}B$ . Also note that if  $f \in \text{Hom}_k(M_1, M_2)$  then  $\partial(f) = 0$  if and only if  $f(\partial m_1) = \partial_2(f(m_2))$ , that is, if and only if  $f$  is a differential module homomorphism.

Referring to the table, it is not hard to show that  $\text{Hom}_k(M_1, M_2) \simeq M_1 \otimes M_2^*$  as differential modules. Furthermore, given  $(M, \partial)$  with  $\dim_k(M) = n$ , corresponding to a differential equation  $Y' = AY$ , we have that  $M \simeq \bigoplus_{i=1}^n \mathbf{1}_k$  if and only if there exist  $y_1, \dots, y_n$  in  $k^n$ , linearly independent over  $k$  such that  $y'_i = Ay_i$ .

Table 1.1.

Construction	$\partial$	Matrix Equation
$(M_1 \oplus M_2, \partial)$	$\partial(m_1 \oplus m_2) = \partial_1 m_1 \oplus \partial_2 m_2$	$Y' = \begin{pmatrix} A_1 & 0 \\ 0 & A_2 \end{pmatrix} Y$
$(M_1 \otimes M_2, \partial)$	$\partial(m_1 \otimes m_2) = \partial_1 m_1 \otimes m_2 + m_1 \otimes \partial_2 m_2$	$Y' = (A_1 \otimes I_{n_2} + I_{n_1} \otimes A_2) Y$
$(\text{Hom}_k(M_1, M_2), \partial)$	$\partial(f)(m) = f(\partial_1 m) - \partial_2(f(m))$	$Y' = A_1 Y - Y A_2$
$\mathbf{1}_k = (k, \partial) = \text{trivial differential module}$	$\partial \equiv 0$	$Y' = 0$
$(M^*, \partial) = \text{Hom}_k(M, \mathbf{1}_k)$	$\partial(f)(m) = f(\partial(m))$	$Y' = -A^T Y$

**From matrix to scalar linear differential equations:** Before I discuss the relationship between matrix and scalar linear differential equations, I need one more concept.

**Definition 1.2.6** *Let  $k$  be a differential field. The ring of differential operators over  $k = k[\partial]$  is the ring of noncommutative polynomials  $\{a_n \partial^n + \dots + a_1 \partial + a_0 \mid a_i \in k\}$  with coefficients in  $k$ , where multiplication is determined by  $\partial \cdot a = a' + a\partial$  for all  $a \in k$ .*

We shall refer to the degree of an element  $L \in k[\partial]$  as its order  $\text{ord} L$ . The following properties are not hard to check ([dPS03], Chapter 2.1):

**Lemma 1.2.7** *Let  $L_1 \neq 0, L_2 \in k[\partial]$ .*

- 1) *There exist unique  $Q, R \in k[\partial]$  with  $\text{ord} R < \text{ord} L_1$  such that  $L_2 = QL_1 + R$ .*
- 2) *Every left ideal of  $k[\partial]$  is of the form  $k[\partial]L$  for some  $L \in k[\partial]$ .*

We note that any differential module  $M$  can be considered a left  $k[\partial]$ -module and conversely, any left  $k[\partial]$ -module that is finite dimensional as a  $k$ -vector space is a differential module. In fact we have a stronger statement:

**Theorem 1.2.8** *(Cyclic Vector Theorem) Assume there exists an  $a \in k$  such that  $a' \neq 0$ . Every differential module is of the form  $k[\partial]/k[\partial]L$  for some  $L \in k[\partial]$ .*

This result has many proofs ([CK02, Cop34, Cop36, Del70, Jac37, Kat87a] to name a few), two of which can be found in Chapter 2.1 of [dPS03].

**Corollary 1.2.9** (*Systems to Scalar equations*) *Let  $k$  be as above. Every system  $Y' = AY$  is equivalent to a scalar equation  $L(y) = 0$ .*

*Proof* Let  $A^* = -A^t$ . Apply the Cyclic Vector Theorem to  $M_{A^*}$  to find an  $L \in k[\partial]$  such that  $M_{A^*} = k[\partial]/k[\partial]L$ . If  $A_L$  is the companion matrix of  $L$ , a calculation shows that  $M_A \simeq M_{A_L}$ .  $\square$

We note that the hypothesis that  $k$  contain an element  $a$  with  $a' \neq 0$  is necessary. If the derivation is trivial on  $k$ , then two matrix equations  $Y' = A_1Y$  and  $Y' = A_2Y$  are equivalent if and only if the matrices are similar. There exist many examples of matrices not similar to a companion matrix.

Before we leave (for now) our discussion of the ring  $k[\partial]$ , I wish to make two remarks.

First, we can define a map  $i : k[\partial] \rightarrow k[\partial]$  by  $i(\sum a_j \partial^j) = \sum (-1)^j \partial^j a_j$ . This map is an involution ( $i^2 = id$ ). Denoting  $i(L) = L^*$ , we have that  $(L_1 L_2)^* = L_2^* L_1^*$ . The operator  $L^*$  is referred to as the *adjoint* of  $L$ . Using the adjoint one sees that there is right euclidean division as well and that every right ideal of  $k[\partial]$  is also principal.

Second, Lemma 1.2.7.2 allows us to define

**Definition 1.2.10** *Let  $L_1, L_2 \in k[\partial]$ .*

- 1) *The least common left multiple  $LCLM(L_1, L_2)$  of  $L_1$  and  $L_2$  is the monic generator of  $k[\partial]L_1 \cap k[\partial]L_2$ .*
- 2) *The greatest common right divisor  $GCRD(L_1, L_2)$  of  $L_1$  and  $L_2$  is the monic generator of  $k[\partial]L_1 + k[\partial]L_2$ .*

A simple modification of the usual euclidean algorithm allows one to find these objects. One can also define least common right multiples and greatest common left divisors using right ideals.

**Solutions** I will give properties of solutions of scalar and matrix linear differential equations and define the notion of the solutions of a differential module. Let  $(k, \partial)$  be a differential field with constants  $C_k$  (see Definition 1.2.1).

**Lemma 1.2.11** *Let  $v_1, \dots, v_r \in k^n$  satisfy  $v'_i = Av_i$ ,  $A \in gl_n(k)$ . If  $v_1, \dots, v_r$  are  $k$ -linearly dependent, then they are  $C_k$ -linearly dependent.*

*Proof* Assume, by induction, that  $v_2, \dots, v_r$  are  $k$ -linearly independent and  $v_1 = \sum_{i=2}^r a_i v_i, a_i \in k$ . We then have

$$0 = v'_1 - Av_1 = \sum_{i=2}^r a'_i v_i + \sum_{i=2}^r a_i (v'_i - Av_i) = \sum_{i=2}^r a'_i v_i$$

so by assumption, each  $a'_i = 0$ .  $\square$

**Corollary 1.2.12** *Let  $(k, \partial)$  be a differential field with constants  $C_k$ ,  $A \in \mathfrak{gl}_n(k)$  and  $L \in k[\partial]$ .*

- 1) *The solution space  $\text{Soln}_k(Y' = AY)$  of  $Y' = AY$  in  $k^n$  is a  $C_k$ -vector space of dimension at most  $n$ .*
- 2) *The elements  $y_1, \dots, y_r \in k$  are linearly dependent over  $C_k$  if and only if the wronskian determinant*

$$wr(y_1, \dots, y_r) = \det \begin{pmatrix} y_1 & \dots & y_r \\ \vdots & \vdots & \vdots \\ y_1^{(r-1)} & \dots & y_r^{(r-1)} \end{pmatrix}$$

*is zero.*

- 3) *The solution space  $\text{Soln}_k(L(y) = 0) = \{y \in k \mid L(y) = 0\}$  of  $L(y) = 0$  in  $k$  is a  $C_k$ -vector space of dimension at most  $n$ .*

*Proof* 1) This follows immediately from Lemma 1.2.11.

2) If  $\sum_{i=1}^r c_i y_i = 0$  for some  $c_i \in C_k$ , not all zero, then  $\sum_{i=1}^r c_i y_i^{(j)} = 0$  for all  $j$ . Therefore,  $wr(y_1, \dots, y_r) = 0$ . Conversely if  $wr(y_1, \dots, y_r) = 0$ , then there exists a nonzero vector  $(a_0, \dots, a_{r-1})$  such that

$$(a_0, \dots, a_{r-1}) \begin{pmatrix} y_1 & \dots & y_r \\ \vdots & \vdots & \vdots \\ y_1^{(r-1)} & \dots & y_r^{(r-1)} \end{pmatrix} = 0$$

Therefore each  $y_i$  satisfies the scalar linear differential equation  $L(y) = a_{r-1}y^{(r-1)} + \dots + a_0y = 0$  so each vector  $v_i = (y_i, y'_i, \dots, y_i^{(r-1)})^t$  satisfies  $Y' = A_L Y$ , where  $A_L$  is the companion matrix. Lemma 1.2.11 implies that the  $v_i$  and therefore the  $y_i$  are linearly dependent over  $C_k$ .

- 3) Apply Lemma 1.2.11 to  $Y' = A_L Y$ .  $\square$

In general, the dimension of the solutions space of a linear differential equation in a field is less than  $n$ . In the next section we will construct



a field such that over this field the solutions space has dimension  $n$ . It will be useful to have the following definition

**Definition 1.2.13** *Let  $(k, \partial)$  be a differential field,  $A \in \mathfrak{gl}_n(k)$  and  $R$  a differential ring containing  $k$ . A matrix  $Z \in \mathrm{GL}_n(R)$  such that  $Z' = AZ$  is called a fundamental solution matrix of  $Y' = AY$ .*

Note that if  $R$  is a field and  $Z$  is a fundamental solution matrix, then the columns of  $Z$  form a  $C_R$ -basis of the solution space of  $Y' = AY$  in  $R$  and that this solution space has dimension  $n$ .

Let  $(M, \partial)$  be a differential module over  $k$ . We define the solution space  $\mathrm{Soln}_k(M)$  of  $(M, \partial)$  to be the kernel  $\ker_M \partial$  of  $\partial$  on  $M$ . As we have noted above, if  $\{e_i\}$  is a basis of  $M$  and  $Y' = AY$  is the associated matrix differential equation in this basis, then  $\sum_i v_i e_i \in \ker \partial$  if and only if  $v' = Av$  where  $v = (v_1, \dots, v_n)^t$ . If  $K \supset k$  is a differential extension of  $k$ , then  $K \otimes_k M$  can be given the structure of a  $K$ -differential module, where  $\partial(a \otimes m) = a' \otimes m + a \otimes \partial m$ . We then define  $\mathrm{Soln}_K(M)$  to be the kernel  $\ker(\partial, K \otimes_k M)$  of  $\partial$  on  $K \otimes_k M$ .

### 1.3 Basic Galois Theory and Applications

**Galois theory of polynomials** The idea behind the Galois theory of polynomials is to associate to a polynomial a group (the group of symmetries of the roots that preserve all the algebraic relations among these roots) and deduce properties of the roots from properties of this group (for example, a solvable group implies that the roots can be expressed in terms of radicals). This idea can be formalized in the following way.

Let  $k$  be a field (of characteristic 0 as usual) and let  $P(X) \in k[X]$  be a polynomial of degree  $n$  without repeated roots (i.e.,  $\mathrm{GCD}(P, P') = 1$ ). Let

$$S = k[X_1, \dots, X_n, \frac{1}{\prod (X_i - X_j)}]$$

(the reason for the term  $\frac{1}{\prod (X_i - X_j)}$  will be explained below) and let  $I = (P(X_1), \dots, P(X_n)) \triangleleft S$  be the ideal generated by the  $P(X_i)$ . The ring  $S/I$  is generated by  $n$  distinct roots of  $P$  but does not yet reflect the possible algebraic relations among these roots. We therefore consider any maximal ideal  $M$  in  $S$  containing  $I$ . This ideal can be thought of as a maximally consistent set of algebraic relations among the roots. We define

**Definition 1.3.1** 1) The splitting ring of the polynomial  $P$  over  $k$  is the ring

$$R = S/M = k[X_1, \dots, X_n, \frac{1}{\prod (X_i - X_j)}] / M.$$

2) The Galois group of  $P$  (or of  $R$  over  $k$ ) is the group of automorphisms  $\text{Aut}(R/k)$ .

Note that since  $M$  is a maximal ideal,  $R$  is actually a field. Furthermore, since  $S$  contains  $\frac{1}{\prod (X_i - X_j)}$ , the images of the  $X_i$  in  $R$  are *distinct* roots of  $P$ . So, in fact,  $R$  coincides with the usual notion of a splitting field (a field generated over  $k$  by the distinct roots of  $P$ ) and as such, is unique up to  $k$ -isomorphism. Therefore,  $R$  is independent of the choice of maximal ideal  $M$  containing  $I$ . We will follow a similar approach with linear differential equations.

**Galois theory of linear differential equations** Let  $(k, \partial)$  be a differential field and  $Y' = AY, A \in \text{gl}_n(k)$  a matrix differential equation over  $k$ . We now want the Galois group to be the group of symmetries of solutions preserving all algebraic and differential relations. We proceed in a way similar to the above.

Let

$$S = k[Y_{1,1}, \dots, Y_{n,n}, \frac{1}{\det(Y_{i,j})}]$$

where  $Y = (Y_{i,j})$  is an  $n \times n$  matrix of indeterminates. We define a derivation on  $S$  by setting  $Y' = AY$ . The columns of  $Y$  form  $n$  independent solutions of the matrix linear differential equation  $Y' = AY$  but we have not yet taken into account other possible algebraic and differential relations. To do this, let  $M$  be any maximal *differential* ideal and let  $R = S/M$ . We have now constructed a ring that satisfies the following definition

**Definition 1.3.2** Let  $(k, \partial)$  be a differential field and  $Y' = AY, A \in \text{gl}_n(k)$  a matrix differential equation over  $k$ . A Picard-Vessiot ring (PV-ring) for  $Y' = AY$  is a differential ring  $R$  over  $k$  such that

- (i)  $R$  is a simple differential ring (i.e., the only differential ideals are  $(0)$  and  $R$ ).
- (ii) There exists a fundamental matrix  $Z \in \text{GL}_n(R)$  for the equation  $Y' = AY$ .
- (iii)  $R$  is generated as a ring by  $k$ , the entries of  $Z$  and  $\frac{1}{\det Z}$ .

One can show that a PV-ring must be a domain and, if  $C_k$  is algebraically closed, then any PV-rings for the same equation are  $k$ -isomorphic as differential rings and  $C_k = C_R$ . Since a PV-ring is a domain we define a *PV-field*  $K$  to be the quotient field of a PV-ring. Assuming that  $C_k$  is algebraically closed this is the same as saying that  $K$  is generated over  $k$  by the entries of a fundamental solution matrix and that  $C_K = C_k$ . These facts are proven in Chapter 1.2 of [dPS03]. *From now on, we will always assume that  $C_k$  is algebraically closed* (see [Dyc05, HvdP95, Kol76, Ulm94] for results when this is not the case) .

**Examples 1.3.3** 1)  $k = \mathbb{C}(x)$   $x' = 1$   $\alpha \in \mathbb{C}$   $Y' = \frac{\alpha}{x}Y$   $S = k[Y, \frac{1}{Y}]$   $Y' = \frac{\alpha}{x}Y$

*Case 1:  $\alpha = \frac{n}{m}$ ,  $\text{GCD}(n, m) = 1$ . I claim that  $\mathcal{R} = k[Y, \frac{1}{Y}]/[Y^m - x^n] = k(x^{\frac{m}{n}})$ . To see this note that the ideal  $(Y^m - x^n) \triangleleft k[Y, \frac{1}{Y}]$  is a maximal ideal and closed under differentiation since  $(Y^m - x^n)' = \frac{n}{x}(Y^m - x^n)$ .*

*Case 2:  $\alpha \notin \mathbb{Q}$ . In this case, I claim that  $(0)$  is a maximal differential ideal and so  $R = k[Y, \frac{1}{Y}]$ . To see this, first note that for any  $f \in \mathbb{C}(x)$ ,*

$$\frac{f'}{f} = \sum \frac{n_i}{(x - \alpha_i)}, n_i \in \mathbb{Z}, \alpha_i \in \mathbb{C}$$

*and this can never equal  $\frac{N\alpha}{x}$ . Therefore  $Y' = \frac{N\alpha}{x}Y$  has no nonzero solutions in  $\mathbb{C}(x)$ . Now assume  $I$  is a proper nonzero differential ideal in  $K[Y, \frac{1}{Y}]$ . One sees that  $I$  is of the form  $I = (f)$  where  $f = Y^N + a_{N-1}Y^{N-1} + \dots + a_0$ ,  $N > 0$ . Since  $Y$  is invertible, we may assume that  $a_0 \neq 0$ . Since  $f' \in (f)$ , by comparing leading terms we have that  $f' = \frac{N\alpha}{x}f$  and so  $a'_0 = \frac{N\alpha}{x}a_0$ , a contradiction.*

2)  $k = \mathbb{C}(x)$   $x' = 1$   $Y' = Y$ . Let  $S = k[Y, \frac{1}{Y}]$ . An argument similar to Case 2 above shows that  $(0)$  is the only proper differential ideal so the PV-ring is  $k[Y, \frac{1}{Y}]$ .

As expected the differential Galois group is defined as

**Definition 1.3.4** Let  $(k, \partial)$  be a differential field and  $R$  a PV-ring over  $k$ . The differential Galois group of  $R$  over  $k$ ,  $\text{DGal}(R/k)$  is the group  $\{\sigma : R \rightarrow R \mid \sigma \text{ is a differential } k\text{-isomorphism}\}$ .

If  $R$  is the PV-ring associated with  $Y' = AY$ ,  $A \in \text{gl}_n(k)$ , we sometimes denote the differential Galois group by  $\text{DGal}(Y' = AY)$ . If  $K$  is a PV-field over  $k$ , then the  $k$ -differential automorphisms of  $K$  over  $k$  are can

be identified with the differential  $k$ -automorphisms of  $R$  over  $k$  where  $R$  is a PV-ring whose quotient field is  $K$ .

Let  $R$  be a PV-ring for the equation  $Y' = AY$  over  $k$  and  $\sigma \in \text{DGal}(R/k)$ . Fix a fundamental solution matrix  $Z$  and let  $\sigma \in \text{DGal}(R/k)$ . We then have that  $\sigma(Z)$  is again a fundamental solution matrix of  $Y' = AY$  and, a calculation shows that  $(Z^{-1}\sigma(Z))' = 0$ . Therefore,  $\sigma(Z) = Zm_\sigma$  for some  $m_\sigma \in \text{GL}_n(C_k)$ . This gives an injective group homomorphism  $\sigma \mapsto m_\sigma$  of  $\text{DGal}(R/k)$  into  $\text{GL}_n(C_\sigma)$ . If we select a different fundamental solution matrix, the images of the resulting maps would be conjugate. The image has a further property.

**Definition 1.3.5** *A subgroup  $G$  of  $\text{GL}_n(C_k)$  is said to be a linear algebraic group if it is a Zariski close the zero set in  $\text{GL}_n(C_k)$  of a system of polynomials over  $C_k$  in  $n^2$  variables.*

With the identification above  $\text{DGal}(R/k) \subset \text{GL}_n(C_k)$  we have

**Proposition 1.3.6**  *$\text{DGal}(R/k) \subset \text{GL}_n(C_k)$  is a linear algebraic group.*

*Proof* Let  $R = k[Y, \frac{1}{\det Y}]/M$ ,  $M = (f_1, \dots, f_m)$ . We may assume that the  $f_i \in k[Y_{1,1}, \dots, Y_{n,n}]$ . We may identify  $\text{DGal}(R/k)$  with the subgroup of  $\sigma \in \text{GL}_n(C_k)$  such that  $\sigma(M) \subset M$ . Let  $W_N$  be the  $C_k$ -space of polynomials in  $k[Y_{1,1}, \dots, Y_{n,n}]$  of degree at most  $N$ . Note that  $\text{GL}_n(C_k)$  acts on  $k[Y_{1,1}, \dots, Y_{n,n}]$  by substitution and leaves  $W_N$  invariant. If  $N$  is the maximum of the degrees of the  $f_i$  then  $V_N = W_N \cap M$  generates  $M$  and we may identify  $\text{DGal}(R/k)$  with the group of  $\sigma \in \text{GL}_n(C_k)$  such that  $\sigma(V_N) \subset V_N$ . Let  $\{e_\alpha\}_\mathcal{A}$  be a  $C$ -basis of  $W_N$  with  $\{e_\alpha\}_{\alpha \in \mathcal{B} \subset \mathcal{A}}$  a  $C$ -basis of  $V_N$ . For any  $\beta \in \mathcal{A}$  there exist polynomials  $p_{\alpha\beta}(x_{ij})$  such that for any  $\sigma = (c_{ij}) \in \text{GL}_n$ ,  $\sigma(e_\beta) = \sum_\alpha p_{\alpha\beta}(c_{ij})e_\alpha$ . We then have that  $\sigma = (c_{ij}) \in \text{DGal}(R/k)$  if and only if  $p_{\alpha\beta}(c_{ij}) = 0$  for all  $\beta \in \mathcal{B}$  and  $\alpha \notin \mathcal{B}$ .  $\square$

**Examples 1.3.7** 1)  $k = \mathbb{C}(x)$   $x' = 1$   $Y' = \frac{\alpha}{x}Y$   $\text{DGal} \subset \text{GL}_1(\mathbb{C})$ . From the above description of the differential Galois group as the group having a certain ideal invariant, we see (using the information of Example 1.3.3):

Case 1:  $\alpha = \frac{n}{m}, \text{GCD}(n, m) = 1$ . We then have  $R = \mathbb{C}[Y, \frac{1}{Y}]/(Y^m - x^n) = \mathbb{C}(x)[x^{\frac{n}{m}}]$ . Therefore  $\text{DGal} = \mathbb{Z}/m\mathbb{Z} = \{(c) \mid c^n - 1 = 0\} \subset \text{GL}_1(\mathbb{C})$

Case 2:  $\alpha \notin \mathbb{Q}$ . we have  $R = \mathbb{C}(x)[Y, \frac{1}{Y}]/(0)$ . Therefore  $\text{DGal} = \text{GL}_1(\mathbb{C})$ .

2)  $k = \mathbb{C}(x)$   $x' = 1$   $Y' = Y$ . As above, one sees that  $\text{DGal} = \text{GL}_1(\mathbb{C})$

In general, it is not easy to compute Galois groups (although there is a complete algorithm due to Hrushovski [Hru02]). We will give more examples in Sections 1.4, 1.5, and 1.6.

One can define the differential Galois group of a differential module in the following way. Let  $M$  be a differential module. Once we have selected a basis, we may assume that  $M \simeq M_A$ , where  $M_A$  is the differential module associated with  $Y' = AY$ . We say that  $Y' = AY$  is a matrix linear differential equation associated with  $M$ .

**Definition 1.3.8** *Let  $M$  be a differential module over  $k$ . The differential Galois group of  $M$  is the differential Galois group of an associated matrix linear differential equation.*

To show that this definition makes sense we must show (see the discussion before Definition 1.2.5) that equivalent matrix differential equations have the same differential Galois groups. Let  $Y' = A_1Y$ ,  $Y' = A_2Y$  be equivalent equations and let  $A_1 = B^{-1}A_2B - B^{-1}B'$ ,  $B \in \text{GL}_n(k)$ . If  $R = k[Z_1, \frac{1}{Z_2}]$  is a PV-ring for  $Y' = A_1Y$ , then a computation shows that  $Z_2 = BZ_1$  is a fundamental solution matrix for  $Y' = A_2Y$  and so  $R$  will also be a PV-ring for  $Y' = A_2Y$ . The equation  $Z_2 = BZ_1$  furthermore shows that the action of the differential Galois group (as a matrix group) on  $Z_1$  and  $Z_2$  is the same.

We now state

**Theorem 1.3.9 (Fundamental Theorem of Differential Galois Theory)** *Let  $k$  be a differential field with algebraically closed constants  $C_k$ . Let  $K$  be a Picard-Vessiot field with differential Galois group  $G$ .*

1) *There is a bijective correspondence between Zariski-closed subgroups  $H \subset G$  and differential subfields  $F$ ,  $k \subset F \subset K$  given by*

$$\begin{aligned} H \subset G &\mapsto K^H = \{a \in K \mid \sigma(a) = a \text{ for all } \sigma \in H\} \\ k \subset F \subset K &\mapsto \text{DGal}(K/F) = \{\sigma \in G \mid \sigma(a) = a \text{ for all } a \in F\} \end{aligned}$$

2) *A differential subfield  $k \subset F \subset K$  is a Picard-Vessiot extension of  $k$  if and only if  $\text{DGal}(K/F)$  is a normal subgroup of  $G$ , in which case  $\text{DGal}(F/k) \simeq \text{DGal}(K/k)\text{DGal}(K/F)$ .*

The Zariski closed sets form the collection of closed sets of a topology on  $\mathrm{GL}_n$  and one can speak of closures, components, etc. in this topology. In this topology, each linear algebraic group  $G$  may be written (uniquely) as the finite disjoint union of connected closed subsets (see Appendix A of [dPS97] or [Spr98] for a further discussion of linear algebraic groups). The connected component containing the identity matrix is denoted by  $G^0$  and is referred to as the identity component. It is a normal subgroup of  $G$ . As a consequence of the Fundamental Theorem, one can show the following:

**Corollary 1.3.10** *Let  $k, K$  and  $G$  be as above.*

- 1) *For  $a \in K$ , we have that  $a \in k \Leftrightarrow \sigma(a) = a \ \forall \sigma \in \mathrm{DGal}(K/k)$*
- 2) *For  $H \subset \mathrm{DGal}(K/k)$ , the Zariski closure  $\overline{H}$  is  $\mathrm{DGal}(K/k)$  if and only if  $K^H = k$*
- 3)  *$K^{G^0}$  is precisely the algebraic closure  $\tilde{k}$  of  $k$  in  $K$ .*

I refer the reader to Chapter 1.3 of [dPS03] for the proof of the Fundamental Theorem and its corollary.

**Example 1.3.11**  $y' = \frac{\alpha}{x}y \quad \alpha \notin \mathbb{Q}, \quad k = \mathbb{C}(x) \quad K = k(x^\alpha) \quad \mathrm{DGal} = \mathrm{GL}_1 = \mathbb{C}^*$

*The Zariski-closed subsets of  $\mathbb{C}^*$  are finite so the closed subgroups are finite and cyclic. We have the following correspondence*

<i>Groups</i>		<i>Fields</i>
$\{1\}$	$\Leftrightarrow$	$k(x^\alpha)$
$\cap$		$\cup$
$\{1, e^{2\pi i/n}, \dots, e^{2(n-1)\pi i/n}\}$	$\Leftrightarrow$	$k(x^{n\alpha})$
$\cap$		$\cup$
$\mathbb{C}^*$	$\Leftrightarrow$	$k$

The Fundamental Theorem also follows from a deeper fact giving geometric meaning to the Picard-Vessiot ring and relating this to the Galois group.

**Galois theory and torsors** Let  $k$  be a field (not necessarily a differential field) and  $\tilde{k}$  its algebraic closure. I wish to refine the notion of a Zariski closed set in order to keep track of the field over which these objects are defined.

**Definition 1.3.12** 1) A Zariski closed subset  $V \subset \bar{k}^n$  is called a variety defined over  $k$  or  $k$ -variety if it is the set of common zeroes of polynomials in  $k[X_1, \dots, X_n]$ .

2) If  $V$  is a variety defined over  $k$ , the  $k$ -ideal of  $V$ ,  $I_k(V)$ , is  $\{f \in k[X_1, \dots, X_n] \mid f(a) = 0 \text{ for all } a \in V\}$ .

3) If  $V$  is a variety defined over  $k$ , the  $k$ -coordinate ring,  $k[V]$ , of  $V$  is the ring  $k[X_1, \dots, X_n]/I_k(V)$ .

4) If  $V \subset \bar{k}^n$  and  $W \subset \bar{k}^n$ , a  $k$ -morphism  $f : V \rightarrow W$  is an  $n$ -tuple of polynomials  $f = (f_1, \dots, f_n)$  in  $k[X_1, \dots, X_n]$  such that  $f(a) \in W$  for all  $a \in V$ .

5) We say two  $k$ -varieties  $V$  and  $W$  are  $k$ -isomorphic if there are  $k$ -morphisms  $f : V \rightarrow W$  and  $g : W \rightarrow V$  such that  $fg$  and  $gf$  are the identity maps.

Note that the coordinate ring is isomorphic to the ring of  $k$ -morphisms from  $V$  to  $k$ . Also note that a  $k$ -morphism  $f : V \rightarrow W$  induces a  $k$ -algebra homomorphism  $f^* : k[W] \rightarrow k[V]$  given by  $f^*(g)(v) = g(f(v))$  for  $v \in V$  and  $g \in k[W]$ .

**Examples 1.3.13** 1) The affine spaces  $k^n$  are  $\mathbb{Q}$ -varieties and, as such, their coordinate rings are just the rings of polynomials  $\mathbb{Q}[X_1, \dots, X_n]$ .

2) The group of invertible matrices  $\text{GL}_n(\bar{k})$  is also a  $\mathbb{Q}$ -variety. In order to think of it as a closed set, we must embed each invertible matrix  $A$  in the  $(n+1) \times (n+1)$  matrix  $\begin{pmatrix} A & 0 \\ 0 & \det(A) \end{pmatrix}$ . The defining ideal in  $\mathbb{Q}[X_{1,1}, \dots, X_{n+1,n+1}]$  is generated by

$$X_{n+1,1}, \dots, X_{n+1,n}, X_{1,n+1}, \dots, X_{n,n+1}, (\det X) - X_{n+1,n+1}$$

where  $X$  is the  $n \times n$  matrix with entries  $X_{i,j}$ ,  $1 \leq i \leq n$ ,  $1 \leq j \leq n$ . The coordinate ring is the ring  $\mathbb{Q}[X_{1,1}, \dots, X_{n,n}, \frac{1}{\det X}]$ .

**Examples 1.3.14** 1) Let  $k = \mathbb{Q}$  and  $V = \{a \mid a^2 - 2 = 0\} \subset \bar{\mathbb{Q}}$ . We have that  $I_{\mathbb{Q}}(V) = (X^2 - 2) \subset \mathbb{Q}[X]$  and  $k[V] = \mathbb{Q}[X]/(X^2 - 2) = \mathbb{Q}(\sqrt{2})$ . Note that in this setting “ $\sqrt{2}$ ” is a function on  $V$ , defined over  $\mathbb{Q}$  with values in  $\bar{\mathbb{Q}}$  and not a number!

2) Let  $k = \mathbb{Q}$  and  $W = \{a \mid a^2 - 1 = 0\} = \{\pm 1\} \subset \bar{\mathbb{Q}}$ . We have that  $I_{\mathbb{Q}}(W) = (X^2 - 1)$  and  $k[W] = \mathbb{Q}[X]/(X^2 - 1) = \mathbb{Q} \oplus \mathbb{Q}$ .

3) Note that  $V$  and  $W$  are not  $\mathbb{Q}$ -isomorphic since such an isomorphism

would induce an isomorphism of their  $\mathbb{Q}$ -coordinate rings. If we consider  $V$  and  $W$  as  $\mathbb{Q}$ -varieties, then the map  $f(x) = \sqrt{2}(x-1) - \sqrt{2}(x+1)$  gives a  $\mathbb{Q}$ -isomorphism of  $W$  to  $V$ .

We shall be interested in certain  $k$ -subvarieties of  $\mathrm{GL}_n(\bar{k})$  that have a group acting on them. These are described in the following definition (a more general definition of torsor is given in Appendix A of [dPS03] but the definition below suffices for our present needs).

**Definition 1.3.15** *Let  $G \subset \mathrm{GL}_n(\bar{k})$  be a linear algebraic group defined over  $k$  and  $V \subset \mathrm{GL}_n(\bar{k})$  a  $k$ -variety. We say that  $V$  is a  $G$ -torsor (or  $G$ -principal homogeneous space) over  $k$  if for all  $v, w \in V$  there exists a unique  $g \in G$  such that  $vg = w$ . Two  $G$ -torsors  $V, W \in \mathrm{GL}_n(\bar{k})$  are said to be  $k$ -isomorphic if there is a  $k$ -isomorphism between the two commuting with the action of  $G$ .*

**Examples 1.3.16** 1)  $G$  itself is always a  $G$ -torsor.

2) Let  $k = \mathbb{Q}$  and  $V = \{a \mid a^2 - 2 = 0\} \subset \bar{\mathbb{Q}}$ . We have that  $V \subset \mathbb{Q}^* = \mathrm{GL}_1(\mathbb{Q})$ . Let  $G = \{\pm 1\} \subset \mathrm{GL}_1(\mathbb{Q})$ . It is easy to see that  $V$  is a  $G$ -torsor. As noted in the previous examples, the torsors  $V$  and  $G$  are not isomorphic over  $\mathbb{Q}$  but are isomorphic over  $\bar{\mathbb{Q}}$ .

We say a  $G$ -torsor over  $k$  is *trivial* if it is  $k$ -isomorphic to  $G$  itself. One has the following criterion.

**Lemma 1.3.17** *A  $G$ -torsor  $V$  over  $k$  is trivial if and only if it contains a point with coordinates in  $k$ . In particular, any  $G$ -torsor over an algebraically closed field is trivial.*

*Proof* If  $V$  contains a point  $v$  with coordinates in  $k$  then the map  $g \mapsto xg$  is defined over  $k$  and gives a  $k$ -isomorphism between  $G$  and  $V$  (with inverse  $v \mapsto x^{-1}v$ ). If  $V$  is trivial and  $f : G \rightarrow V$  is an isomorphism defined over  $k$ , then  $f(id)$  has coordinates in  $k$  and so is a  $k$ -point of  $V$ .  $\square$

For connected groups  $G$ , there are other fields for which all torsors are trivial.

**Proposition 1.3.18** *Let  $G$  be a connected linear algebraic group defined over  $k$  and  $V$  a  $G$ -torsor over  $k$ . If  $k$  is an algebraic extension of  $\mathbb{C}(x)$ ,  $\mathbb{C}((x))$ , or  $\mathbb{C}(\{x\})$ , then  $V$  is a trivial  $G$ -torsor.*



*Proof* This result is due to Steinberg, see [Ser64], Chapter III.2.3  $\square$

Another way of saying that  $V$  is a  $G$  torsor is to say that  $V$  is defined over  $k$  and is a left coset of  $G$  in  $\mathrm{GL}_n(\bar{k})$ . Because of this one would expect that  $V$  and  $G$  share many geometric properties. The one we will be interested in concerns the dimension. We say that a  $k$ -variety  $V$  is *irreducible* if it is not the union of two proper  $k$ -varieties. This is equivalent to saying that  $I_k(V)$  is a prime ideal or, equivalently, that  $k[V]$  is an integral domain. For an irreducible  $k$ -variety  $V$ , one defines the *dimension of  $V$* ,  $\dim_k(V)$  to be the transcendence degree of  $k(V)$  over  $k$ , where  $k(V)$  is the quotient field of  $k[V]$ . For example, if  $k = \mathbb{C}$  and  $V$  is a  $k$ -variety that is nonsingular as a complex manifold, then the dimension in the above sense is just the dimension as a complex manifold (this happens for example when  $V$  is a subgroup of  $\mathrm{GL}_n(\mathbb{C})$ ). In general, any  $k$ -variety  $V$  can be written as the irredundant union of irreducible  $k$ -varieties, called the irreducible components of  $V$  and the dimension of  $V$  is defined to be the maximum of the dimensions of its irreducible components. One can show that if  $k \subset K$ , then  $\dim_k(V) = \dim_K(V)$  and in particular,  $\dim_k(V) = \dim_{\bar{k}} V$  (see [dPS03], Appendix A for more information about  $k$ -varieties).

**Proposition 1.3.19** *If  $V$  is a  $G$ -torsor, then  $\dim_k(V) = \dim_k(G)$ .*

*Proof* As noted above, any  $G$ -torsor over  $k$  is trivial when considered as a  $\bar{k}$  torsor. We therefore have  $\dim_k(V) = \dim_{\bar{k}}(V) = \dim_{\bar{k}}(G) = \dim_k(G)$ .  $\square$

The reason for introducing the concept of a  $G$ -torsor in this paper is the following proposition. We use the fact that if  $V$  is a  $G$ -torsor over  $k$  and  $g \in G$ , then the isomorphism  $\rho_g : v \mapsto vg$  induces an isomorphism  $\rho_g^* : k[V] \rightarrow k[V]$ .

**Proposition 1.3.20** *Let  $R$  be a Picard-Vessiot extension with differential Galois group  $G$ . Then  $R$  is the coordinate ring of a  $G$ -torsor  $V$  defined over  $k$ . Furthermore, for  $\sigma \in G$ , the Galois action of  $\sigma$  on  $R = k[V]$  is the same as  $\rho_\sigma^*$ .*

**Corollary 1.3.21** *1) If  $K$  is a Picard-Vessiot field over  $k$  with Galois group  $G$ , then  $\dim_{C_k} G = \mathrm{tr.deg.}(K/k)$ .*

*2) If  $R$  is a Picard-Vessiot ring over  $k$  with Galois group  $G$  where  $k$  is either algebraically closed,  $\mathbb{C}(x)$ ,  $\mathbb{C}((x))$ , or  $\mathbb{C}(\{x\})$ , then  $R \simeq k[G]$ .*

We refer to Chapter 1.3 of [dPS03] for a proof of the proposition and note that the Fundamental Theorem of Galois Theory is shown there to follow from this proposition.

**Applications and ramifications.** I shall discuss the following applications and ramifications of the differential Galois theory

- Monodromy
- Factorization of linear differential operators
- Transcendental numbers
- Why doesn't  $\sec x$  satisfy a linear differential equation?
- Systems of linear partial differential equations

Another application concerning solving linear differential equations in terms of special functions and algorithms to do this will be discussed in Section 1.5. The Picard-Vessiot theory has also been used to identify obstructions to complete integrability of Hamiltonian systems (see [Aud01] or [MR99])

Monodromy. We will now consider differential equations over the complex numbers, so let  $k = \mathbb{C}(x)$ ,  $x' = 1$  and

$$\frac{dY}{dx} = A(x)Y, \quad A(x) \in \mathfrak{gl}_n(\mathbb{C}(x)) \quad (1.1)$$

**Definition 1.3.22** *A point  $x_0 \in \mathbb{C}$  is an ordinary point of equation (1.1) if all the entries of  $A(x)$  are analytic at  $x_0$ , otherwise  $x_0$  is a singular point.*

One can also consider the point at infinity,  $\infty$ . Let  $t = \frac{1}{x}$ . We then have that  $\frac{dY}{dt} = -\frac{1}{t^2}A(\frac{1}{t})Y$ . The point  $x = \infty$  is an ordinary or singular point precisely when  $t = 0$  is an ordinary or singular point of the transformed equation.

**Example 1.3.23**  $\frac{dy}{dx} = \frac{\alpha}{x}y \Rightarrow \frac{dy}{dt} = -\frac{\alpha}{t}y$  so the singular points are  $\{0, \infty\}$ .

Let  $S = \mathbb{CP}^1$  be the Riemann Sphere and let  $X = S^2 - \{\text{singular points of (1.1)}\}$  and let  $x_0 \in X$ . From standard existence theorems, we know that in a small neighborhood  $\mathcal{O}$  of  $x_0$ , there exists a fundamental matrix  $Z$  whose entries are analytic in  $\mathcal{O}$ . Let  $\gamma$  be a closed curve in  $X$  based at  $x_0$ .

Analytic continuation along  $\gamma$  yields a new fundamental solution matrix  $Z_\gamma$  at  $x_0$  which must be related to the old fundamental matrix as

$$Z_\gamma = ZD_\gamma$$

where  $D_\gamma \in \text{GL}_n(\mathbb{C})$ . One can show that  $D_\gamma$  just depends on the homotopy class of  $\gamma$ . This defines a homomorphism

$$\begin{aligned} \text{Mon} : \pi_1(X, x_0) &\rightarrow \text{GL}_n(\mathbb{C}) \\ \gamma &\mapsto D_\gamma \end{aligned}$$

**Definition 1.3.24** *The homomorphism Mon is called the monodromy map and its image is called the monodromy group.*

Note that the monodromy group depends on the choice of  $Z$  so it is only determined up to conjugation. Since analytic continuation preserves analytic relations, we have that the monodromy group is contained in the differential Galois group.

**Examples 1.3.25** 1)  $y' = \frac{\alpha}{x}y$  ( $y = e^{\alpha \log x} = x^\alpha$ ) Singular points =  $\{0, \infty\}$

$$\text{Mon}(\pi_1(X, x_0)) = \{(e^{2\pi i \alpha})^n \mid n \in \mathbb{Z}\}$$

*If  $\alpha \in \mathbb{Q}$  this image is finite and so equals the differential Galois group. If  $\alpha \notin \mathbb{Q}$ , then this image is infinite and so Zariski dense in  $\text{GL}_1(\mathbb{C})$ . In either case the monodromy group is Zariski dense in the differential Galois group.*

2)  $y' = y$  ( $y = e^x$ ), Singular points =  $\{\infty\}$

$$\text{Mon}(\pi_1(X, x_0)) = \{1\}$$

*In this example the differential Galois group is  $\text{GL}_1(\mathbb{C})$  and the monodromy group is not Zariski dense in this group.*

The second example shows that the monodromy group is not necessarily Zariski dense in the differential Galois group but the first example suggests that under certain conditions it may be. To state these conditions we need the following definitions.

**Definition 1.3.26** 1) *An open sector  $\mathcal{S} = \mathcal{S}(a, b, \rho)$  at a point  $p$  is the set of complex numbers  $z \neq p$  such that  $\arg(z - p) \in (a, b)$  and  $|z - p| < \rho(\arg(z - p))$  where  $\rho : (a, b) \rightarrow \mathbb{R}_{>0}$  is a continuous function.*

2) *A singular point  $p$  of  $Y' = AY$  is a regular singular point if for any*

open sector  $\mathcal{S}(a, b, \rho)$  at  $p$  with  $|a - b| < 2\pi$ , there exists a fundamental solution matrix  $Z = (z_{ij})$ , analytic in  $\mathcal{S}$  such that for some positive constant  $c$  and integer  $N$ ,  $|z_{ij}| < c|x^N|$  as  $x \rightarrow p$  in  $\mathcal{S}$ .

Note that in the first example above, the singular points are regular while in the second they are not. For scalar linear differential equations  $L(y) = y^{(n)} + a_{n-1}y^{(n-1)} + \dots + a_0y = 0$ , the criterion of Fuchs states that  $p$  is a regular singular point if and only if each  $a_i$  has a pole of order  $\leq n - i$  at  $p$  (see Chapter 5.1 of [dPS03] for a further discussion of regular singular points). For equations with only regular singular points we have

**Proposition 1.3.27** (*Schlesinger*) *If  $Y' = AY$  has only regular singular points, then  $\text{Mon}(\pi_1(X, x_0))$  is Zariski dense in  $D\text{Gal}$ .*

*Proof* Let  $K = \text{PV}$  extension field of  $k$  for  $Y' = AY$ . We have  $\text{Mon}(\pi_1(X, x_0)) \subset D\text{Gal}(K/k)$ . By Corollary 1.3.10 it is enough to show that if  $f \in K$  is left fixed by  $\text{Mon}(\pi_1(X, x_0))$  then  $f \in k = \mathbb{C}(x)$ . If  $f$  is left invariant by analytic continuation on  $X$ , then  $f$  is a single valued meromorphic function on  $X$ . Let  $Z = (z_{ij})$ ,  $z_{ij} \in K$  be a fundamental solution matrix and let  $f \in \mathbb{C}(x, z_{ij})$ . Regular singular points imply that  $f$  has polynomial growth near the singular points of  $Y' = AY$  and is at worst meromorphic at other points on the Riemann Sphere. Cauchy's Theorem implies that  $f$  is meromorphic on the Riemann Sphere and Liouville's Theorem implies that  $f$  must be a rational function.  $\square$

In Section 1.4, we shall see what other analytic information one needs to determine the differential Galois group in general.

Factorization of linear differential operators. In Section 1.5, we will see that factorization properties of linear differential operators and differential modules play a crucial role in algorithms to calculate properties of Galois groups. Here I shall relate factorization of operators to the Galois theory.

Let  $k$  be a differential field,  $k[\partial]$  the ring of differential operators over  $k$  and  $L \in k[\partial]$ . Let  $K$  be the PV-field over  $k$  corresponding to  $L(y) = 0$ ,  $G$  its Galois group and  $V = \text{Soln}_K(L(y) = 0)$ .

**Proposition 1.3.28** *There is a correspondence among the following objects*

- (i) *Monic  $L_0 \in k[\partial]$  such that  $L = L_1 \cdot L_0$ .*
- (ii) *Differential submodules of  $k[\partial]/k[\partial]L$ .*
- (iii)  *$G$ -invariant subspaces of  $V$ .*

*Proof* Let  $W \subset V$  be  $G$ -invariant and  $B = \{b_1, \dots, b_t\}$  a  $C_k$ -basis of  $W$ . For  $\sigma \in G$ , let  $[\sigma]_B$  be the matrix of  $\sigma|_W$  with respect to  $B$ ,

$$\sigma\left(\begin{bmatrix} b_1 & \dots & b_t \\ \vdots & \vdots & \vdots \\ b_1^{(t-1)} & \dots & b_t^{(t-1)} \end{bmatrix}\right) = \begin{bmatrix} b_1 & \dots & b_t \\ \vdots & \vdots & \vdots \\ b_1^{(t-1)} & \dots & b_t^{(t-1)} \end{bmatrix} [\sigma]_B.$$

Taking determinants we have  $\sigma(wr(b_1, \dots, b_t)) = wr(b_1, \dots, b_t) \det([\sigma]_B)$ . Define  $L_W(Y) \stackrel{\text{def}}{=} \frac{wr(Y, b_1, \dots, b_t)}{wr(b_1, \dots, b_t)}$ . Using the above, we have that the coefficients of  $L_W$  are left fixed by the differential Galois group and so lie in  $k$ . Therefore  $L_W \in k[\partial]$ . Furthermore, Corollary 1.2.12 implies that  $L_W$  vanishes on  $W$ . If we write  $L = L_1 \cdot L_W + R$  where  $\text{ord}(R) < \text{ord}(L_W)$ , one sees that  $R$  also vanishes on  $W$ . Since the dimension of  $W = \text{ord}(L_W)$ , Corollary 1.2.12 again implies that  $R = 0$ . Therefore  $W$  is the solution space of the right factor  $L_W$  of  $L$ .

Let  $L_0 \in k[\partial]$  with  $L = L_1 \cdot L_0$ . Let  $W = \text{Soln}_K(L_0(y) = 0)$ . We have that  $L_0$  maps  $V$  into  $\text{Soln}_K(L_1(y) = 0)$ . Furthermore we have the following implications

- $\text{Im}(L_0) \subset \text{Soln}_K(L_1(y) = 0) \Rightarrow \dim_{C_k}(\text{Im}(L_0)) \leq \text{ord}(L_1)$
- $\text{Ker}(L_0) = \text{Soln}_K(L_0(y) = 0) \Rightarrow \dim_{C_k}(\text{Ker}(L_0)) \leq \text{ord}(L_0)$
- $n = \dim_{C_k}(V) = \dim_{C_k}(\text{Im}(L_0)) + \dim_{C_k}(\text{Ker}(L_0)) \leq \text{ord}(L_1) + \text{ord}(L_0) = n$

Therefore  $\dim_{C_k} \text{Soln}_K(L_0(y) = 0) = \text{ord}(L_0)$ . This now implies that the association of  $G$ -invariant subspaces of  $V$  to right factors of  $L$  is a bijection.

I will now describe the correspondence between right factors of  $L$  and submodules of  $k[\partial]/k[\partial]L$ . One easily checks that if  $L = L_1 L_0$ , then  $k[\partial]L \subset k[\partial]L_0$  so  $M = k[\partial]L_0/k[\partial]L \subset k[\partial]/k[\partial]L$ . Conversely let  $M$  be a differential submodule of  $k[\partial]/k[\partial]L$ . Let  $\pi : k[\partial] \rightarrow k[\partial]/k[\partial]L$  and  $\overline{M} = \pi^{-1}(M)$ .  $\overline{M} = k[\partial]L_0$  for some  $L_0 \in k[\partial]$ . Since  $L \in \overline{M}$ ,  $L = L_1 L_0$ .  $\square$

By induction on the order, one can show that any differential operator can be written as a product of irreducible operators. This need not be unique (for example  $\partial^2 = \partial \cdot \partial = (\partial + \frac{1}{x})(\partial - \frac{1}{x})$ ) but a version of the Jordan-Hölder Theorem implies that if  $L = L_1 \cdot \dots \cdot L_r = \tilde{L}_1 \cdot \dots \cdot \tilde{L}_s$  the  $r = s$  and, after possibly renumbering,  $k[\partial]/k[\partial]L_i \simeq k[\partial]/k[\partial]\tilde{L}_i$  ([dPS03], Chapter 2).

One can also characterize submodules of differential modules in terms of corresponding matrix linear differential equations. Let  $Y' = AY$  correspond to the differential module  $M$  of dimension  $n$ . One then has that  $M$  has a proper submodule of dimension  $t$  if and only if  $Y' = AY$  is equivalent to  $Y' = BY$ ,  $B = \begin{pmatrix} B_0 & B_1 \\ 0 & B_2 \end{pmatrix}$   $B_0 \in \mathfrak{gl}_t(k)$ .

Transcendental numbers. Results of Siegel and Shidlovski (and subsequent authors) reduce the problem of showing that the values of certain functions are algebraically independent to showing that these functions are themselves algebraically independent. The Galois theory of linear differential equations can be used to do this.

**Definition 1.3.29**  $f = \sum_{i=0}^{\infty} a_n \frac{z^n}{n!} \in \mathbb{Q}[[z]]$  is an E-function if  $\exists c \in \mathbb{Q}$  such that, for all  $n$ ,  $|a_n| \leq c^n$  and the least common denominator of  $a_0, \dots, a_n \leq c^n$ .

**Examples 1.3.30**  $e^x = \sum \frac{z^n}{n!}$ ,  $J_0(z) = \sum \frac{z^{2n}}{(n!)^2}$

**Theorem 1.3.31** (Siegel-Shidlovski, c.f., [Lan66]): If  $f_1, \dots, f_s$  are E-functions, algebraically independent over  $\mathbb{Q}(z)$  and satisfy a linear differential equation

$$L(y) = y^{(m)} + a_{m-1}y^{(n-1)} + \dots + a_0y, \quad a_i \in \mathbb{Q}(z)$$

then for  $\alpha \in \mathbb{Q}$ ,  $\alpha \neq 0$  and  $\alpha$  not a pole of the  $a_i$ , the numbers  $f_1(\alpha), \dots, f_s(\alpha)$  are algebraically independent over  $\mathbb{Q}$ .

The differential Galois group of  $L(y) = 0$  measures the algebraic relations among solutions. In particular, Corollary 1.3.21 implies that if  $y_1, \dots, y_n$  are a  $\mathbb{Q}$ -basis of the solution space of  $L(y) = 0$  then the transcendence degree of  $\mathbb{Q}(x)(y_1, \dots, y_n, y'_1, \dots, y'_n, \dots, y_1^{(n-1)}, \dots, y_n^{(n-1)})$  over  $\mathbb{Q}(x)$  is equal to the dimension of the differential Galois group of  $L(y) = 0$ .

**Example 1.3.32** *The differential Galois group of the Bessel equations  $y'' + \frac{1}{z}y' + (1 - \frac{\lambda^2}{z^2})y = 0$ ,  $\lambda - \frac{1}{2} \notin \mathbb{Z}$  over  $\mathbb{Q}(x)$  can be shown to be  $\mathrm{SL}_2$  ([Kol68]) and examining the recurrence defining a power series solution of this equation, one sees that the solutions are  $E$ -functions. Given a fundamental set of solutions  $\{J_\lambda, Y_\lambda\}$  we have that the transcendence degree of  $\bar{\mathbb{Q}}(x)(J_\lambda, Y_\lambda, J'_\lambda, Y'_\lambda)$  over  $\bar{\mathbb{Q}}(x)$  is 3 (one can show that  $J_\lambda Y'_\lambda - J'_\lambda Y_\lambda \in \bar{\mathbb{Q}}(x)$ ). Therefore one can apply the Siegel-Shidlovski Theorem to show that certain values of  $J_\lambda, Y_\lambda$  and  $J'_\lambda$  are algebraically independent over  $\mathbb{Q}$ .*

Why does  $\cos x$  satisfy a linear differential equation over  $\mathbb{C}(x)$ , while  $\sec x$  does not? There are many reasons for this. For example,  $\sec x = \frac{1}{\cos x}$  has an infinite number of poles which is impossible for a solution of a linear differential equation over  $\mathbb{C}(x)$ . The following result, originally due to Harris and Sibuya [HS85], also yields this result and can be proven using differential Galois theory [Sin86].

**Proposition 1.3.33** *Let  $k$  be a differential field with algebraically closed constants and  $K$  a PV-field of  $k$ . Assume  $y \neq 0, 1/y \in K$  satisfy linear differential equations over  $k$ . The  $\frac{y'}{y}$  is algebraic over  $k$ .*

*Proof (Outline)* We may assume that  $k$  is algebraically closed and that  $K$  is the quotient field of a PV-ring  $R$ . Corollary 1.3.21 implies that  $R = k[G]$  where  $G$  is an algebraic group whose  $C_k$  points form the Galois group of  $R$  over  $k$ . Since both  $y$  and  $\frac{1}{y}$  satisfy linear differential equations over  $k$ , their orbits under the action of the Galois group span a finite dimensional vector space over  $C_k$ . This in turn implies the orbit of that  $y$  and  $\frac{1}{y}$  induced by the action of  $G(k)$  on  $G(k)$  by right multiplication is finite dimensional. The general theory of linear algebraic groups [Spr98] tells us that this implies that these elements must lie in  $k[G]$ . A result of Rosenlicht [Mag76, Ros61] implies that  $y = af$  where  $a \in k$  and  $f \in k[G]$  satisfies  $f(gh) = f(g)f(h)$  for all  $g, h \in G$ . In particular, for  $\sigma \in G(C_k) = \mathrm{Aut}(K/k)$ , we have  $\sigma(f(x)) = f(x\sigma) = f(x)f(\sigma)$ . If  $L \in k[\partial]$  is of minimal positive order such that  $L(f) = 0$ , then  $0 = \sigma(L(f)) = L(f \cdot f(\sigma))$  and a calculation shows that this implies that  $f(\sigma) \in C_k$ .

We therefore have that, for all  $\sigma \in \mathrm{Aut}(K/k)$  there exists a  $c_\sigma \in C_k$  such that  $\sigma(y) = c_\sigma y$ . This implies that  $\frac{y'}{y}$  is left fixed by  $\mathrm{Aut}(K/k)$  and so must be in  $k$ .  $\square$

If  $y = \cos x$  and  $\frac{1}{y} = \sec x$  both satisfy linear differential equations over  $\mathbb{Q}(x)$ , then the above result implies that  $\tan x$  would be algebraic over  $\mathbb{Q}(x)$ . One can show that this is not the case by noting that a periodic function that is algebraic over  $\mathbb{Q}(x)$  must be constant, a contradiction. Another proof can be given by starting with a putative minimal polynomial  $P(Y) \in \mathbb{C}(x)[Y]$  and noting that  $\frac{dP}{dx} + \frac{dP}{dY}Y' = \frac{dP}{dx} + \frac{dP}{dY}(Y^2 + 1)$  must be divisible by  $P$ . Comparing coefficients, one sees that the coefficients of powers of  $Y$  must be constant and so  $\tan x$  would be constant.

*Systems of linear partial differential equations.* Let  $(k, \Delta = \{\partial_1, \dots, \partial_m\})$  be a partial differential field. One can define the ring of linear partial differential operators  $k[\partial_1, \dots, \partial_m]$  as the noncommutative polynomial ring in  $\partial_1, \dots, \partial_m$  where  $\partial_i \cdot a = \partial_i(a) + a\partial_i$  for all  $a \in k$ . In this context, a differential module is a finite dimensional  $k$ -space that is a left  $k[\partial_1, \dots, \partial_m]$ -module. We have

**Proposition 1.3.34** *Assume  $k$  contains an element  $a$  such that  $\partial_i(a) \neq 0$  for some  $i$ . There is a correspondence between*

- *differential modules,*
- *left ideals  $I \subset k[\partial_1, \dots, \partial_m]$  such that  $\dim_k k[\partial_1, \dots, \partial_m]/I < \infty$ ,*  
*and*
- *systems  $\partial_i Y = A_i Y, A \in \mathfrak{gl}_n(k), i = 1, \dots, m$  such that*

$$\partial_i(A_j) + A_i A_j = \partial_j(A_i) + A_j A_i$$

*i.e., integrable systems*

If

$$\partial_i Y = A_i Y, A \in \mathfrak{gl}_n(k), i = 1, \dots, m$$

is an integrable system, the solution space over any differential field is finite dimensional over constants. Furthermore, one can define PV extensions, differential Galois groups, etc. for such systems and develop a Galois theory whose groups are linear algebraic groups. For further details see [Kol76, CS06] and Appendix D of [dPS03].

## 1.4 Local Galois Theory

In the previous section, we showed that for equations with only regular singular points the analytically defined monodromy group determines the differential Galois group. In this section we shall give an idea of what



analytic information determines the Galois group for general singular points. We shall use the following notation:

- $\mathbb{C}[[x]] = \{\sum_{i=0}^{\infty} a_i x^i \mid a_i \in \mathbb{C}\}, \quad x' = 1$
- $\mathbb{C}((x)) = \text{quotient field of } \mathbb{C}[[x]] = \mathbb{C}[[x]][x^{-1}]$
- $\mathbb{C}\{\{x\}\} = \{f \in \mathbb{C}[[x]] \mid f \text{ converges in a neighborhood } \mathcal{O}_f \text{ of } 0\}$
- $\mathbb{C}(\{x\}) = \text{quotient field of } \mathbb{C}\{\{x\}\} = \mathbb{C}\{\{x\}\}[x^{-1}]$

Note that

$$\mathbb{C}(x) \subset \mathbb{C}(\{x\}) \subset \mathbb{C}((x))$$

We shall consider equations of the form

$$Y' = AY \tag{1.2}$$

with  $A \in \text{gl}_n(\mathbb{C}(\{x\}))$  or  $\text{gl}_n(\mathbb{C}((x)))$  and discuss the form of solutions of such equations as well as the Galois groups relative to the base fields  $\mathbb{C}(\{x\})$  and  $\mathbb{C}((x))$ . We start with the simplest case:

**Ordinary points.** We say that  $x = 0$  is an *ordinary point* of equation (1.2) if  $A \in \text{gl}_n(\mathbb{C}[[x]])$ . The standard existence theory (see [Poo60] for a classical introduction or [Har64]) yields the following:

**Proposition 1.4.1** *If  $A \in \text{gl}_n(\mathbb{C}[[x]])$  then there exists a unique  $Y \in \text{GL}_n(\mathbb{C}[[x]])$  such that  $Y' = AY$  and  $Y(0)$  is the identity matrix. If  $A \in \text{gl}_n(\mathbb{C}(\{x\}))$  then we furthermore have that  $Y \in \text{GL}_n(\mathbb{C}(\{x\}))$ .*

This implies that for ordinary points with  $A \in \text{gl}_n(k)$  where  $k = \mathbb{C}(\{x\})$  or  $\mathbb{C}((x))$ , the Galois group of the associated Picard-Vessiot extension over  $k$  is trivial. The next case to consider is

**Regular singular points.** From now on we shall consider equations of the the form (1.2) with  $A \in \mathbb{C}(\{x\})$  (although some of what we say can be adapted to the case when  $A \in \mathbb{C}((x))$ ). We defined the notion of a regular singular point in Definition 1.3.26. The following proposition gives equivalent conditions (see Chapter 5.1.1 of [dPS03] for a proof of the equivalences).

**Proposition 1.4.2** *Assume  $A \in \text{gl}_n(\mathbb{C}(\{x\}))$ . The following are equivalent:*

- (i)  $Y' = AY$  is equivalent over  $\mathbb{C}((x))$  to  $Y' = \frac{A_0}{x}Y, A_0 \in \text{gl}_n(\mathbb{C})$ .
- (ii)  $Y' = AY$  is equivalent over  $\mathbb{C}(\{x\})$  to  $Y' = \frac{A_0}{x}Y, A_0 \in \text{gl}_n(\mathbb{C})$ .

- (iii) For any small sector  $\mathcal{S}(a, b, \rho)$ ,  $|a-b| < 2\pi$ , at 0,  $\exists$  a fundamental solution matrix  $Z = (z_{ij})$ , analytic in  $\mathcal{S}$  such that  $|z_{ij}| < |x|^N$  for some integer  $N$  as  $x \rightarrow 0$  (regular growth of solutions).
- (iv)  $Y' = AY$  has a fundamental solution matrix of the form  $Y = x^{A_0} Z$ ,  $A_0 \in \mathfrak{gl}_n(\mathbb{C})$ ,  $Z \in \mathrm{GL}_n(\mathbb{C}(\{x\}))$ .
- (v)  $Y' = AY$  is equivalent to  $y^{(n)} + a_{n-1}y^{(n-1)} + \dots + a_0y$ ,  $a_i \in \mathbb{C}(\{x\})$ , where  $\mathrm{ord}_{x=0} a_i \geq i - n$ .
- (vi)  $Y' = AY$  is equivalent to  $y^{(n)} + a_{n-1}y^{(n-1)} + \dots + a_0y$ ,  $a_i \in \mathbb{C}(\{x\})$ , having  $n$  linearly independent solutions of the form  $x^\alpha \sum_{i=0}^r c_i(x)(\log x)^i$ ,  $\alpha \in \mathbb{C}$ ,  $c_i(x) \in \mathbb{C}\{\{x\}\}$ .

We now turn to the Galois groups. We first consider the Galois group of a regular singular equation over  $\mathbb{C}(\{x\})$ . Let  $\mathcal{O}$  be a connected open neighborhood of 0 such that the entries of  $A$  are regular in  $\mathcal{O}$  with at worst a pole at 0. Let  $x_0 \neq 0$  be a point of  $\mathcal{O}$  and  $\gamma$  a simple loop around 0 in  $\mathcal{O}$  based at  $x_0$ . There exists a fundamental solution matrix  $Z$  at  $x_0$  and the field  $K = k(Z, \frac{1}{\det Z})$ ,  $k = \mathbb{C}(\{x\})$  is a Picard-Vessiot extension of  $k$ . Analytic continuation around  $\gamma$  gives a new fundamental solution matrix  $Z_\gamma = ZM_\gamma$  for some  $M_\gamma \in \mathrm{GL}_n(\mathbb{C})$ . The matrix  $M_\gamma$  depends on the choice of  $Z$  and so is determined only up to conjugation. Nonetheless, it is referred to as the *monodromy matrix at 0*. As in our discussion of monodromy in Section 1.5, we see that  $M_\gamma$  is in the differential Galois group  $\mathrm{DGal}(K/k)$  and, arguing as in Proposition 1.3.27, one sees that  $M_\gamma$  generates a cyclic group that is Zariski dense in  $\mathrm{DGal}(K/k)$ . From the theory of linear algebraic groups, one can show that a linear algebraic group have a Zariski dense cyclic subgroup must be of the form  $(\mathbb{C}^*, \times)^r \times (\mathbb{C}, +)^s \times \mathbb{Z}/m\mathbb{Z}$ ,  $r \geq 0$ ,  $s \in \{0, 1\}$  and that any such group occurs as a differential Galois group of a regular singular equation over  $k$ . (Exercise 5.3 of [dPS03]). This is not too surprising since property (vi) of the above proposition shows that  $K$  is contained in a field of the form  $k(x^{\alpha_1}, \dots, x^{\alpha_r}, \log x)$  for some  $\alpha_i \in \mathbb{C}$ .

Since  $\mathbb{C}(\{x\}) \subset \mathbb{C}((x)) = \tilde{k}$ , we can consider the the field  $\tilde{K} = \tilde{k}(Z, \frac{1}{\det Z})$  where  $Z$  is as above. Again, using (vi) above, we have that  $\tilde{K} \subset F = \tilde{k}(x^{\alpha_1}, \dots, x^{\alpha_r}, \log x)$  for some  $\alpha_i \in \mathbb{C}$ . Since we are no longer dealing with analytic functions, we cannot appeal to analytic continuation but we can define a *formal* monodromy, that is a differential automorphism of  $F$  over  $\tilde{k}$  that sends  $x^\alpha \mapsto x^\alpha e^{2\pi i \alpha}$  and  $\log x \mapsto \log x + 2\pi i$ . One can show that the only elements of  $F$  that are left fixed by this automorphism lie in  $\tilde{k}$ . This automorphism restricts to an automorphism of  $\tilde{K}$ . Therefore the Fundamental Theorem implies that the group gener-

ated by the formal monodromy is Zariski dense in the differential Galois group  $\mathrm{DGal}(\tilde{K}/\tilde{k})$ . The restriction of the formal monodromy to  $K$  (as above) is just the usual monodromy, so one sees that the *convergent Galois group*  $\mathrm{DGal}(K/k)$  equals the *formal Galois group*  $\mathrm{DGal}(\tilde{K}/\tilde{k})$ .

We note that the existence of  $n$  linearly independent solutions of the form described in (vi) goes back to Fuchs in the 19<sup>th</sup> Century and can be constructed using the Frobenius method (see [Gra00] for an historical account of the development of linear differential equations in the 19<sup>th</sup> Century).

**Irregular singular points.** We say  $x = 0$  is an irregular singular point if it is not ordinary or regular. Fabry, in the 19<sup>th</sup> Century constructed a fundamental set of formal solutions of a scalar linear differential equation at an irregular singular point. In terms of linear systems, every equation of the form  $Y' = AY$ ,  $A \in \mathrm{gl}_n(\mathbb{C}((x)))$  has a fundamental solution matrix of the form

$$\hat{Y} = \hat{\phi}(t)x^L e^{Q(1/t)}$$

where

- $t^\nu = x$ ,  $\nu \in \{1, 2, 3, \dots, n!\}$
- $\hat{\phi}(t) \in \mathrm{GL}_n(\mathbb{C}((t)))$
- $L \in \mathrm{gl}_n(\mathbb{C})$
- $Q = \mathrm{diag}(q_1, \dots, q_n)$ ,  $q_i \in \frac{1}{t}\mathbb{C}[\frac{1}{t}]$
- $L$  and  $Q$  commute.

One refers to the integer  $\nu$  as the *ramification* at 0 and the polynomial  $q_i$  as the *eigenvalues* at 0 of the equation. We note that even if  $A \in \mathrm{gl}_n(\mathbb{C}(\{x\}))$ , the matrix  $\hat{\phi}(t)$  does not necessarily lie in  $\mathrm{gl}_n(\mathbb{C}(\{t\}))$ . There are various ways to make this representation unique but we will not deal with this here. We also note that by adjusting the term  $x^L$ , we can always assume that no ramification occurs in  $\hat{\phi}(t)$ , that is,  $\hat{\phi} \in \mathrm{GL}_n(\mathbb{C}((x)))$  (but one may lose the last commutativity property above). As in the regular case, there are algorithms to determine a fundamental matrix of this form. We refer to Chapter 3 of [dPS03] for a detailed discussion of the existence of these formal solutions and algorithms to calculate them and p. 98 of [dPS03] for historical references and references to some recent work.

I will now give two examples (taken from [LR95]).

**Example 1.4.3** (Euler Eqn)  $x^2y' + y = x$

This inhomogeneous equation has a solution  $\hat{y} = \sum_{n=0}^{\infty} (-1)^n n! x^{n+1}$ . By applying the operator  $x \frac{d}{dx} - 1$  to both sides of this equation we get a second order homogeneous equation  $x^3y'' + (x^2 + x)y' - y = 0$  which, in matrix form becomes

$$\frac{dY}{dx} = \begin{pmatrix} 0 & 1 \\ \frac{1}{x^3} & -(\frac{1}{x} + \frac{1}{x^2}) \end{pmatrix} Y, \quad Y = \begin{pmatrix} y \\ y' \end{pmatrix}$$

This matrix equation has a fundamental solution matrix of the form  $\hat{Y} = \hat{\phi}(x)e^Q$ , where

$$Q = \begin{pmatrix} \frac{1}{x} & 0 \\ 0 & 0 \end{pmatrix} \quad \hat{\phi}(x) = \begin{pmatrix} 1 & \hat{f} \\ -\frac{1}{x^2} & \hat{f}' \end{pmatrix} \quad \hat{f} = \sum_{n=0}^{\infty} (-1)^n n! x^{n+1}$$

There is no ramification here (i.e.,  $\nu = 1$ ).

**Example 1.4.4** (Airy Equation)  $y'' = xy$

The singular point of this equation is at  $\infty$ . If we let  $z = \frac{1}{x}$  the equation becomes  $z^5y'' + 2z^4y' - y = 0$  or in matrix form

$$Y' = \begin{pmatrix} 0 & 1 \\ \frac{1}{z^5} & -\frac{2}{z} \end{pmatrix} Y$$

This equation has a fundamental matrix of the form  $\hat{Y} = \hat{\phi}(z)z^J U e^{Q(t)}$  where

$$\hat{\phi}(z) = \dots, \quad U = \begin{pmatrix} 1 & 1 \\ 1 & -1 \end{pmatrix}, \quad J = \begin{pmatrix} \frac{1}{4} & 0 \\ 0 & -\frac{3}{4} \end{pmatrix}$$

$$t = z^{\frac{1}{2}}, \quad Q = \begin{pmatrix} -\frac{2}{3t^3} & 0 \\ 0 & \frac{2}{3t^3} \end{pmatrix}$$

The ramification is  $\nu = 2$ . The precise form of  $\hat{\phi}(z)$  is not important at the moment but we do note that it lies in  $\text{GL}_n(\mathbb{C}((z)))$  (see [LR95] for the precise form).

We now turn to the Galois groups over  $\mathbb{C}((x))$  and  $\mathbb{C}(\{x\})$ .

Galois Group over  $\mathbb{C}((x))$ : the Formal Galois Group at  $x = 0$ . We consider the differential equation  $Y' = AY$ ,  $A \in \text{gl}_n(\mathbb{C}((x)))$ . To simplify the presentation while capturing many of the essential ideas we shall assume that there is no ramification in  $\hat{\phi}$ , that is, the equation has a fundamental

solution matrix  $\hat{Y} = \hat{\phi}(x)x^L e^{Q(1/t)}$ , with  $\hat{\phi}(x) \in \text{gl}_n(\mathbb{C}((x)))$ . The map  $Y_0 \mapsto \hat{\phi}(x)Y_0$  defines an equivalence over  $\mathbb{C}((x))$  between the equations

$$Y'_0 = A_0 Y_0 \text{ and } Y' = AY$$

where  $A_0 = \hat{\phi}^{-1} A \hat{\phi} - \hat{\phi}^{-1} \hat{\phi}'$ .

**Definition 1.4.5** *The equation  $Y' = A_0 Y$  is called a Normal Form of  $Y' = AY$ . It has  $Y_0 = x^L e^{Q(1/x)}$  as a fundamental solution matrix.*

**Example 1.4.6 (Euler Equation)** *For this equation (Example 1.4.3), the normal form is  $\frac{dY_0}{dx} = A_0 Y_0$ ,  $A_0 = \begin{pmatrix} -\frac{1}{x^2} & 0 \\ 0 & 0 \end{pmatrix}$   $Y_0 = e^{\begin{pmatrix} \frac{1}{x} & 0 \\ 0 & 0 \end{pmatrix}}$*

**Example 1.4.7 (Airy Equation)** *Using the solution in Example 1.4.4, we get a normal form  $\frac{dY_0}{dx} = A_0 Y_0$  with solution  $Y_0 = x^L U e^{Q(1/t)}$  with  $L, Q, t$  as in Example 1.4.4.*

Since the map  $Y_0 \mapsto \hat{\phi}(x)Y_0$  defines an isomorphism between the differential modules associated with  $Y'_0 = A_0 Y_0$  and  $Y' = AY$ , we have (Table 1.2) that  $F = \hat{\phi}(x)$  satisfies the differential equation

$$\frac{dF}{dx} = AF - FA_0$$

Since the normal form is equivalent over  $\mathbb{C}((x))$  to the original equation, its differential Galois group over  $\mathbb{C}((x))$  is the same as that of the original equation.

Let  $K$  be the Picard-Vessiot extension of  $\mathbb{C}((x))$  for  $\frac{dY_0}{dx} = A_0 Y_0$  (and therefore also for  $\frac{dY}{dx} = AY$ ). We have that

$$K \subset E = \mathbb{C}((x))(x^{\frac{1}{\nu}}, x^{\alpha_1}, \dots, x^{\alpha_n}, \log x, e^{q_1(x^{1/\nu})}, \dots, e^{q_n(x^{1/\nu})}).$$

We shall describe some differential automorphisms of  $E$  over  $\mathbb{C}((x))$ :

(1) Formal Monodromy  $\gamma$ : This is a differential automorphism of  $E$  given by the following conditions

- $\gamma(x^r) = e^{2\pi i r} x^r$
- $\gamma(\log x) = \log x + 2\pi i$
- $\gamma(e^{q(x^{1/\nu})}) = e^{q(\zeta x^{1/\nu})}$ ,  $\zeta = e^{\frac{2\pi i}{\nu}}$

(2) Exponential Torus  $\mathcal{T}$ : This is the group of differential automorphisms  $\mathcal{T} = \text{DGal}(E/F)$  where  $E = F(e^{q_1(x^{1/\nu})}, \dots, e^{q_n(x^{1/\nu})})$  and  $F = \mathbb{C}((x))(x^{\frac{1}{\nu}}, x^{\alpha_1}, \dots, x^{\alpha_n}, \log x)$ . Note that for  $\sigma \in \mathcal{T}$ :

- $\sigma(x^r) = x^r$
- $\sigma(\log x) = \log x$
- $\sigma(e^{q(x^{1/\nu})}) = c_\sigma e^{q(x^{1/\nu})}$

so we may identify  $\mathcal{T}$  with a subgroup of  $(\mathbb{C}^*)^n$ . One can show (Proposition 3.25 of [dPS03])

**Proposition 1.4.8** *If  $f \in E$  is left fixed by  $\gamma$  and  $\mathcal{T}$ , then  $f \in \mathbb{C}((x))$ .*

The formal monodromy and the elements of the exponential torus restrict to differential automorphisms of  $K$  over  $\mathbb{C}((x))$  and are again called by these names. Therefore the above proposition and the Fundamental Theorem imply

**Theorem 1.4.9** *The differential Galois group of  $Y' = AY$  over  $\mathbb{C}((x))$  is the Zariski closure of the group generated by the formal monodromy and the exponential torus.*

One can furthermore show (Proposition 3.25 of [dPS03]) that  $x = 0$  is a regular singular point if and only if the exponential torus is trivial.

**Example 1.4.10 (Euler Equation)** *We continue using the notation of Examples 1.4.3 and 1.4.6. One sees that  $K = E = \mathbb{C}((x))(e^{\frac{1}{x}})$  in this case. Furthermore, the formal monodromy is trivial. Therefore  $\text{DGal}(K/\mathbb{C}((x)))$  is equal to the exponential torus  $\mathcal{T} = \left\{ \begin{pmatrix} c & 0 \\ 0 & 1 \end{pmatrix} \mid c \in \mathbb{C}^* \right\}$*

**Example 1.4.11 (Airy Equation)** *We continue using the notation of Examples 1.4.4 and 1.4.7. In this case  $F = \mathbb{C}((z))(z^{\frac{1}{4}})$  and  $E = F(e^{\frac{2}{3z^{3/2}}})$ .*

*The formal monodromy  $\gamma = \begin{pmatrix} 0 & \sqrt{-1} \\ \sqrt{-1} & 0 \end{pmatrix}$  and the exponential torus*

*$\mathcal{T} = \left\{ \begin{pmatrix} c & 0 \\ 0 & c^{-1} \end{pmatrix} \mid c \neq 0 \right\}$ . Therefore*

$$\text{DGal}(K/\mathbb{C}((x))) = D_\infty = \left\{ \begin{pmatrix} c & 0 \\ 0 & c^{-1} \end{pmatrix} \mid c \neq 0 \right\} \cup \left\{ \begin{pmatrix} 0 & c \\ -c^{-1} & 0 \end{pmatrix} \mid c \neq 0 \right\}$$

Galois Group over  $\mathbb{C}(\{x\})$ : the Convergent Galois Group at  $x = 0$ . We now assume that we are considering a differential equation with  $\frac{dY}{dx} = AY$  with  $A \in \mathbb{C}(\{x\})$  and we let  $K$  be the Picard-Vessiot extension of  $\mathbb{C}(\{x\})$  for this equation. Since  $\mathbb{C}(\{x\}) \subset \mathbb{C}((x))$ , we have that the formal Galois group  $\mathrm{DGal}(K/\mathbb{C}((x)))$  is a subgroup of the convergent Galois group  $\mathrm{DGal}(K/\mathbb{C}(\{x\}))$ . If  $x = 0$  is a regular singular point then these Galois groups are the same but in general the convergent group is larger than the formal group. I will describe analytic objects that measure the difference. The basic idea is the following. We have a formal solution  $\hat{Y} = \hat{\phi}(x)x^Le^{Q(1/t)}$  for the differential equation. For each small enough sector  $\mathcal{S}$  at  $x = 0$  we can give the terms  $x^L$  and  $e^{Q(1/t)}$  meaning in an obvious way. The matrix  $\hat{\phi}(x)$  is a little more problematic but it has been shown that one can find a matrix  $\Phi_{\mathcal{S}}$  of functions, analytic in  $\mathcal{S}$ , “asymptotic” to  $\hat{\phi}(x)$  and such that  $\Phi_{\mathcal{S}}(x)x^Le^{Q(1/t)}$  satisfies the differential equation. In overlapping sectors  $\mathcal{S}_1$  and  $\mathcal{S}_2$  the corresponding  $\Phi_{\mathcal{S}_1}$  and  $\Phi_{\mathcal{S}_2}$  may not agree but they will satisfy  $\Phi_{\mathcal{S}_1} = \Phi_{\mathcal{S}_2}St_{12}$  for some  $St_{12} \in \mathrm{GL}_n(\mathbb{C})$ . The  $St_{i,j}$  ranging over a suitable choice of small sectors  $\mathcal{S}_{\ell}$  together with the formal Galois group should generate a group that is Zariski dense in the convergent Galois group. There are two (related) problems with this idea. The first is what do we mean by asymptotic and the second is how do we select the sectors and the  $\Phi_{\ell}$  in some canonical way? We shall first consider these questions for fairly general formal series and then specialize to the situation where the series are entries of matrices  $\hat{\phi}(x)$  that arise in the formal solutions of linear differential equations.

Asymptotics and Summability. In the 19<sup>th</sup> century, Poincaré proposed the following notion of an asymptotic expansion.

**Definition 1.4.12** *Let  $f$  be analytic on a sector  $\mathcal{S} = \mathcal{S}(a, b, \rho)$  at  $x = 0$ . The series  $\sum_{n \geq n_0} c_n x^n$  is an asymptotic expansion of  $f$  if  $\forall N \geq 0$  and every closed sector  $W \subset \mathcal{S}$  there exists a constant  $C(N, W)$  such that*

$$|f(x) - \sum_{n_0 \leq n \leq N-1} c_n x^n| \leq C(N, W)|x|^N \quad \forall x \in W.$$

A function  $f$  analytic on a sector can have at most one asymptotic expansion on that sector and we write  $J(f)$  for this series if it exists. We denote by  $\mathcal{A}(\mathcal{S}(a, b, \rho))$  the set of functions analytic on  $\mathcal{S}(a, b, \rho)$  having asymptotic expansions on this sector. We define  $\mathcal{A}(a, b)$  to be the direct limit of the  $\mathcal{A}(\mathcal{S}(a, b, \rho))$  over all  $\rho$  (here we say  $f_1 \in \mathcal{A}(\mathcal{S}(a, b, \rho_1))$  and

$f_2 \in \mathcal{A}(\mathcal{S}(a, b, \rho_2))$  are equivalent if they agree on the intersection of the sectors). If one thinks of  $a, b \in S^1$ , the unit circle, then one can see that the sets  $\mathcal{A}(a, b)$  form a sheaf on  $S^1$ . We have the following facts (whose proofs can be found in Chapters 7.1 and 7.2 of [dPS03] or in [Bal94]).

- $\mathcal{A}(a, b)$  is a differential  $\mathbb{C}$ -algebra.
- $\mathcal{A}(S^1) = \mathbb{C}(\{x\})$ .
- (Borel-Ritt) For all  $(a, b) \neq S^1$ , the map  $J : \mathcal{A}(a, b) \rightarrow \mathbb{C}((x))$  is surjective.

Given a power series in  $\mathbb{C}((x))$  that satisfies a linear differential equation, one would like to say that it is the asymptotic expansion of an analytic function that also satisfies the differential equation. Such a statement is contained in the next result, originally due to Hukuhara and Turrittin, with generalizations and further developments concerning nonlinear equations due to Ramis and Sibuya (see Chapter 7.2 of [dPS03] for a proof and references).

**Theorem 1.4.13** *Let  $A \in \text{gl}_n(\mathbb{C}(\{x\}))$ ,  $w \in (\mathbb{C}(\{x\}))^n$  and  $\hat{v} \in (\mathbb{C}((x)))^n$  such that*

$$\hat{v}' - A\hat{v} = w .$$

*For any direction  $\theta \in S^1$  there exist  $a, b$ ,  $a < \theta < b$  and  $v \in (\mathcal{A}(a, b))^n$  such that  $J(v) = \hat{v}$  and*

$$v' - Av = w$$

In general, there may be many functions  $v$  satisfying the conclusion of the theorem. To guarantee uniqueness we will refine the notion of asymptotic expansion and require that this be valid on large enough sectors. The refined notion of asymptotics is given by

**Definition 1.4.14** (1) *Let  $k$  be a positive real number,  $\mathcal{S}$  an open sector. A function  $f \in \mathcal{A}(\mathcal{S})$  with  $J(f) = \sum_{n \geq n_0} c_n x^n$  is Gevrey of order  $k$  if for all closed  $W \subset \mathcal{S}$ ,  $\exists A > 0, c > 0$  such that  $\forall N \geq 1$ ,  $x \in W$ ,  $|x| \leq c$  one has*

$$|f(z) - \sum_{n_0 \leq n \leq N-1} c_n x^n| \leq A^N (N!)^{\frac{1}{k}} |x|^N$$

*(One usually uses  $\Gamma(1 + \frac{N}{k})$  instead of  $(N!)^{\frac{1}{k}}$  but this makes no difference.)*



(2) We denote by  $\mathcal{A}_{\frac{1}{k}}(\mathcal{S})$  the ring of all Gevrey functions of order  $k$  on  $\mathcal{S}$  and let  $\mathcal{A}_{\frac{1}{k}}^0(\mathcal{S}) = \{f \in \mathcal{A}_{\frac{1}{k}}(\mathcal{S}) \mid J(f) = 0\}$

The following is a useful criterion to determine if a function is in  $\mathcal{A}_{\frac{1}{k}}^0(\mathcal{S})$ .

**Lemma 1.4.15**  $f \in \mathcal{A}_{\frac{1}{k}}^0(\mathcal{S})$  if and only if  $\forall$  closed  $W \subset \mathcal{S}, \exists A, B > 0$  such that

$$|f(x)| \leq A \exp(-B|x|^{-k}) \quad \forall x \in W$$

**Example 1.4.16**  $f(x) = e^{-\frac{1}{x^k}} \in \mathcal{A}_{\frac{1}{k}}^0(\mathcal{S})$ , for  $\mathcal{S} = \mathcal{S}(-\frac{\pi}{2k}, \frac{\pi}{2k})$  but not on a larger sector. This is a hint of what is to come.

Corresponding to the notion of a Gevrey function is the notion of a Gevrey series.

**Definition 1.4.17**  $\sum_{n \geq n_0} c_n x^n$  is a Gevrey series of order  $k$  if  $\exists A > 0$  such that  $\forall n > 0 \ |c_n| \leq A^n (n!)^{\frac{1}{k}}$ .

(2)  $\mathbb{C}((x))_{\frac{1}{k}}$  = Gevrey series of order  $k$ .

Note that if  $k < \ell$  then  $\mathbb{C}((x))_{\frac{1}{\ell}} \subset \mathbb{C}((x))_{\frac{1}{k}}$ . Using these definitions one can prove the following key facts:

- (Improved Borel-Ritt) If  $|b - a| < \frac{\pi}{k}$ , the map  $J : \mathcal{A}_{\frac{1}{k}}(a, b) \rightarrow \mathbb{C}((x))_{\frac{1}{k}}$  is surjective but not injective.
- (Watson's Lemma) If  $|b - a| > \frac{\pi}{k}$ , the map  $J : \mathcal{A}_{\frac{1}{k}}(a, b) \rightarrow \mathbb{C}((x))_{\frac{1}{k}}$  is injective but not surjective.

Watson's lemma motivates the next definition.

**Definition 1.4.18** (1)  $\hat{y} \in \mathbb{C}((x))$  is  $k$ -summable in the direction  $d \in S^1$  if there exists an  $f \in \mathcal{A}_{\frac{1}{k}}(d - \frac{\alpha}{2}, d + \frac{\alpha}{2})$  with  $J(f) = \hat{y}$ ,  $\alpha > \frac{\pi}{2}$ .

(2)  $\hat{y} \in \mathbb{C}((x))$  is  $k$ -summable if it is  $k$ -summable in all but a finite number of directions

A  $k$ -summable function has unique analytic “lifts” in many large sectors (certainly enough to cover a deleted neighborhood of the origin). I would like to say that the entries of the matrix  $\hat{\phi}(x)$  appearing in the formal solution of a linear differential equation are  $k$ -summable for some  $k$  but regrettably this is not exactly true and the situation is more complicated (each is a sum of  $k_i$ -summable functions for possible several  $k_i$ ). Before I describe what does happen, it is useful to have criteria for

deciding if a series  $\hat{y} \in \mathbb{C}((x))$  is  $k$ -summable. Such criteria can be given in terms of Borel and Laplace transforms.

**Definition 1.4.19** *The Formal Borel Transform  $\mathcal{B}_k$  of order  $k$ :*

$$\mathcal{B}_k : \mathbb{C}[[x]] \rightarrow \mathbb{C}[[\zeta]]$$

$$\mathcal{B}_k\left(\sum c_n x^n\right) = \sum \frac{c_n}{\Gamma(1 + \frac{n}{k})} \zeta^n$$

Note: If  $\hat{y} = \sum c_n x^n \in \mathbb{C}((x))_{\frac{1}{k}} \Rightarrow \mathcal{B}(\hat{y})$  has a nonzero radius of convergence.

**Example 1.4.20**  $\mathcal{B}_1(\sum (-1)^n n! x^{n+1}) = \sum (-1)^n \frac{\zeta^{n+1}}{n+1} = \log(1 + \zeta)$

**Definition 1.4.21** *The Laplace Transform  $\mathcal{L}_{k,d}$  of order  $k$  in the direction  $d$ : Let  $f$  satisfy  $|f(\zeta)| \leq A \exp(B|\zeta|^k)$  along the ray  $r_d$  from 0 to  $\infty$  in direction  $d$ . Then*

$$\mathcal{L}_{k,d} f(x) = \int_{r_d} f(\zeta) \exp(-(\frac{\zeta}{x})^k) d((\frac{\zeta}{x})^k)$$

Note:  $\mathcal{L}_{k,d} \circ \mathcal{B}_k(x^n) = x^n$  and  $\mathcal{L}_{k,d} \circ \mathcal{B}_k(f) = f$  for  $f \in \mathbb{C}\{\{x\}\}$ .

**Example 1.4.22** *For any ray  $r_d$  not equal to the negative real axis, the function  $f(\zeta) = \log(1 + \zeta)$  has an analytic continuation along  $r_d$ . Furthermore, for such a ray, we have*

$$\mathcal{L}_{1,d}(\log(1 + \zeta))(x) = \int_d \log(1 + \zeta) e^{-\frac{\zeta}{x}} d(\frac{\zeta}{x}) = \int_d \frac{1}{1 + \zeta} e^{-\frac{\zeta}{x}} d\zeta$$

Note that there are several slightly different definitions of the Borel and Laplace transforms scattered in the literature but for our purposes these difference do not affect the final results. I am following the definitions in [Bal94] and [dPS03]. The following gives useful criteria for a Gevrey series of order  $k$  to be  $k$ -summable (Chapter 3.1 of [Bal94] or Chapter 7.6 of [dPS03]).

**Proposition 1.4.23**  $\hat{y} \in \mathbb{C}[[x]]_{\frac{1}{k}}$  is  $k$ -summable in direction  $d$

$$\Updownarrow$$

$\mathcal{B}_k(\hat{y})$  has an analytic continuation  $h$  in a full sector  $\{\zeta \mid 0 < |\zeta| < \infty, |\arg \zeta - d| < \epsilon\}$  and has exponential growth of order  $\leq k$  at  $\infty$  in this sector ( $|h(\zeta)| \leq A \exp(B|\zeta|^k)$ ). In this case,  $f = \mathcal{L}_{k,d}(h)$  is its  $k$ -sum.

**Example 1.4.24** Consider  $\hat{y} = \sum (-1)^n n! x^{n+1}$ , a power series of Gevrey order 1 that is also a solution of the Euler equation. We have seen that  $f(\zeta) = \mathcal{B}_1(\hat{y}) = \sum (-1)^n \frac{\zeta^{n+1}}{n+1} = \log(1 + \zeta)$  for  $|\zeta| < 1$ . As we have noted in Example 1.4.22, this function can be continued outside the unit disk in any direction other than along the negative real axis. On any ray except this one, it has exponential growth of order at most 1 and its Laplace transform is

$$y_d(x) = \mathcal{L}_{1,d}(\log(1 + \zeta))(x) = \int_d \log(1 + \zeta) e^{-\frac{\zeta}{x}} d\left(\frac{\zeta}{x}\right) = \int_d \frac{1}{1 + \zeta} e^{-\frac{\zeta}{x}} d\zeta.$$

We note that this function again satisfies the Euler equation. To see this note that  $y(x) = Ce^{\frac{1}{x}}$  is a solution of  $x^2 y' + y = 0$  and so variation of constants gives us that  $f(x) = \int_0^x e^{\frac{1}{x} - \frac{1}{t}} \frac{dt}{t}$  is a solution of  $x^2 y' + y = x$ . for convenience let  $d = 0$  and define a new variable  $\zeta$  by  $\frac{\zeta}{x} = \frac{1}{t} - \frac{1}{x}$  gives  $f(x) = \int_{d=0} \frac{1}{1 + \zeta} e^{-\frac{\zeta}{x}} d\zeta$ . Therefore, for any ray except the negative real axis, we are able to canonically associate a function, analytic in a large sector around this ray, with the divergent series  $\sum (-1)^n n! x^{n+1}$  and so this series is 1-summable. Furthermore these functions will again satisfy the Euler equation. This is again a hint for what is to come.

Linear Differential Equations and Summability. It has been known since the early 20<sup>th</sup> Century that a formal series solution of an analytic differential equation  $F(x, y, y', \dots, y^{(n)}) = 0$  must be Gevrey of order  $k$  for some  $k$  ([Mai03]). Regretably, even for linear equations, these formal solutions need not be  $k$  summable (see [LR95]). Nonetheless, if  $\hat{y}$  is a vector of formal series that satisfy a linear differential equation it can be written as a sum  $\hat{y} = \hat{y}_1 + \dots + \hat{y}_t$  where each  $\hat{y}_i$  is  $k_i$  summable for some  $k_i$ . This result is the culmination of the work of several authors including Ramis, Malgrange, Martinet, Sibuya and Écale (see [LR95], [Ram93] and Chapter 7.8 of [dPS03]). The  $k_i$  that occur as well as the directions along which the  $\hat{y}_i$  are not summable can be seen from the normal form of the differential equation. We need the following definitions. Let  $Y' = AY$ ,  $A \in \text{gl}_n(\mathbb{C}(\{x\}))$  and let  $\hat{Y} = \hat{\phi}(x) x^L e^{Q(1/t)}$  be a formal solution with  $t^\nu = x$  and  $Q = \text{diag}(q_1, \dots, q_n)$ ,  $q_i \in \frac{1}{t} \mathbb{C}[\frac{1}{t}]$ .

**Definition 1.4.25** (1) The rational number  $k_i = \frac{1}{\nu}( \text{the degree of } q_i \text{ in } \frac{1}{t} )$  is called a slope of  $Y' = AY$ .

(2) A Stokes direction  $d$  for a  $q_i = cx^{-k_i} + \dots$  is a direction such that  $\text{Re}(cx^{-k_i}) = 0$ , i.e., where  $e^{q_i}$  changes behavior.

(3) Let  $d_1, d_2$  be consecutive Stokes directions. We say that  $(d_1, d_2)$  is a negative Stokes pair if  $\operatorname{Re}(cx^{-k_i}) < 0$  for  $\arg(x) \in (d_1, d_2)$ .

(4) A singular direction is the bisector of a negative Stokes pair.

**Example 1.4.26** For the Euler equation we have that  $\nu = 1$  and  $Q(x) = \operatorname{diag}(\frac{1}{x}, 0)$ . Therefore the slopes are  $\{0, 1\}$ . The Stokes directions are  $\{\frac{\pi}{2}, \frac{3\pi}{2}\}$ . The pair  $(\frac{\pi}{2}, \frac{3\pi}{2})$  is a negative Stokes pair and so  $\pi$  is the only singular direction.

With these definitions we can state


**Theorem 1.4.27** (*Multisummation Theorem*) Let  $1/2 < k_1 < \dots < k_r$  be the slopes of  $Y' = AY$  and let  $\hat{Y} \in (\mathbb{C}((x)))^n$  satisfy this differential equation. For all directions  $d$  that are not singular directions, there exist  $\hat{Y}_i \in \mathbb{C}((x))$  such that  $\hat{Y} = \hat{Y}_1 + \dots + \hat{Y}_r$ , each  $\hat{Y}_i$  is  $k_i$ -summable in direction  $d$  and, for  $y_i =$  the  $k_i$ -sum of  $\hat{Y}_i$ ,  $y_d = y_1 + \dots + y_r$  satisfies  $y' = Ay$ .

The condition that  $1/2 < k_1$  is a needed technical assumption but, after a suitable change of coordinates, one can always reduce to this case. We shall refer to the element  $y_d$  as above as the *multisum* of  $\hat{Y}$  in the direction  $d$ .

*Stokes Matrices and the Convergent Differential Galois Group.* We are now in a position to describe the analytic elements which, together with the formal Galois group determine the analytic Galois group. Once again, let us assume that we have a differential equation  $Y' = AY$  with  $A \in \operatorname{gl}_n(\mathbb{C}(\{x\}))$  and assume this has a formal solution  $\hat{Y} = \hat{\phi}(x)x^L e^{Q(1/t)}$  with  $\hat{\phi}(x) \in \operatorname{GL}_n(\mathbb{C}((x)))$ . The matrix  $F = \hat{\phi}(x)$  is a formal solution of the  $n^2 \times n^2$  system

$$\frac{dF}{dx} = AF - FA_0 \quad (1.3)$$

where  $Y' = A_0Y$  is the normal form of  $Y' = AY$ . We will want to associate functions, analytic in sectors, to  $\hat{\phi}(x)$  and therefore will apply the Multisummation Theorem to equation 1.3 (and not to  $Y' = AY$ ). One can show that the eigenvalues of equation (1.3) are of the form  $q_i - q_j$  where the  $q_i, q_j$  are eigenvalues of  $Y' = AY$ .



Let  $d$  be a singular direction of  $Y' = AY - YA_0$ . Select  $d^+$  and  $d^-$  nearby so that the associated sectors overlap and contain  $d$ . Let  $\phi^+, \phi^-$  be the associated multisums. We must have  $\phi^+ x^L e^{Q(1/t)} = \phi^- x^L e^{Q(1/t)} \cdot St_d$  for some  $St_d \in GL_n(\mathbb{C})$  on the overlap of the regions (the shaded area in the figure). The matrix  $St_d$  is independent of the choice of  $d^+$  and  $d^-$ .

**Definition 1.4.28** The matrix  $St_d$  is called a Stokes matrix in the direction  $d$ .

**Example 1.4.29** (Euler equation)

- Eigenvalues of  $Y' = AY - YA_0$ :  $\{\frac{1}{x}, 0\}$
- Singular direction of  $Y' = AY - YA_0$ :  $d = \pi$
- $\hat{\phi}(x) = \begin{pmatrix} 1 & \hat{f} \\ -\frac{1}{x^2} & \hat{f}' \end{pmatrix}$   $\hat{f} = \sum_{n=0}^{\infty} (-1)^n x^{n+1}$

For  $d^+$  and  $d^-$  close to the negative real axis, we have

$$f^+ = \int_{d^+} \frac{1}{1+\zeta} e^{-\frac{\zeta}{x}} d\zeta \text{ and } f^- = \int_{d^-} \frac{1}{1+\zeta} e^{-\frac{\zeta}{x}} d\zeta.$$

Since  $f^+ - f^- = 2\pi i \operatorname{Res}_{\zeta=-1} \left( \frac{e^{-\frac{\zeta}{x}}}{1+\zeta} \right) = 2\pi i$ , we have

$$\begin{pmatrix} 1 & f^+ \\ -\frac{1}{x^2} & (f^+)' \end{pmatrix} = \begin{pmatrix} 1 & f^- \\ \frac{1}{x^2} & (f^-)' \end{pmatrix} \begin{pmatrix} 1 & 2\pi i \\ 0 & 1 \end{pmatrix}$$

$$\text{so } St_\pi = \begin{pmatrix} 1 & 2\pi i \\ 0 & 1 \end{pmatrix}$$

The following result is due to Ramis [Ram85a, Ram85c, Ram85b, MR89]. Another proof appears in the work of Loday-Richaud [LR94]. An exposition of this result and its proof appears in Chapter 8 of [dPS03].

**Theorem 1.4.30** The convergent differential Galois group  $\operatorname{DGal}(K/\mathbb{C}(\{x\}))$  is the Zariski closure of the group generated by

- the formal differential Galois group  $\operatorname{DGal}(K/\mathbb{C}((x)))$ , and
- The collection of Stokes matrices  $\{St_d\}$  where  $d$  runs over the set of singular directions of the polynomials  $\{q_i - q_j\}$ ,  $q_i, q_j$  eigenvalues of  $Y' = AY$ .

**Examples 1.4.31** (1) (Euler equation)

- The formal differential Galois group:

$$\left\{ \begin{pmatrix} c & 0 \\ 0 & 1 \end{pmatrix} \mid c \neq 0 \right\}$$

- Stokes matrix:  $\begin{pmatrix} 1 & 2\pi i \\ 0 & 1 \end{pmatrix}$

Therefore the convergent differential Galois group is

$$\left\{ \begin{pmatrix} c & d \\ 0 & 1 \end{pmatrix} \mid c \neq 0, d \in \mathbb{C} \right\}.$$

(2) (Airy equation)

- The formal differential Galois group:

$$D_\infty = \left\{ \begin{pmatrix} c & 0 \\ 0 & c^{-1} \end{pmatrix}, \begin{pmatrix} 0 & c \\ -c^{-1} & 0 \end{pmatrix} \mid c \neq 0 \right\}$$

- The Stokes matrices:

$$St_0 = \begin{pmatrix} 1 & * \\ 0 & 1 \end{pmatrix}, St_{\frac{2\pi}{3}} = \begin{pmatrix} 1 & 0 \\ * & 1 \end{pmatrix}, St_{\frac{4\pi}{3}} = \begin{pmatrix} 1 & * \\ 0 & 1 \end{pmatrix}$$

(see Example 8.15 of [dPS97] for the calculation of the Stokes matrices.)

Therefore the convergent differential Galois group is  $SL_2(\mathbb{C})$

We highly recommend the book [Ram93] of Ramis, the papers of Loday-Richaud [LR90, LR95] and the papers of Varadarajan [Var91, Var96] for further introductions to this analytic aspect of linear differential equations. One can find references to the original papers in these works as well as in [dPS03].

## 1.5 Algorithms

In this section we shall consider the question: Can we compute differential Galois groups and their properties? We shall not give a complete algorithm to determine the Galois groups although one exists - see [Hru02]. This general algorithm has not been implemented at present and, in its present form seems not very efficient. We will rather give a flavor of the various techniques that are presently implemented and used.

**Categories motivate algorithms.** We will show how results in a categorical setting concerning representations of groups lead to algorithms to determine Galois groups. We begin with an example from the Galois theory of polynomials.

**Example 1.5.1** *Let*

$$P(Y) = Y^3 + bY + c = 0 \in \mathbb{Q}[Y].$$

*The usual Galois group  $G$  of this equation is a subgroup of the full permutation group  $\mathcal{S}_3$  on 3 elements. We will describe how to compute  $G$ . If  $P(Y) = (Y + \alpha)(Y^2 + \beta Y + \gamma)$ ,  $\alpha, \beta, \gamma \in \mathbb{Q}$ , then  $G$  is the Galois group of  $Y^2 + \beta Y + \gamma$ , which we assume we know how to compute. Therefore we will assume that  $P(Y)$  is irreducible over  $\mathbb{Q}$ . This implies that  $G$  acts transitively on the roots  $\{\rho_1, \rho_2, \rho_3\}$  of  $P$ . From group theory we know that the only transitive subgroups of  $\mathcal{S}_3$  are  $\mathcal{A}_3$ , the alternating group on 3 elements, and  $\mathcal{S}_3$ . To decide which we have, consider the following associated polynomial*

$$Z^2 + 4b^3 + 27c^2 = (Z + \delta)(Z - \delta) \quad \delta = (r_1 - r_2)(r_2 - r_3)(r_3 - r_1).$$

*One then has that  $G = \mathcal{A}_3$  if  $Z^2 + 4b^3 + 27c^2$  factors over  $\mathbb{Q}$  and  $G = \mathcal{S}_3$  if  $Z^2 + 4b^3 + 27c^2$  is irreducible over  $\mathbb{Q}$ .*

In the above example we determined the Galois group by considering factorization properties of the polynomial and associated polynomials. Why does this work?

To answer this question, let us consider a square-free polynomial  $f(Y) \in \mathbb{Q}[Y]$  with roots  $r_1, \dots, r_n \in \mathbb{Q}$ , the algebraic closure of  $\mathbb{Q}$ . Let  $G$  be the Galois group of  $f$  over  $\mathbb{Q}$ . The group  $G$  acts as a group of permutations on the set  $S = \{r_1, \dots, r_n\}$ . We have that  $G$  acts transitively on  $S$  if and only if the polynomial  $f$  is irreducible. More generally,  $G$  leaves a subset  $\{r_{i_1}, \dots, r_{i_t}\}$  invariant if and only if the polynomial  $g(Y) = \prod_{j=1}^t (Y - r_{i_j}) \in \mathbb{Q}[Y]$ . Therefore the orbits of  $G$  in the set  $S$  correspond to irreducible factors of  $f$ . In fact, for any  $h \in \mathbb{Q}[Y]$  having all its roots in the splitting field of  $f$ ,  $G$ -orbits in the set of roots correspond to irreducible factors of  $h$ . In the above example, we were able to distinguish between groups because they have different orbits in certain “well-selected” sets and we were able to see this phenomenon via factorization properties of associated polynomials. The key is that, in general,

*one can distinguish between finite groups by looking at the sets on which they act.*

This strange looking statement can be formalized in the following way (see Appendix B1 of [dPS03] for details). For any finite set, let  $\text{Perm}(S)$  be the group of permutations of  $S$ . One constructs a category  $\text{Perm}_G$  of  $G$ -sets whose objects are pairs  $(S, \rho)$ , where  $\rho : G \rightarrow \text{Perm}(S)$  is the map defining an action of  $G$  on  $S$ . The morphisms of this category are just maps between  $G$ -sets that commute with the action of  $G$ . One has the forgetful functor  $\omega : \text{Perm}_G \rightarrow \text{Sets}$  which takes  $(S, \rho)$  to  $S$ . One then has the formalization of the above statement as: one can recover the group from the pair  $(\text{Perm}_G, \omega)$ . A consequence of this is that given two groups  $H \subset G$  there is a set  $S$  on which  $G$  (and therefore  $H$ ) acts so that the orbit structures with respect to these two groups are different.

These facts lead to the following strategy to compute Galois groups of a polynomial  $f \in \mathbb{Q}[X]$ . For each pair of groups  $H \subset G \subset \mathcal{S}_n$  we construct a polynomial  $f_{H,G} \in \mathbb{Q}[X]$  whose roots can be expressed in terms of the roots of  $f$  and such that  $H$  and  $G$  can be distinguished by their orbit structures on the roots of  $f_{H,G}$  (by the above facts, we know that there is a set that distinguishes  $H$  from  $G$  and one can show that this set can be constructed from the roots of  $f$ ). The factorization properties of  $f_{H,G}$  will then distinguish  $H$  from  $G$ . This is what was done in Example 1.5.1.

To generalize the above ideas to calculating differential Galois groups of linear differential equations, we must first see what determines a linear algebraic group. In fact, just as finite groups are determined by their *permutation* representations, linear algebraic groups are determined by their *linear* representations. To make this more precise, let  $G$  be a linear algebraic group defined over an algebraically closed field  $C$ . I will denote by  $\text{Rep}_G$  the category of linear representations of  $G$ . The objects of this category are pairs  $(V, \rho)$  where  $V$  is a finite dimensional vector space over  $C$  and  $\rho : G \rightarrow \text{GL}_n(V)$  is a representation of  $G$ . The morphisms  $m : (V_1, \rho_1) \rightarrow (V_2, \rho_2)$  are linear maps  $m : V_1 \rightarrow V_2$  such that  $m \circ \rho_1 = \rho_2 \circ m$ . One can define subobjects, quotients, duals, direct sums and tensor products in obvious ways (see Appendix B2 and B3 of [dPS03] for details). One again has a forgetful functor  $\omega : \text{Rep}_G \rightarrow \text{Vect}_C$  (where  $\text{Vect}_C$  is the category of vector spaces over  $C$ ) given by  $\omega(V, \rho) = V$ . In analogy to the finite group case, we have Tannaka's Theorem, that says that we can recover  $G$  from the pair  $(\text{Rep}_G, \omega)$ .



Let  $\mathcal{T}$  be a category with a notion of quotients, duals, direct sums, tensor products and a functor  $\tilde{\omega} : \mathcal{T} \rightarrow \text{Vect}_C$ . An example of such a category is the category of differential modules  $\mathcal{D}$  over a differential field  $k$  with algebraically closed constants and  $\tilde{\omega}(M) = \ker(\partial, K \otimes M)$  (see the discussion following Definition 1.2.13) where  $K$  is a differential field large enough to contain solutions of the differential equations corresponding to the modules in  $\mathcal{D}$ . One can give axioms that guarantee that  $\tilde{\omega}(\mathcal{T}) = \omega(\text{Rep}_G)$  for some linear algebraic group  $G$  (see [DM82], [Del90] or Appendix B3 of [dPS03]). Such a pair  $(\mathcal{T}, \tilde{\omega})$  is called a *neutral tannakian category*. Applying this to a subcategory of  $\mathcal{D}$ , one has:

**Theorem 1.5.2** *Let  $(k, \partial)$  be a differential field with algebraically closed constants, let  $Y' = AY$  be a linear differential equation over  $(k, \partial)$ , let  $M_A$  be the associated differential module and  $K$  the associated Picard-Vessiot extension. Let  $\mathcal{T} = \{\{M_A\}\}$  be the smallest subcategory of  $\mathcal{D}$  containing  $M_A$  and closed under the operations of taking submodules, quotients, duals, direct sums, and tensor products. For  $N \in \{\{M_A\}\}$ , let  $\tilde{\omega}(N) = \ker(\partial, N \otimes K)$ .*

*Then  $(\mathcal{T}, \tilde{\omega})$  is a neutral tannakian category and the image of  $\tilde{\omega}$  is the category of vector spaces on which  $\text{DGal}(K/k)$  acts.*

This theorem and Tannaka's Theorem have the following implications:

- One can construct all representations of  $\text{DGal}(K/k)$  by:
  - using the constructions of linear algebra (submodules, quotients, duals, direct sums, tensor products) on  $M_A$ , and
  - taking the solution spaces of the corresponding differential equations.
- $\text{DGal}(K/k)$  is determined by knowing which subspaces of these solution spaces are  $\text{DGal}(K/k)$ -invariant.

At first glance, it seems that one would need to consider an infinite number of differential modules but in many cases, it is enough to consider only a finite number of modules constructed from  $M_A$ . Regretably, there are groups (for example, any group of the form  $C^* \oplus \dots \oplus C^*$ ) for which one needs an infinite number of representations to distinguish it from other groups. Nonetheless, there are many groups where a finite number of representations suffice. Furthermore, Proposition 1.3.28 states that the  $\text{DGal}$ -invariant subspaces of the solution space of a differential module  $N$  correspond to factors of the associated scalar equation  $L_N(Y) = 0$ .

Therefore, in many cases, one can give criteria for a differential equation to have a given group as Galois group in terms of factorization properties of certain associated differential operators. We shall give examples for second order equations. We begin with a definition.

Let  $k$  be a differential field with algebraically closed constants  $C$ , let  $L = \partial^2 - s \in k[\partial]$ , and let  $K$  be the Picard-Vessiot extension corresponding to  $L(Y) = 0$ . Let  $\{y_1, y_2\}$  be a basis for the solution space  $V$  of  $L(Y) = 0$  in  $K$  and let  $V_m = C - \text{span of } \{y_1^m, y_1^{m-1}y_2, \dots, y_2^m\}$ . One can see that  $V_m$  is independent of the selected basis of  $V$  and that it is invariant under the action of  $\text{DGal}(K/k)$ . As in the proof of Proposition 1.3.28 one can show that this implies that  $V_m$  is precisely the solution space of a scalar linear differential equation over  $k$ .

**Definition 1.5.3** *The  $m^{\text{th}}$  symmetric power  $L^{\odot m}$  of  $L$  is the monic operator whose solution space is  $V_m$ .*

One has that the order of  $L^{\odot m}$  is  $m + 1$ . To see this it is enough to show that the  $y_1^{m-i}y_2^i$ ,  $0 \leq i \leq m$  are linearly independent over  $C$ . Since any homogeneous polynomial in two variables factors over  $C$ , if we had  $0 = \sum c_i y_1^{m-i}y_2^i = \prod (a_i y_1 + b_i y_2)$ , then  $y_1$  and  $y_2$  would be linearly dependent over  $C$ . One can calculate  $L^{\odot m}$  by starting with  $y^m$  and formally differentiating it  $m$  times. One uses the relation  $y'' - sy = 0$  to replace  $y^{(j)}$ ,  $j > 1$  with combinations of  $y$  and  $y'$ . One then has  $m+2$  expressions  $y^m, (y^m)', \dots, (y^m)^{(m+1)}$  in the  $m+1$  “indeterminates”  $y^{m-i}(y')^i$ ,  $0 \leq i \leq m$ . These must be linearly dependent and one can find a dependence  $(y^m)^{(m+1)} + b_m(y^m)^{(m)} + \dots + b_0(y^m) = 0$ . One can show that  $L^{\odot m} = \partial^{m+1} + b_m \partial^m + \dots + b_0$  (see Chapter 2.3 of [dPS03]).

**Example 1.5.4**  $m = 2$ :

$$\begin{aligned} y^2 &= y^2 \\ (y^2)' &= 2yy' \\ (y^2)'' &= 2(y')^2 + 2sy^2 \\ (y^2)''' &= 8syy' + 2s'y^2 \end{aligned} \quad \Rightarrow \quad L^{\odot 2}(y) = y''' - 4sy' - 2s'y$$

The vector spaces  $V_m$  correspond to representations of  $\text{DGal}(K/k)$  on the  $m^{\text{th}}$  symmetric power of  $V$  and are well understood for many groups [FH91] (the  $m^{\text{th}}$  symmetric power of a vector space is the vector space with a *basis* formed by the monomials of degree  $m$  in a basis of  $V$ ). Using representation theory, one can show:

**Proposition 1.5.5** *Let  $L(y) = y'' - sy$  and let  $G = \text{DGal}(K/k)$ . Assume  $L$  and  $L^{\otimes 2}$  are irreducible.*

- *If  $L^{\otimes 3}$  is reducible, then  $G/\pm I \simeq A_4$ .*
- *If  $L^{\otimes 3}$  is reducible and  $L^{\otimes 4}$  is irreducible then  $G/\pm I \simeq S_4$ .*
- *If  $L^{\otimes 4}$  is reducible, and  $L^{\otimes 6}$  is irreducible, then  $\Rightarrow G/\pm I \simeq A_5$ .*
- *If  $L^{\otimes 6}$  is irreducible then  $G \simeq \text{SL}_2(C)$ .*

In order to implement these criteria, one needs good algorithms to factor linear operators. Such algorithms exist and I refer to Chapter 4.2 of [dPS03] for descriptions of some algorithms that factor linear operators as well as further references to the extensive literature. The calculation of symmetric powers and factorization of operators has been implemented in MAPLE in the DETools package.

One can develop similar criteria for higher order operators. Given a differential operator  $L$  of order  $n$  with  $\{y_1, \dots, y_n\}$  being a basis of the solution space  $V$  in some Picard-Vessiot extension, one can consider the  $C$ -space  $V_m$  of homogeneous polynomials of degree  $m$  in the  $y_i$ . This space has dimension at most  $\binom{m+n-1}{n-1}$  but may have dimension that is strictly smaller and so is not isomorphic to the  $m^{\text{th}}$  symmetric power of  $V$  (although it is isomorphic to a quotient). This complicates matters but this situation can be dealt with by casting things directly in terms of differential modules, forming the symmetric powers of differential modules and developing algorithms to find submodules of differential modules. This is explained in Chapter 4.2 of [dPS03]. Examples of criteria for third order equations are given in Chapter 4.3.5 of [dPS03] where further references are given as well.

### **Solving $L(y) = 0$ in terms of exponentials, integrals and**

**algebraics.** I shall give a formal definition that captures the meaning of this term.

**Definition 1.5.6** *Let  $k$  be a differential field,  $L \in k[\partial]$ , and  $K$  be the associated Picard-Vessiot extension.*

1) *A liouvillian tower over  $k$  is a tower of differential fields  $k = K_0 \subset \dots \subset K_m$  with  $K \subset K_m$  and, for each  $i, 0 \leq i < m$ ,  $K_{i+1} = K_i(t_i)$ , with*

- *$t_i$  algebraic over  $K_i$ , or*
- *$t'_i \in K_i$ , i.e.,  $t_i = \int u_i, u_i \in K_i$ , or*
- *$t'_i/t_i \in K_i$ , i.e.,  $t_i = e^{\int u_i}, u_i \in K_i$*

- 2) An element of liouvillian tower is said to be liouvillian over  $k$ .  
 3)  $L(y) = 0$  is solvable in terms of liouvillian functions if the associated PV-extension  $K$  lies in a liouvillian tower over  $k$ .

Picard and Vessiot stated the following Galois theoretic criteria and this was given a formal modern proof by Kolchin [Kol48, Kol76].

**Theorem 1.5.7** *Let  $k, L$ , and  $K$  be as above.  $L(y) = 0$  is solvable in terms of liouvillian functions if and only if the identity component (in the Zariski topology) of  $\mathrm{DGal}(K/k)$  is solvable.*

Recall that a group  $G$  is solvable if there exists a tower of subgroups  $G = G_0 \supset G_1 \supset \dots \supset G_m = \{e\}$  such that  $G_{i+1}$  is normal in  $G_i$  and  $G_{i+1}/G_i$  is abelian. This above theorem depends on the

**Theorem 1.5.8 (Lie-Kolchin Theorem)** *Let  $C$  be an algebraically closed field. A Zariski connected solvable group  $G \subset \mathrm{GL}_n(C)$  is conjugate to a group of triangular matrices. In particular,  $G$  leaves a one dimensional subspace invariant.*

I shall show how the Lie-Kolchin Theorem can be strengthened resulting in a strengthened version of Theorem 1.5.7 that leads to an algorithm to decide if a given linear differential equation (over  $C(x)$ ) can be solved in terms of liouvillian functions. I begin with

**Proposition 1.5.9** *Let  $k$  be a differential field,  $L \in k[\partial]$ ,  $K$  the corresponding Picard-Vessiot extensions and  $G$  the differential Galois group of  $K$  over  $k$ . The following are equivalent:*

- (i)  $L(y) = 0$  has a liouvillian solution  $\neq 0$ .
- (ii)  $G$  has a subgroup  $H$ ,  $|G : H| = m < \infty$  such that  $H$  leaves a one dimensional subspace invariant.
- (iii)  $L(y) = 0$  has a soln  $z \neq 0$  such that  $z'/z$  algebraic over  $k$  of degree  $\leq m$ .

*Proof* (Outline; see Chapter 4.3 of [dPS03]) (iii)  $\Rightarrow$  (i): Clear.

(i)  $\Rightarrow$  (ii): One can reduce this to the case where all solutions are liouvillian. In this case, the Lie-Kolchin Theorem implies that the identity component  $G^0$  of  $G$  leaves a one dimensional subspace invariant.

(ii)  $\Rightarrow$  (iii): Let  $V = \mathrm{Soln}(L)$  and  $v \in V$  span an  $H$ -invariant line. We

then have that  $\forall \sigma \in H, \exists c_\sigma \in C_k$  such that  $\sigma(v) = c_\sigma v$ . This implies that  $\forall \sigma \in H, \sigma(\frac{v'}{v}) = \frac{(c_\sigma v)'}{c_\sigma v} = \frac{v'}{v}$ . Therefore, the Fundamental Theorem implies that  $\frac{v'}{v} \in E = \text{fixed field of } H$ . One can show that  $[E : k] = |G : H| = m$  so  $\frac{v'}{v}$  is algebraic over  $k$  of degree at most  $m$ .  $\square$

In order to use this result, one needs a bound on the integer  $m$  that can appear in (ii) and (iii) above. This is supplied by the following group theoretic result (see Chapter 4.3.1 of [dPS03] for references).

**Lemma 1.5.10** *There exists a function  $I(n)$  such that if  $G \subset \text{GL}_n(\mathbb{C})$ ,  $H \subset G$  satisfies*

- (i)  $|G : H| < \infty$  and
- (ii)  $H$  leaves a one dimension subspace invariant

*then there exists a subgroup  $\tilde{H} \subset G$  such that*

- (i)  $|G : \tilde{H}| < I(n)$  and
- (ii)  $\tilde{H}$  leaves a one dimensional subspace invariant.

In general we know that  $I(n) \leq n^{2n^2+2}$ . For small values of  $n$  we have exact values, (e.g.,  $I(2) = 12$ ,  $I(3) = 360$ ). This clearly allows us to deduce the following corollary to Proposition 1.5.9.

**Corollary 1.5.11** *Let  $k, L, K$  be as in Proposition 1.5.9 with the order of  $L$  equal to  $n$ . The following are equivalent*

- (i)  $L(y) = 0$  has a liouvillian solution  $\neq 0$ .
- (ii)  $G$  has a subgroup  $H$ ,  $|G : H| \leq I(n)$  such that  $H$  leaves a one dimensional subspace invariant.
- (iii)  $L(y) = 0$  has a soln  $z \neq 0$  such that  $z'/z$  algebraic /  $k$  of deg  $\leq I(n)$ .

This last result leads to several algorithms to decide if a differential equation  $L(y) = 0$ ,  $L \in C(x)[\partial]$ ,  $C$  a finitely generated subfield of  $\mathbb{C}$ , has a liouvillian solution. The one presented in [Sin81] (containing ideas going back to [Bou98]) searches for a putative minimal polynomial  $P(u) = a_m u^m + a_{m-1} u^{m-1} + \dots + a_0$ ,  $a_i \in \mathbb{C}[x]$  of an element  $u = z'/z$  where  $z$  is a solution of  $L(y) = 0$  and  $m \leq I(n)$ . This algorithm shows how the degrees of the  $a_i$  in  $x$  can be bounded in terms of information calculated at each singular point of  $L$ . Once one has degree bounds, the actual coefficients  $a_i$  can be shown to satisfy a system of polynomial equations and one can (in theory) use various techniques (e.g., Gröbner bases) to solve these. Many improvements and new ideas have been

given since then (see Chapter 4 of [dPS03]). We shall present criteria that form the basis of one method, describe what one needs to do to use this in general and give details for finding liouvillian solutions of second order differential equations.

**Proposition 1.5.12** (1) *Let  $G$  be a subgroup of  $\mathrm{GL}_n$ . There exists a subgroup  $H \subset G$ ,  $|G : H| \leq I(n)$  with  $H$  leaving a one dimensional space invariant if and only if  $G$  permutes a set of at most  $I(n)$  one dimensional subspaces.*

(2) *Let  $k, L, K, G$  be as in Proposition 1.5.9 with the order of  $L$  equal to  $n$ . Then  $L(y) = 0$  has a liouvillian solution if and only if  $G \subset \mathrm{GL}_n$  permutes a set of at most  $I(n)$  one dimensional subspaces.*

*Proof* (1) Let  $\ell$  be a one dimensional space left invariant by  $H$ . The orbit of  $\ell$  under the action of  $G$  has dimension at most  $|G : H|$ . Now, let  $\ell$  be a one dimensional space whose orbit under  $G$  is at most  $I(n)$  and let  $H$  be the stabilizer of  $\ell$ . We then have  $|G : H| \leq I(n)$ . (2) is an immediate consequence of (1) and the previous proposition.  $\square$

To apply this result, we need the following definition:

**Definition 1.5.13** *Let  $V$  be a vector space of dimension  $n$  and  $t \geq 1$  be an integer. The  $t^{\text{th}}$  symmetric power  $\mathrm{Sym}^t(V)$  of  $V$  is the quotient of the  $t$ -fold tensor product  $V^{\otimes t}$  by the subspace generated by elements  $v_1 \otimes \dots \otimes v_t - v_{\pi(1)} \otimes \dots \otimes v_{\pi(t)}$ , for all  $v_i \in V$  and  $\pi$  a permutation.*

We denote by  $v_1 v_2 \dots v_t$  the image of  $v_1 \otimes \dots \otimes v_t$  in  $\mathrm{Sym}^t(V)$ . If  $e_1, \dots, e_n$  is a basis of  $V$  and  $1 \leq t \leq n$ , then  $\{e_{i_1} e_{i_2} \dots e_{i_t} \mid i_1 < i_2 < \dots < i_t\}$  is a basis of  $\mathrm{Sym}^t(V)$ . Furthermore, if  $G \subset \mathrm{GL}_n(V)$ , then  $G$  acts on  $\mathrm{Sym}^t(V)$  as well via the formula  $\sigma(v_1 \dots v_t) = \sigma(v_1) \dots \sigma(v_t)$ ,  $\sigma \in G$ .

**Definition 1.5.14** (1) *An element  $w \in \mathrm{Sym}^t(V)$  is decomposable if  $w = w_1 \dots w_t$  for some  $w_i \in V$ .*

(2) *A one dimensional subspace  $\ell \subset \mathrm{Sym}^t(V)$  is decomposable if  $\ell$  is spanned by a decomposable vector.*

We note that if we fix a basis  $\{e_1, \dots, e_n\}$  of  $V$  then an element

$$\sum_{i_1 < \dots < i_t} c_{i_1 \dots i_t} e_{i_1} \dots e_{i_t}$$

is decomposable if and only if the  $c_{i_1 \dots i_t}$  satisfy a system of equations called the *Brill equations* ([GKZ94], p. 120,140). Using these definitions, it is not hard to show

**Proposition 1.5.15** (1) *A group  $G \subset \mathrm{GL}_n(V)$  permutes a set of  $t$  one dimensional subspaces if and only if  $G$  leaves a decomposable one dimensional subspace of  $\mathrm{Sym}^t(V)$  invariant.*

(2) *Let  $k, L, K, G$  be as in Proposition 1.5.9 with the order of  $L$  equal to  $n$ . Then  $L(y) = 0$  has a liouvillian solution if and only if for some  $t \leq I(n)$   $G$  leaves invariant a one dimensional subspace of  $\mathrm{Sym}^t(\mathrm{Soln}(L))$ .*

Note that  $\mathrm{Sym}^t(V)$  is constructed using tensor products and quotients. When we apply this construction to differential modules, we get

**Definition 1.5.16** *Let  $k$  be a differential field and  $M$  a differential module over  $k$ . The  $t^{\mathrm{th}}$  symmetric power  $\mathrm{Sym}^t(M)$  of  $M$  is the quotient of the  $t$ -fold tensor product  $M^{\otimes t}$  by the  $k$ -subspace generated by elements  $v_1 \otimes \dots \otimes v_t - v_{\pi(1)} \otimes \dots \otimes v_{\pi(t)}$ , for all  $v_i \in V$  and  $\pi$  a permutation (note that this subspace is a differential submodule as well).*

At the end of Section 1.2, I defined the solution space of a  $k$ -differential module  $M$  in a differential field  $K \supset k$  to be  $\mathrm{Soln}_K(M) = \ker(\partial, K \otimes_k M)$ . From our discussion of tannakian categories, we have that  $\mathrm{Soln}_K(\mathrm{Sym}^t(M)) = \mathrm{Sym}^t(\mathrm{Soln}_K(M))$ . This latter fact suggests the following algorithm to decide if a linear differential equation has a liouvillian solution (we continue to work with scalar equations although everything generalizes easily to systems). Let  $M_L$  be the differential module associated to the equation  $L(y) = 0$ . For each  $t \leq I(n)$

- Calculate  $N_t = \mathrm{Sym}^t(M)$  and find all one dimensional submodules.
- Decide if any of these is decomposable as a subspace of  $N_t$ .

The tannakian formalism implies that a one dimensional submodule of  $N_t$  is decomposable if and only if its solution space is a decomposable subspace  $G$ -invariant subspace of  $\mathrm{Sym}^t(\mathrm{Soln}_K(M_L))$ .

Much work has been done on developing algorithms for these two step. The problem of finding one dimensional submodules of a differential module was essentially solved in the 19<sup>th</sup> and early 20<sup>th</sup> Centuries (albeit in a different language). More recently work of Barkatou, Bronstein van Hoeij, Li, Weil, Wu, Zhang and others has produced good algorithms to solve this problem (see [HW96], [HRUW99], [LSWZ06] and Chapter

4.3.2 of [dPS03] for references). As noted above, the second problem can be solved (in principal) using the Brill equations (see [SU97] and Chapter 4.3.2 of [dPS03] for a fuller discussion). Finally, one can modify the above to produce the liouvillian solutions as well.

Before I explicate these ideas further in the context of second order equations, I will say a few words concerning finding invariant one dimensional subspaces of solution spaces of linear scalar equations.

**Proposition 1.5.17** *Let  $k$  be a differential field with algebraically closed constants  $C$ .  $L \in k[\partial]$ ,  $K$  the Picard-Vessiot extension of  $k$  for  $L(Y) = 0$  and  $G$  the differential Galois group of  $K$  over  $k$ . An element  $z$  in the solution space  $V$  of  $L(Y) = 0$  spans a one dimensional  $G$ -invariant subspace if and only if  $u = z'/z$  is left fixed by  $G$ . In this case,  $\partial - u$  is a right divisor of  $L$ . Conversely if  $\partial - u, u \in k$  is a right divisor of  $L$ , then there exists a solution  $z \in K$  such that  $z'/z = u$ , in which case,  $z$  spans a  $G$ -invariant space.*

*Proof* This follows from Proposition 1.3.28 and its proof applied to first order factors of  $L$ .  $\square$

Therefore, to find one dimensional  $G$ -invariant subspaces of the solution space of  $L(Y) = 0$ , we need to be able to find elements  $u \in k$  such that  $z = e^{\int u}$  satisfies  $L(z) = 0$ . This is discussed in detail in Chapter 4.1 of [dPS03] and I will give a taste of the idea behind a method to do this for  $L \in \mathbb{C}(x)$  assuming  $L$  has only regular singular points.

Let  $u \in \mathbb{C}(x)$  satisfy  $L(e^{\int u}) = 0$  and let  $x = \alpha$  be a pole of  $u$ . The assumption that  $x_0$  is at worst a regular singular point implies that  $e^{\int u}$  is dominated by a power of  $(x - \alpha)$  near  $\alpha$ . We must therefore have that  $u$  has a pole of order at most 1 at  $x = x_0$  and so  $z = e^{\int u} = (x - \alpha)^a h(x)$  where  $a \in \mathbb{C}$  and  $h$  is analytic near  $\alpha \in \mathbb{C}$ . To determine  $a$ , we write

$$L(Y) = Y^{(n)} + \left(\frac{b_{n-1}}{(x-\alpha)} + \text{h.o.t.}\right)Y^{(n-1)} + \dots + \left(\frac{b_0}{(x-\alpha)^{n-1}} + \text{h.o.t.}\right)Y.$$

Substituting  $Y = (x - \alpha)^a(c_0 + c_1(x - \alpha) + \text{h.o.t.})$  and setting the coefficient of  $(x - \alpha)^{a-n}$  equal to zero, we have

$$c_0(a(a-1)\dots(a-(n-1)) + b_{n-1}(a(a-1)\dots(a-(n-2)) + \dots + b_0) = 0$$

or

$$a(a-1)\dots(a-(n-1)) + b_{n-1}(a(a-1)\dots(a-(n-2)) + \dots + b_0 = 0$$



This latter equation is called the *indicial equation* at  $x = \alpha$  and its roots are called the *exponents at  $\alpha$* . If  $\alpha$  is an ordinary point, the  $b_{n-1} = \dots = b_0 = 0$  so  $a \in \{0, \dots, n-1\}$ . One can also define exponents at  $\infty$ . We therefore have that

$$y = \prod_{\alpha_i = \text{finite sing. pt.}} (x - \alpha_i)^{a_i} P(x)$$

where the  $a_i$  are exponents at  $\alpha_i$  and  $-\sum a_i - \deg P$  is an exponent at  $\infty$ . The  $a_i$  and the degree of  $P$  are therefore determined up to a finite set of choices. Note that  $P$  is a solution of  $\tilde{L}(Y) = \prod (x - \alpha_i)^{-a_i} L(\prod (x - \alpha_i)^{a_i} Y)$ . Finding polynomial solutions of  $\tilde{L}(Y) = 0$  of a fixed degree can be done by substituting a polynomial of that degree with undetermined coefficients and equating powers of  $x$  to reduce this to a problem in linear algebra.

**Liouvillian Solutions of Second Order Equations.** The method outlined above can be simplified for second order equations  $L(Y) = Y'' - sY$  because of several facts that we summarize below (see Chapter 4.3.4 of [dPS03]). The resulting algorithm is essentially the algorithm presented by Kovacic in [Kov86] but put in the context of the general algorithm mentioned above. Kovacic's algorithm predated and motivated much of the work on liouvillian solutions of general linear differential equation presented above.

- It can be shown that the fact that no  $Y'$  term appears implies that the differential Galois group must be a subgroup of  $\text{SL}_2$  (Exercise 1.35.5, p. 27 [dPS03]).
- An examination of the algebraic subgroups of  $\text{SL}_2$  implies that  $\text{SL}_2$  has no subgroup of finite index leaving a one dimensional subspace invariant and any proper algebraic subgroup of  $\text{SL}_2$  has a subgroup of index 1, 2, 4, 6, or 12 that leaves a one dimensional subspace invariant (*c.f.*, [Kov86]).
- As shown in the paragraph following Definition 1.5.3, the dimension of the solution space of  $L^{\otimes t}$  is the same as the dimension of  $\text{Sym}^t(\text{Soln}(L))$  and so these two spaces are the same.
- Any element  $z = \sum_{i=0}^t y_1^{t-i} y_2^i \in \text{Sym}^t(\text{Soln}(L))$  can be written as a product  $\prod_{i=0}^t (c_i y_1 + d_i y_2)$  and so all elements of  $\text{Sym}^t(\text{Soln}(L))$  are decomposable.

Combining these facts with the previous results we have

$L(y) = 0$  has a nonzero liouvillian solution

$\Updownarrow$

The differential Galois group  $G$  permutes 1, 2, 4, 6 or 12 one dimensional subspaces of in  $Sol(L)$

$\Updownarrow$

For  $t = 1, 2, 4, 6$ , or  $12$ ,  $G$  leaves invariant a line in  $Sol(L^{\otimes t})$

$\Updownarrow$

For  $t = 1, 2, 4, 6$ , or  $12$ ,  $L^{\otimes t}$  has a solution  $z$  such that  $u = z'/z \in k$

So, to check if  $Y'' - sY = 0$  has a nonzero liouvillian solution (and therefore, by variation of parameters, only liouvillian solutions), one needs to check this last condition.

### Example 1.5.18

$$L(Y) = Y'' + \frac{3(x^2 - x + 1)}{16(x - 1)^2 x^2} Y$$

This is an equation with only regular singular points. One can show that  $L(Y) = 0$  has no nonzero solutions  $z$  with  $z'/z \in \mathbb{C}(x)$ . The second symmetric power is

$$L^{\otimes 2}(Y) = Y'' + 3/4 \frac{(x^2 - x + 1)}{x^2 (x - 1)^2} Y' - 3/8 \frac{(2x^3 - 3x^2 + 5x - 2)}{x^3 (x - 1)^3} Y.$$

The singular points of this equation are at 0, 1 and  $\infty$ . The exponents there are

$$\text{At } 0 : \{1, \frac{1}{2}, \frac{3}{2}\}$$

$$\text{At } 1 : \{1, \frac{1}{2}, \frac{3}{2}\}$$

$$\text{At } \infty : \{-1, -\frac{1}{2}, -\frac{3}{2}\}$$

For  $Y = x^{a_0}(x - 1)^{a_1}P(x)$ , try  $a_0 = \frac{1}{2}, a_1 = 1, \deg P = 0$ . One sees that  $Y = x^{\frac{1}{2}}(x - 1)$  is a solution of  $L^{\otimes 2}(y) = 0$ . Therefore  $L(Y) = 0$  is solvable in terms of liouvillian functions.

For second order equations one can also easily see how to modify the above to find liouvillian solutions when they exist. Assume that one has found a nonzero element  $u \in k$  such that  $L^{\otimes m}(Y) = 0$  has a solution  $z$  with  $z'/z = u$ . We know that

$$z = y_1 \cdots y_m, \quad y_1, \dots, y_m \in \text{Soln}(L).$$

Since  $\sigma(z) \in C \cdot z$  for all  $\sigma \in G$ , we have that for any  $\sigma \in \text{Gal}(L)$ , there exists a permutation  $\pi$  and constants  $c_i$  such that  $\sigma(y_i) = c_i y_{\pi(i)}$ .

Therefore, for  $v_i = \frac{y'_i}{y_i}$  and  $\sigma \in G$  we have  $\sigma(v_i) = v_{\pi(i)}$ , i.e.,  $G$  permutes the  $v_i$ . Let

$$P(Y) = \prod_{i=1}^m (Y - v_i) = Y^m + a_{m-1}Y^{m-1} + \frac{a_{m-2}}{2!}Y^{m-2} + \dots + \frac{a_0}{m!}.$$

The above reasoning implies that the  $a_i \in k$  and

$$a_{m-1} = -(v_1 + \dots + v_m) = -\left(\frac{y'_1}{y_1} + \dots + \frac{y'_m}{y_m}\right) = -\frac{(y_1 \cdot \dots \cdot y_m)'}{y_1 \cdot \dots \cdot y_m} = -\frac{z'}{z} = -u$$

The remaining  $a_i$  can be calculated from  $a_{m-1} = -u$  using the following fact from [BMW97], [Kov86] and [UW96] (see also Chapter 4.3.4 of [dPS03]).

**Lemma 1.5.19** *Using the notation above, we have*

$$a_{i-1} = -a'_i - a_{m-i}a_i - (m-i)(i+1)sa_{i+1}, \quad i = m-1, \dots, 0$$

where  $a_{-1} = 0$ .

In particular we can find a nonzero polynomial satisfied by  $y'/y$  for some nonzero solution of  $L(y) = 0$ . This gives a liouvillian solution and variation of parameters yields another.

**Example 1.5.20** *We continue with the example*

$$L(Y) = Y'' + \frac{3(x^2 - x + 1)}{16(x-1)^2x^2}Y$$

We have that  $y = x^{\frac{1}{2}}(x-1)$  is a soln of  $L^{\otimes 2}(Y) = 0$ . Let

$$a_1 = -\frac{y'}{y} = -\left(\frac{1}{2x} + \frac{1}{x-1}\right) = -\frac{3x-1}{2x(x-1)},$$

then if  $v$  is a root of

$$P(Y) = Y^2 - \frac{3x-1}{2x(x-1)}Y + \frac{9x^2-7x+1}{16x^2(x-1)^2}$$

we have  $y = e^{\int v}$  satisfies  $L(y) = 0$ . This yields

$$\sqrt[4]{x-x^2}\sqrt{1+\sqrt{x}} \text{ and } \sqrt[4]{x-x^2}\sqrt{1-\sqrt{x}}$$

as solutions.  $DGal = D_4 = \text{symmetries of a square}$ .

Another approach to second order linear differential equations was discovered by Felix Klein. It has been put in modern terms by Dwork and Baldassari and made more effective recently by van Hoeij and Weil (see [HW05] for references).

**Solving in Terms of Lower Order Equations.** Another way of defining the notion of “solving in terms of liouvillian functions” is to say that a linear differential equation can be solved in terms of solutions of first order equations  $Y' + aY = b$  and algebraic functions. With this in mind it is natural to make the following definition.

**Definition 1.5.21** *Let  $k$  be a differential field,  $L \in k[\partial]$  and  $K$  the associated Picard-Vessiot extension. The equation  $L(Y) = 0$  is solvable in terms of lower order linear equations if there exists a tower  $k = K_0 \subset K_1 \subset \dots \subset K_m$  with  $K \subset K_m$  and for each  $i, 0 \leq i < m$ ,  $K_{i+1} = K_i \langle t_i \rangle$  ( $K_{i+1}$  is generated as a differential field by  $t_i$  and  $K_i$ ), with*

- $t_i$  algebraic over  $K_i$ , or
- $L_i(t_i) = b_i$  for some  $L_i \in K_i[\partial]$   $\text{ord}(L_i) < \text{ord}(L)$ ,  $b_i \in K_i$

One has the following characterization of this property in terms of the differential Galois group.

**Theorem 1.5.22** ([Sin88]) *Let  $k, L, K$  be as above with  $\text{ord}(L) = n$ . The equation  $L(Y) = 0$  is not solvable in terms of lower order linear equations if and only if the lie algebra of  $\text{DGal}(K/k)$  is simple and has no faithful representations of dimension less than  $n$ .*

If  $\text{ord}(L) = 3$ , then an algorithm to determine if this equation is solvable in terms of lower order linear equations is given in [Sin85]. Refinements of this algorithm and extensions to higher order  $L$  are given in [H02], [H07], [Ngu07], [NdP07], [Per02].

## 1.6 Inverse Problems

In this section we consider the following problem:

*Given a differential field  $k$ , characterize those linear differential algebraic groups that appear as differential Galois groups of Picard-Vessiot extensions over  $k$ .*

We begin with considering the inverse problem over fields of constants.

**Differential Galois groups over  $\mathbb{C}$ .** Let  $L \in \mathbb{C}[\partial]$  be a linear differential operator with constant coefficients. All solutions of  $L(Y) = 0$ ,  $L \in \mathbb{C}[\partial]$  are of the form  $\sum_i P_i(x)e^{\alpha_i x}$  where  $P_i(x) \in \mathbb{C}[x]$ ,  $\alpha_i \in \mathbb{C}$ ,  $x' = 1$ . This implies that the associated Picard-Vessiot  $K$  is a subfield of a field  $E = \mathbb{C}(x, e^{\alpha_1 x}, \dots, e^{\alpha_t x})$ . One can show that  $\text{DGal}(E/k) = (\mathbb{C}, +) \times ((\mathbb{C}^*, \cdot) \times \dots \times (\mathbb{C}^*, \cdot))$ . The group  $\text{DGal}(K/\mathbb{C})$  is a quotient of this latter group and one can show that it therefore must be of the form  $(\mathbb{C}, +)^a \times (\mathbb{C}^*, \cdot)^b$ ,  $a = 0, 1$ ,  $b \in \mathbb{N}$ .

One can furthermore characterize in purely group theoretic term the groups that appear as differential Galois groups over  $\mathbb{C}$ . Note that

$$(\mathbb{C}, +) \simeq \left\{ \begin{pmatrix} 1 & a \\ 0 & 1 \end{pmatrix} \mid a \in \mathbb{C} \right\}$$

and that all these elements are unipotent matrices (i.e.,  $(A - I)^m = 0$  for some  $m \neq 0$ ). This motivates the following definition

**Definition 1.6.1** *If  $G$  is a linear differential algebraic group, the unipotent radical  $R_u$  of  $G$  is the largest normal subgroup of  $G$  all of whose elements are unipotent.*

Note that for a group of the form  $G = \mathbb{C} \times (\mathbb{C}^* \times \dots \times \mathbb{C}^*)$  we have that  $R_u(G) = \mathbb{C}$ . Using facts about linear algebraic groups (see [Hum75] or [Spr98]) one can characterize those linear algebraic groups that appear as differential Galois groups over an algebraically closed field of constants  $C$  as

**Proposition 1.6.2** *A linear algebraic group  $G$  is a differential Galois group of a Picard-Vessiot extension of  $(\mathbb{C}, \partial)$ ,  $\partial c = \forall c \in \mathbb{C}$  if and only if  $G$  is connected, abelian and  $R_u(G) \leq 1$ .*

All of the above holds equally well for any algebraically closed field  $C$ .

**Differential Galois groups over  $\mathbb{C}(x)$ ,  $x' = 1$ .** The inverse problem for this field was first solved by C. and M. Tretkoff, who showed

**Theorem 1.6.3** [TT79] *Any linear algebraic group is a differential Galois group of a Picard-Vessiot extension of  $\mathbb{C}(x)$ .*

Their proof (which I will outline below) depends on the solution of Hilbert's 21<sup>st</sup> Problem. This problem has a weak and strong form. Let  $\mathcal{S} = \{\alpha_1, \dots, \alpha_m, \infty\} \subset S^2$ ,  $\alpha_0 \in S^2 - \mathcal{S}$  and  $\rho : \pi_1(S^2 - \mathcal{S}, \alpha_0) \rightarrow$

$\mathrm{GL}_n(\mathbb{C})$  be a homomorphism.

Weak Form Does there exist  $A \in \mathrm{gl}_n(\mathbb{C}(x))$  such that

- $Y' = AY$  has only regular singular points, and
- The monodromy representation of  $Y' = AY$  is  $\rho$ .

Strong Form Do there exist  $A_i \in \mathrm{GL}_n(\mathbb{C})$  such that the monodromy rep of

$$Y' = \left( \frac{A_1}{x - \alpha_1} + \dots + \frac{A_m}{x - \alpha_m} \right) Y$$

is  $\rho$ .

We know that a differential equation  $Y' = AY$  that has a regular singular point is locally equivalent to one with a simple pole. The strong form of the problem insists that we find an equation that is *globally* equivalent to one with only simple poles. A solution of the strong form of the problem of course yields a solution of the weak form. Many special cases of the strong form were solved before a counterexample to the general case was found by Bolibruch (see [Bea93], [AB94] or Chapters 5 and 6 of [dPS03] for a history and exposition of results and [BMM06] for more recent work and generalizations).

- (i) Let  $\gamma_1, \dots, \gamma_m$  be generators of  $\pi_1(S^2 - \mathcal{S}, \alpha_0)$ , each enclosing just one  $\alpha_i$ . If some  $\rho(\gamma_i)$  is diagonalizable, then the answer is yes. (Plemelj)
- (ii) If all  $\rho(\gamma_i)$  are sufficiently close to  $I$ , the answer is yes. (Lappo-Danilevsky)
- (iii) If  $n = 2$ , the answer is yes. (Dekkers)
- (iv) If  $\rho$  is irreducible, the answer is yes. (Kostov, Bolibruch)
- (v) Counterexample for  $n = 3, |\mathcal{S}| = 4$  and complete characterization for  $n = 3, 4$  (Bolibruch, Gladyshev)

In a sense, the positive answer to the weak form follows from Plemelj's result above - one just needs to add an additional singular point  $\alpha_{m+1}$  and let  $\gamma_{m+1}$  be the identity matrix. A modern approach to give a positive answer to the weak form was given by Röhrl and later Deligne (see [Del70]). We now turn to

*Proof of Theorem 1.6.3* (outline, see [TT79] of Chapter 5.2 or [dPS03] for details) Let  $G$  be a linear algebraic group. One can show that there exist  $g_1, \dots, g_m \in G$  that generate a Zariski dense subgroup  $H$ . Select points  $\mathcal{S} = \{\alpha_1, \dots, \alpha_m\} \subset S^2$  and define  $\rho : \pi_1(S^2 - \mathcal{S}, \alpha_0) \rightarrow H$  via

$\rho(\gamma_i) = g_i$ . Using the solution of the weak form of Hilbert's 21<sup>st</sup>, there exist  $A \in \text{gl}_n(\mathbb{C}(x))$  such that  $Y' = AY$  has regular singular points and monodromy gp  $H$ . Schlesinger's Theorem, Theorem 1.3.27, implies that the differential Galois group of this equation is the Zariski closure of  $H$ .  $\square$

The above result leads to the following two questions:

- What is the minimum number of singular points (not necessarily regular singular points) that a linear differential system must have to realize a given group as its Galois group?
- Can we realize all linear algebraic group as differential Galois groups over  $C(x), x' = 1$  where  $C$  is an arbitrary algebraically closed field?

To answer the first question, we first will consider

**Differential Galois groups over  $\mathbb{C}(\{x\})$ .** To characterize which groups occur as differential Galois groups over this field, we need the following definitions.

**Definition 1.6.4** *Let  $C$  be an algebraically closed field.*

(1) *A torus is a linear algebraic group isomorphic to  $(C^*, \cdot)^r$  for some  $r$ .*

(2) *If  $G$  is a linear algebraic group then we define  $L(G)$  to be the group generated by all tori in  $G$ . This is a normal, Zariski closed subgroup of  $G$  (see [Ram96] and Chapter 11.3 of [dPS03]).*

**Examples 1.6.5** (1) *If  $G$  is reductive and connected (e.g., tori,  $\text{GL}_n, \text{SL}_n$ ), then  $L(G) = G$  (A group  $G$  is reductive if  $R_u(G) = \{I\}$ ).*

(2) *If  $G = (C, +)^r, r \geq 2$ , then  $L(G) = \{I\}$ .*

Ramis [Ram96] showed the following (see also Chapter 11.4 of [dPS03]).

**Theorem 1.6.6** *A linear algebraic group  $G$  is a differential Galois group of a Picard-Vessiot extension of  $\mathbb{C}(\{x\})$  if and only if  $G/L(G)$  is the Zariski closure of a cyclic group.*

Any linear algebraic group  $G$  with  $G/G^0$  cyclic and  $G$  reductive satisfies these criteria while  $G = (C, +)^r, r \geq 2$  does not. Ramis showed that his characterization of local Galois groups in terms of formal monodromy, exponential torus and Stokes matrices yields a group of this type and conversely any such group can be realized as a local Galois group. Ramis

was furthermore able to use analytic patching techniques to get a global version of this result.

**Theorem 1.6.7** *A linear algebraic group  $G$  is a differential Galois group of a Picard-Vessiot extension of  $\mathbb{C}(x)$  with at most  $r - 1$  regular singular points and one (possibly) irregular singular point if and only if  $G/L(G)$  is the Zariski closure of a group generated by  $r - 1$  elements.*

**Examples 1.6.8** (1) *We once again have that any linear algebraic group is a differential Galois group over  $\mathbb{C}(x)$  since such a group satisfies the above criteria for some  $r$ .*

(2)  $\mathrm{SL}_n$  can be realized as a differential Galois group of a linear differential equation over  $\mathbb{C}(x)$  with only one singular point.

(3) One needs  $r$  singular points to realize the group  $(\mathbb{C}, +)^{r-1}$  as a differential Galois group over  $\mathbb{C}(x)$ .

**Differential Galois groups over  $C(x)$ ,  $C$  an algebraically closed**

**field.** The proofs of the above results depend on analytic techniques over  $\mathbb{C}$  that do not necessarily apply to general algebraically closed fields  $C$ . The paper [Sin93] examines the question of the existence of “transfer principals” and shows that for certain groups  $G$ , the existence of an equation over  $\mathbb{C}(x)$  having differential Galois group  $G$  over implies the existence of an equation over  $C(x)$  having differential Galois group  $G$  over  $C(x)$  but these results do not apply to all groups. Algebraic proofs for various groups over  $C(x)$  (in fact over any differential field finitely generated over its algebraically closed field of constants) have been given over the years by Bialynicki-Birula (nilpotent groups), Kovacic (solvable groups), Mitschi/Singer (connected groups) (see [MS96] for references). Finally, Hartmann [Har05] showed that any linear algebraic group can be realized over  $C(x)$  (see also [Obe03]). An exciting recent development is the announcement of a new proof by Harbater and Hartmann [Har07a, Har07c] of this fact based on a generalization of algebraic patching techniques. I will give two examples of a constructive technique (from [MS96]) that allows one to produce explicit equations for connected linear algebraic groups. This technique is based on (see Proposition 1.31 of [dPS03])

**Proposition 1.6.9** *Let  $Y' = AY$  be a differential equation over a differential field  $k$  with algebraically closed constants  $C$ . Let  $G, H$  be linear algebraic subgroups of  $\mathrm{GL}_n(C)$  with Lie algebras  $\mathcal{G}, \mathcal{H}$ . Assume  $G$  is con-*



nected.

1) Let  $Y' = AY$  is a differential equation with  $A \in \mathcal{G} \otimes_C k$  and assume that  $G$  is connected. Then the differential Galois group of this equation is conjugate to a subgroup of  $G$ .

(2) Assume  $k = C(x)$ , that  $A \in \mathcal{G} \otimes_C C(x)$  and that the differential Galois group of  $Y' = AY$  is  $H \subset G$ , where  $H$  is assumed to be connected. Then there exists  $B \in G(C(x))$  such that  $\tilde{A} = B^{-1}AB - B^{-1}B \in \mathcal{H} \otimes_C C(x)$ , that is  $Y' = AY$  is equivalent to an equation  $Y' = \tilde{A}Y$  with  $\tilde{A} \in \mathcal{G} \otimes_C C(x)$  and the equivalence is given by an element of  $G(C(x))$ .

These results allow us to formulate the following strategy to construct differential equations with differential Galois group over  $C(x)$  a given connected group  $G$ . Select  $A \in \mathcal{G} \otimes_C C(x)$  such that

- (i) the differential Galois group of  $Y' = AY$  over  $C(x)$  is connected, and
- (ii) for any proper connected linear algebraic subgroup  $H$  of  $G$ , there is no  $B \in G(C(x))$  such that  $B^{-1}AB - B^{-1}B \in \mathcal{H} \otimes_C C(x)$ .

We will assume for convenience that  $C \subset \mathbb{C}$ . To insure that the differential Galois group is connected, we will select an element  $A \in \mathcal{G} \otimes_C C[x]$ , that is a matrix with *polynomial entries*. This implies that there is a fundamental solution matrix whose entries are entire functions. Since these functions and their derivatives generate the associated Picard-Vessiot extension  $K$ , any element of  $K$  will be a function with at worst poles on the complex plane. If  $E, C(x) \subset E \subset K$  is the fixed field of  $G^0$ , then the elements of  $E$  are algebraic functions that can only be ramified at the singular points of  $Y' = AY$ , that is, only at  $\infty$ . But such an algebraic function must be rational so  $E = C(x)$  and therefore  $G = G^0$ .

The next step is to select an  $A \in \mathcal{G} \otimes_C C[x]$  satisfying condition (ii) above. This is the heart of [MS96] and we shall only give two examples.

**Example 1.6.10**  $G = C^* \times C^* =$

$$\left\{ \begin{pmatrix} a & 0 \\ 0 & b \end{pmatrix} \mid ab \neq 0 \right\}$$

If  $\mathcal{G}$  is the lie algebra of  $G$  then  $\mathcal{G} \otimes_C C[x] =$

$$\left\{ \begin{pmatrix} f_1 & 0 \\ 0 & f_2 \end{pmatrix} \mid f_1, f_2 \in C[x] \right\}$$

The proper closed subgroups of  $G$  are of the form

$$G_{m,n} = \left\{ \begin{pmatrix} c & 0 \\ 0 & d \end{pmatrix} \mid c^n d^m = 1 \right\}$$

$m, n$  integers, not both zero. The connected subgroups are those  $G_{m,n}$  with  $m, n$  relatively prime. If  $\mathcal{G}_{m,n}$  is the lie algebra of  $G_{m,n}$  then  $\mathcal{G}_{m,n} \otimes_C C(x) =$

$$\left\{ \begin{pmatrix} g_1 & 0 \\ 0 & g_2 \end{pmatrix} \mid mg_1 + ng_2 = 0 \right\}$$

We need to find  $A = \begin{pmatrix} f_1 & 0 \\ 0 & f_2 \end{pmatrix}$ ,  $f_1, f_2 \in C[x]$  such that for any  $B = \begin{pmatrix} u & 0 \\ 0 & v \end{pmatrix}$ ,  $u, v \in C(x)$ ,  $BAB^{-1} - BB^{-1} \notin \mathcal{G}_{m,n} \otimes_C C(x)$ , that is

$$m(f_1 - \frac{u'}{u}) + n(f_2 - \frac{v'}{v}) = 0 \Rightarrow m = n = 0.$$

It is sufficient to find  $f_1, f_2 \in C[x]$  such that

$$mf_1 + nf_2 = \frac{h'}{h} \text{ for some } h \in C(x) \Rightarrow m = n = 0$$

Selecting  $f_1 = 1$  and  $f_2 = \sqrt{2}$  will suffice so

$$Y' = \begin{pmatrix} 1 & 0 \\ 0 & \sqrt{2} \end{pmatrix} Y$$

has Galois group  $G$ .

**Example 1.6.11**  $G = \mathrm{SL}_2$ . Its lie algebra is  $\mathfrak{sl}_2 = \text{trace } 0 \text{ matrices}$ . Let

$$A = A_0 + xA_1 = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} + x \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$$

It can be shown that any proper subalgebra of  $\mathfrak{sl}_2$  is solvable and therefore leaves a one dimensional subspace invariant. A calculation shows that no  $U'U^{-1} + UAU^{-1}$  leaves a line invariant (see [MS96]) for details) and so this equation must have Galois group  $\mathrm{SL}_2$ .

This latter example can be generalized to any semisimple linear algebraic group  $G$ . Such a group can be realized as the differential Galois group of an equation of the form  $Y' = (A_0 + xA_1)Y$  where  $A_0$  is a sum of generators of the root spaces and  $A_1$  is a sufficiently general element of the Cartan subalgebra. In general, we have

**Theorem 1.6.12** *Let  $G$  be a connected lin alg gp. defined over alg closed field  $C$ . One can construct  $A_i \in \mathfrak{gl}_n(C)$  and  $A_\infty \in C[x]$  such that for distinct  $\alpha_i \in C$*

$$Y' = \left( \frac{A_1}{x - \alpha_1} + \dots + \frac{A_d}{x - \alpha_d} + A_\infty \right) Y$$

*has Galois group  $G$ .*

Furthermore, the number  $d$  in the above result coincides with the number of generators of a Zariski dense subgroup of  $G/L(G)$  and the degree of the polynomials in  $A_\infty$  can be bounded in terms of the groups as well. For non-connected groups [MS02] gives a construction to realize solvable-by-finite groups and [CMS05] gives a construction to realize certain semisimple-by-finite groups. Finally, many linear algebraic groups can be realized as differential Galois groups of members of classical families of differential equations [BBH88, BH89, Kat87b, Kat90, Kat96, DM89, Mit89, Mit96].

To end this section, I will mention a general inverse problem. We know that any Picard-Vessiot ring is the coordinate ring of a torsor for the differential Galois group. One can ask if every coordinate ring of a torsor for a linear algebraic group can be given the structure of a Picard-Vessiot ring. To be more precise, let  $k$  be a differential field with derivation  $\partial$  and algebraically closed constants  $C$ . Let  $G$  be a linear algebraic group defined over  $C$  and  $V$  a  $G$ -torsor defined over  $k$ .

*Does there exist a derivation on  $R = k[V]$  extending  $\partial$  such that  $R$  is a Picard-Vessiot ring with differential Galois group  $G$  and such that the Galois action of  $G$  on  $R$  corresponds to the action induced by  $G$  on the torsor  $V$ ?*

When  $k = C(x)$ , this has can be answered affirmatively due to the work of [Har02]. For a general  $k$ , finitely generated over  $C$ , I do not know the answer. For certain groups and fields, Juan and Ledet have given positive answers, see [Jua07, JL07b, JL07a, JL07c].

Finally, the inverse problem over  $\mathbb{R}(x)$  has been considered by [Dyc07].

## 1.7 Families of Linear Differential Equations

In this section we will consider a family of parameterized linear differential equations and discuss how the differential Galois group depends on

the parameter. I will start by describing the situation for polynomials and the usual Galois groups. Let  $C$  be a field and  $\overline{C}$  its algebraic closure

Let  $G \subset \mathcal{S}_m$  = fixed group of permutations and  $\mathcal{P}(n, m, G) =$

$$\{P = \sum_{i=0}^n \sum_{j=0}^m a_{i,j} x^i y^j \mid a_{i,j} \in C, \text{Gal}(P/C(x)) = G\} \subset C^{(n+1)(m+1)}$$

To describe the structure of  $\mathcal{P}(n, m, G)$ , we need the following definition

**Definition 1.7.1** A set  $S \subset \overline{C}^N$  is  $C$ -constructible if it is the finite union of sets

$$\{a \in \overline{C}^N \mid f_1(a) = \dots = f_t(a) = 0, g(a) \neq 0\}$$

where  $f_i, g \in C[X_1, \dots, X_N]$ .

For example, any Zariski closed set defined over  $C$  is  $C$ -constructible and a subset  $S \subset \overline{C}$  is  $\overline{C}$ -constructible if and only if it is finite or cofinite. In particular,  $\mathbb{Q}$  is not  $\mathbb{C}$ -constructible in  $\mathbb{C}$ . We have the following results (see [dDR79] and the references given there)

**Theorem 1.7.2**  $\mathcal{P}(m, n, G)$  is  $\mathbb{Q}$ -constructible.

From this we can furthermore deduce

**Corollary 1.7.3** Any finite group  $G$  is a Galois group over  $\overline{\mathbb{Q}}(x)$ .

*Proof* (Outline) For  $G \subset \mathcal{S}_m$ , one can construct an  $m$ -sheeted normal covering of the Riemann Sphere with this as the group of deck transformations, [Tre71]. The Riemann Existence Theorem implies that this Riemann Surface is an algebraic curve and that the the Galois group of its function field over  $\mathbb{C}(x)$  is  $G$ . Therefore for some  $n$ ,  $\mathcal{P}(n, m, G)$  has a point in  $C^{(n+1)(m+1)}$ . Hilbert's Nullstellensatz implies it will have a point in  $\overline{\mathbb{Q}}^{(n+1)(m+1)}$  and therefore that  $G$  is a Galois group over  $\overline{\mathbb{Q}}(x)$ .  $\square$

### Differential Galois Groups of Families of Linear Differential

**Equations.** I would like to apply the same strategy to linear differential equations. We begin by fixing a linear algebraic group  $G \subset \text{GL}_m$  and set  $\mathcal{L}(m, n, G) =$

$$\{L = \sum_{i=0}^n \sum_{j=0}^m a_{i,j} x^i \partial^j \mid a_{i,j} \in \mathbb{C}, \text{Gal}(L/\mathbb{C}(x)) = G\} \subset \mathbb{C}^{(n+1)(m+1)}$$

Regrettably, this set is not necessarily a  $\mathbb{Q}$  or even a  $\mathbb{C}$ -constructible set.

**Example 1.7.4** We have seen that the Galois group of  $\frac{dy}{dx} - \frac{\alpha}{x}y$  is  $\mathrm{GL}_1(\mathbb{C})$  if and only if  $\alpha \notin \mathbb{Q}$ . If  $\mathcal{L}(1, 1, \mathrm{GL}_1)$  were  $\mathbb{C}$ -constructible then

$$\mathcal{L}(1, 1, \mathrm{GL}_1) \cap \{x\partial - \alpha \mid \alpha \in \mathbb{C}\} = \mathbb{C} \setminus \mathbb{Q}$$

would be constructible. Therefore, even the set  $\mathcal{L}(1, 1, \mathrm{GL}_1)$  is not  $\mathbb{C}$ -constructible.

**Example 1.7.5** Consider the family

$$\frac{d^2y}{dx^2} - (\alpha_1 + \alpha_2)\frac{dy}{dx} + \alpha_1\alpha_2y = 0 \quad G = \mathbb{C}^* \oplus \mathbb{C}^*$$

This equation has solution  $\{e^{\alpha_1 x}, e^{\alpha_2 x}\}$ . Assume  $\alpha_2 \neq 0$ . If  $\alpha_1/\alpha_2 \notin \mathbb{Q}$  then  $D\mathrm{Gal} = \mathbb{C}^* \oplus \mathbb{C}^*$  while if  $\alpha_1/\alpha_2 \in \mathbb{Q}$  then  $D\mathrm{Gal} = \mathbb{C}^*$ . Therefore,  $\mathcal{L}(2, 0, \mathbb{C}^* \oplus \mathbb{C}^*)$  is not  $\mathbb{C}$ -constructible.

Recall that for  $L \in \mathbb{C}(x)[\partial]$  and  $x_0$  a singular point  $L(y) = 0$  has a fundamental set of solutions:

$$y_i = (x - \alpha)^{\rho_i} e^{P_i(1/t)} \left( \sum_{j=0}^{s_i} b_{ij} (\log(x - x_0))^j \right)$$

where  $\rho_i \in \mathbb{C}$ ,  $t^\ell = (x - x_0)$  for some  $\ell \leq n!$ ,  $P_i$  are polynomials without constant term,  $b_{ij} \in \mathbb{C}[[x - x_0]]$ .

**Definition 1.7.6** The set  $\mathcal{D}_{x_0} = \{\rho_1, \dots, \rho_n, P_1, \dots, P_n\}$  is called the local data of  $L$  at  $x_0$ .

In Example 1.7.4, 0 and  $\infty$  are the singular points and at both of these  $\{\alpha\}$  is the local data. In Example 1.7.5,  $\infty$  is the only singular point and the local data is  $\{P_1 = \frac{\alpha_1}{x}, P_2 = \frac{\alpha_2}{x}\}$ . In both cases, the parameters allow us to vary the local data. One would hope that by fixing the local data, parameterized families of linear differential equations with fixed Galois group  $G$  would be constructible. This turns out to be true for many *but not all* groups  $G$ .

**Definition 1.7.7** Let  $\mathcal{D} = \{\rho_1, \dots, \rho_r, P_1, \dots, P_s\}$  be a finite set with  $\rho_i \in \mathbb{C}$  and  $P_i$  polynomials without constant terms. We define

$$\mathcal{L}(m, n, \mathcal{D}, G) = \{L \in \mathcal{L}(m, n, G) \mid \mathcal{D}_a \subset \mathcal{D} \text{ for all sing pts } a \in S^2\}$$

Note that we do not fix the position of the singular points; we only fix the local data at putative singular points. Also note that  $\mathcal{L}(m, n, \mathcal{D}) = \{L = \sum_{i=0}^n \sum_{j=0}^m a_{i,j} x^i \partial^j \mid a_{i,j} \in \mathbb{C}, \mathcal{D}_{x_0} \subset \mathcal{D}, \text{ for all sing pts } x_0\}$  is a constructible set. To describe the linear algebraic groups for which  $\mathcal{L}(m, n, \mathcal{D}, G)$  is a constructible set, we need the following definitions.

**Definition 1.7.8** *Let  $G$  a linear algebraic group defined over  $C$ , an algebraically closed field and let  $G^0$  be the identity component.*

- 1) *A character  $\chi$  is a polynomial homomorphism  $\chi : G \rightarrow C^*$ .*
- 2)  *$\text{Ker}X(G^0)$  is defined to be the intersection of kernels of all characters  $\chi : G^0 \rightarrow C^*$ .*

**Examples 1.7.9** 1) *If  $G$  is finite, then  $\text{Ker}X(G^0)$  is trivial.*

2) *If  $G^0$  is semisimple or unipotent, then  $\text{Ker}X(G^0) = G^0$ .*

One can show [Sin93] that  $\text{Ker}X(G^0)$  is the smallest normal subgroup such that  $G^0/\text{Ker}X(G^0)$  is a torus. In addition it is the subgroup generated by all unipotent elements of  $G^0$ . Furthermore,  $\text{Ker}X(G^0)$  is normal in  $G$  and we have

$$1 \rightarrow G^0/\text{Ker}X(G^0) \rightarrow G/\text{Ker}X(G^0) \rightarrow G/G^0 \rightarrow 1$$

Finally,  $G^0/\text{Ker}X(G^0)$  is abelian, so we have that  $G/G^0$  acts on  $G^0/\text{Ker}X(G^0)$ .

**Theorem 1.7.10** [Sin93] *Assume that the action of  $G/G^0$  on  $G^0/\text{Ker}X(G^0)$  is trivial. Then  $\mathcal{L}(m, n, \mathcal{D}, G)$  is constructible.*

**Examples 1.7.11** *The groups  $G$  that satisfy the hypothesis of this theorem include all finite groups, all connected groups and all groups that are either semisimple or unipotent. An example of a group that does not satisfy the hypothesis is the group  $\mathbb{C}^* \rtimes \{1, -1\}$  where  $-1$  sends  $c$  to  $c^{-1}$ .*

For groups satisfying the hypothesis of this theorem we can prove a result similar to Corollary 1.7.3.

**Corollary 1.7.12** *Let  $G$  be as in Theorem 1.7.10, defined over  $\overline{\mathbb{Q}}$ . Then  $G$  can be realized as a differential Galois group over  $\overline{\mathbb{Q}}(x)$ .*

*Proof* (Outline) Let  $G$  be as in Theorem 1.7.10, defined over  $\overline{\mathbb{Q}}$ . We know that  $G$  can be realized as the differential Galois group of a linear

differential equation over  $\mathbb{C}(x)$ , *e.g.*, Theorem 1.6.3. A small modification of Theorem 1.6.3 allows one to assume that the local data of such an equation is defined over  $\overline{\mathbb{Q}}$  (see [Sin93]). Therefore for some  $n, m, \mathcal{D}$ ,  $\mathcal{L}(m, n, \mathcal{D}, G)$  has a point in  $\mathbb{C}^{(n+1)(m+1)}$ . The Hilbert Nullstellensatz now implies that  $\mathcal{L}(m, n, \mathcal{D}, G)$  has a point in  $\overline{\mathbb{Q}}^{(n+1)(m+1)}$ . Therefore  $G \subset \mathrm{GL}_m$  is a differential Galois gp over  $\overline{\mathbb{Q}}(x)$ .  $\square$

The above proof ultimately depends on analytic considerations. We again note that Harbater and Hartman [Har07a, Har07c] have given a direct algebraic proof showing all linear algebraic groups defined over  $\overline{\mathbb{Q}}$  can be realized as a differential Galois group over  $\overline{\mathbb{Q}}(x)$ .

The hypotheses of Theorem 1.7.10 are needed. In fact one can construct (see [Sin93], p. 384-385)) a two-parameter family of second order linear differential equations  $L_{a,t}(Y) = 0$  such that

- $L_{a,t}(Y) = 0$  has 4 regular singular points  $0, 1, \infty, a$  and exponents independent of the parameters  $a, t$ ,
- for all values of the parameters, the differential Galois group is a subgroup of  $\mathbb{C}^* \rtimes \{1, -1\}$  where  $-1$  sends  $c$  to  $c^{-1}$ .
- there is an elliptic curve  $\mathcal{C} : y^2 = f(x)$ ,  $f$  cubic with  $(f, f') = 1$  such that if  $(a, y(a))$  is a point of finite order on  $\mathcal{C}$  then there exists a unique  $t$  such that the differential Galois group of  $L_{a,t}(Y) = 0$  is finite and conversely, the differential Galois group being finite implies that  $(a, y(a))$  is a point of finite order on  $\mathcal{C}$ .

One already sees in the Lamé equation

$$L_{n,B,e}(y) = f(x)y'' + \frac{1}{2}f'(x)y' - (n(n+1)x + B)y = 0$$

where  $f(x) = 4x(x-1)(x-e)$ , that the finiteness of the differential Galois group depends on relations between the fourth singular point  $e$  and the other parameters. For example, Brioschi showed that if  $n + \frac{1}{2} \in \mathbb{Z}$ , then there exists a  $P \in \mathbb{Q}[u, v]$  such that  $L_{n,B,e}(y) = 0$  has finite differential Galois group if and only if  $P(e, B) = 0$ . Algebraic solutions of the Lamé equation are a continuing topic of interest, see [Bal81, Bal87, Dwo90, Lit02, Mai04, BvdW04, LvdPU05].

Finally, the results of [Sin93] have been recast and generalized in [Ber02, Ber04]. In particular the conditions of Theorem 1.7.10 are shown to be necessary and sufficient and existence and properties of moduli spaces are examined (see also Chapter 12 of [dPS03]).

**Parameterized Picard-Vessiot Theory.** In the previous paragraphs, I examined how differential Galois groups depend algebraically on parameters that may appear in families of differential equations. Now I will examine the differential dependence and give an introduction to [CS06] where a Galois theory is developed that measures this. To make things concrete, let us consider parameterized differential equations of the form

$$\frac{\partial Y}{\partial x} = A(x, t_1, \dots, t_m)Y \quad A \in \text{gl}_n(\mathbb{C}(x, t_1, \dots, t_m)). \quad (1.4)$$

If  $x = x_0, t_1 = \tau_1, \dots, t_m = \tau_m$  is a point where the entries of  $A$  are analytic, standard existence theory yields a solution  $Y = (y_{i,j}(x, t_1, \dots, t_m))$ , with  $y_{i,j}$  analytic near  $x = x_0, t_1 = \tau_1, \dots, t_m = \tau_m$ . Loosely speaking, we want to define a Galois group that is the group of transformations that preserve all algebraic relations among  $x, t_1, \dots, t_m$ , the  $y_{i,j}$  and the derivatives of the  $y_{i,j}$  and their derivatives with respect to *all the variables*  $x, t_1, \dots, t_m$ .

To put this in an algebraic setting, we let  $k = \mathbb{C}(x, t_1, \dots, t_m)$  be a (partial) differential field with derivations  $\Delta = \{\partial_0, \partial_1, \dots, \partial_m\}$ ,  $\partial_0 = \frac{\partial}{\partial x}$ ,  $\partial_i = \frac{\partial}{\partial t_i}$   $i = 1, \dots, m$ . We let

$$K = k(y_{1,1}, \dots, y_{n,n}, \dots, \partial_1^{n_1} \partial_2^{n_2} \dots \partial_m^{n_m} y_{i,j}, \dots) = k\langle y_{1,1}, \dots, y_{n,n} \rangle_\Delta.$$

Note that the fact that the  $y_{i,j}$  appear as entries of a solution of (1.4) implies that this field is stable under the derivation  $\partial_0$ . We shall define the parameterized Picard-Vessiot group (PPV-group)  $\text{DGal}_\Delta(K/k)$  to be the group of  $k$ -automorphisms  $\sigma$  of  $K$  that commute with all  $\partial_i$ ,  $i = 0, \dots, m$ .

**Example 1.7.13** Let  $n = 1, m = 1, k = \mathbb{C}(x, t), \Delta = \Delta = \{\partial_x, \partial_t\}$  and

$$\frac{\partial y}{\partial x} = \frac{t}{x}y.$$

This equation has  $y = x^t = e^{t \log x}$  as a solution. Differentiating with respect to  $t$  and  $x$  shows that  $K = \mathbb{C}(x, t, x^t, \log x)$ . Let  $\sigma \in \text{DGal}_\Delta(K/k)$ .

Claim 1:  $\sigma(x^t) = ax^t$ ,  $a \in K$ ,  $\partial_x(a) = 0$  (and so  $a \in \mathbb{C}(t)$ ). To see this note that  $\partial_x \sigma(x^t) = \frac{t}{x} \sigma(x^t)$  so  $\partial_x(\sigma(x^t)/x^t) = 0$ .

Claim 2:  $\sigma(\log x) = \log x + c$ ,  $c \in \mathbb{C}$ . To see this, note that  $\partial_x \sigma(\log x) = \frac{1}{x}$  so  $\partial_x(\sigma(\log x) - \log x) = 0$ . Since  $\partial_t(\log x) = 0$  we also have that  $\partial_t(\sigma(\log x) - \log x) = 0$ .

Claim 3:  $\sigma(x^t) = ax^t, \partial_x a = 0, \partial_t a = ca$ ,  $c \in \mathbb{C}$ . To see this note



that  $\sigma(\partial_t x^t) = \sigma(\log x \ x^t) = (\log x + c)ax^t = a \log x \ x^t + cax^t$  and  $\partial_t \sigma(x^t) = \partial_t(ax^t) = (\partial_t a)x^t + a \log x \ x^t$ .

Therefore,

$$\begin{aligned} \mathrm{DGal}_\Delta(K/k) &= \{a \in k^* \mid \partial_x a = 0, \partial_t(\frac{\partial_t a}{a}) = 0\} \\ &= \{a \in k^* \mid \partial_x a = 0, \partial_t^2(a)a - (\partial_t(a))^2 = 0\} \end{aligned}$$

Remarks: (1) Let  $k_0 = \ker \partial_x = \mathbb{C}(t)$ . This is a  $\partial_t$ -field. The PPV-group  $\mathrm{DGal}_\Delta(K/k)$  can be identified with a certain subgroup of  $\mathrm{GL}_1(k_0)$ , that is, with a group of matrices whose entries satisfy a differential equations. This is an example of a linear differential algebraic group (to be defined more fully below).

(2) In fact, using partial fraction decompositions, one can see that

$$\{a \in k^* \mid \partial_x a = 0, \partial_t(\frac{\partial_t a}{a}) = 0\} = \mathbb{C}^*$$

This implies that for any  $\sigma \in \mathrm{DGal}_\Delta(K/k)$ ,  $\sigma(\log x) = \log x$ . If we want a Fundamental Theorem for this new Galois theory, we will want to have enough automorphisms so that any element  $u \in K \setminus k$  is moved by some automorphism. This is not the case here.

(3) If we specialize  $t$  to be an element  $\tau \in \mathbb{C}$ , we get a linear differential equation whose monodromy group is generated by the map  $y \mapsto e^{2\pi\tau}y$ . One can think of the map  $y \mapsto e^{2\pi\tau}y$  as the “parameterized monodromy map”. I would like to say that this map lies in the PPV-group and that this map generates a group that is, in some sense, dense in the PPV-group. We cannot do this yet.

Remarks (2) and (3) indicate that the PPV group does not have enough elements. When we considered the Picard-Vessiot theory, we insisted that the constants be algebraically closed. This guaranteed that (among other things) the differential Galois group (defined by polynomial equations) had enough elements. Since our PPV-groups will be defined by differential equations, it seems natural to require that the field of  $\partial_0$ -constants is “differentially closed”. More formally,

**Definition 1.7.14** Let  $k_0$  be a differential field with derivations  $\Delta_0 = \{\partial_1, \dots, \partial_m\}$ . We say that  $k_0$  is differentially closed if any system of polynomial differential equations

$$f_1(y_1, \dots, y_n) = \dots = f_r(y_1, \dots, y_n) = 0, g(y_1, \dots, y_n) \neq 0$$

with coefficients in  $k_0$  that has a solution in some  $\Delta_0$ -differential extension field has a solution in  $k_0$ .

Differentially closed fields play the same role in differential algebra as algebraically closed fields play in algebraic geometry and share many (but not all) of the same general properties and, using Zorn's Lemma, they are not hard to construct. They have been studied by Kolchin (under the name of constrained closed differential fields [Kol74]) and by many logicians [Mar00]. We need two more definitions before we state the main facts concerning parameterized Picard-Vessiot theory.

**Definition 1.7.15** Let  $k_0$  be a differential field with derivations  $\Delta_0 = \{\partial_1, \dots, \partial_m\}$ .

(1) A set  $X \subset k_0^n$  is Kolchin closed if it is the zero set of a system of differential polynomials over  $k_0$ .

(2) A linear differential algebraic group is a subgroup  $G \subset \mathrm{GL}_n(k_0)$  that is Kolchin closed in  $\mathrm{GL}_n(k_0) \subset k_0^{n^2}$ .

Kolchin closed sets form the closed sets of a topology called the Kolchin topology. I will therefore use topological language (open, dense, etc) to describe their properties. Examples and further description of linear differential algebraic groups are given after the following definition and result.

**Definition 1.7.16** Let  $k$  be a differential field with derivations  $\Delta = \{\partial_0, \partial_1, \dots, \partial_m\}$  and let

$$\partial_0 Y = AY, \quad A \in \mathrm{gl}_n(k) \quad (1.5)$$

A Parameterized Picard-Vessiot extension (PPV-extension) of  $k$  for (1.5) is a  $\Delta$ -ring  $R \supset k$  such that  $R$  is a simple  $\Delta$ -ring (i.e., its only  $\Delta$ -ideals are  $(0)$  and  $R$ ) and  $R$  is generated as a  $\Delta$ -ring by the entries of some  $Z \in \mathrm{GL}_n(R)$  and  $\frac{1}{\det Z}$  where  $\partial_0 Z = AZ$ . A PPV-field is defined to be the quotient field of a PPV-ring.

As in the usual Picard-Vessiot theory, such rings exist and are domains. Under the assumption that  $k_0 = \ker \partial_0$  is a differentially closed  $\Delta_0 = \{\partial_1, \dots, \partial_m\}$ -field, PPV-rings can be shown to be unique (up to appropriate isomorphism).

**Theorem 1.7.17** Let  $k$  be a differential field with derivations  $\Delta = \{\partial_0, \partial_1, \dots, \partial_m\}$  and assume that  $k_0 = \ker \partial_0$  is a differentially closed

$\Delta_0 = \{\partial_1, \dots, \partial_m\}$ -field. Let  $K$  be the PPV-field for an equation

$$\partial_0 Y = AY, \quad A \in \mathfrak{gl}_n(k).$$

(1) The group  $\mathrm{DGal}_\Delta(K/k)$  (called the PPV-group) of  $\Delta$ -differential  $k$ -automorphisms of  $K$  has the structure of a linear differential algebraic group.

(2) There is a Galois correspondence between  $\Delta$ -subfields  $E$  of  $K$  containing  $k$  and Kolchin  $\Delta_0$  closed subgroups  $H$  of  $\mathrm{DGal}_\Delta(K/k)$  with normal subgroups  $H$  corresponding to field  $K$  that are again PPV-extensions of  $k$ .

In the context of this theorem we can recast the above example. Let  $k_0, \mathbb{C}(t) \subset k_0$  be a differentially closed  $\partial_t$ -field and  $k = k_0(x)$  be a  $\Delta = \{\partial_x, \partial_t\}$  field where  $\partial_x(k_0) = 0, \partial_x(x) = 1, \partial_t(x) = 0$ . One can show that the field  $K = k \langle x^t \rangle_\Delta = k(x^t, \log)$  is a PPV-field for the equation  $\partial_x Y = \frac{t}{x} Y$  with PPV group as described above. Because  $k_0$  is differentially closed, this PPV group has enough elements so that the Galois correspondence holds. Furthermore the element  $e^{2\pi i t} \in k_0$  is an element of this group and, in fact, generates a Kolchin dense subgroup.

Remarks: (1) In general, let  $\partial_x Y = AY$  is a parameterized equation with  $A \in \mathbb{C}(x, t_1, \dots, t_m)$ . For each value of  $\bar{t}$  of  $t = (t_1, \dots, t_m)$  at which the entries of  $A$  are defined and generators  $\{\gamma_i\}$  of the Riemann Sphere punctured at the poles of  $A(x, \bar{t}_1, \dots, \bar{t}_m)$ , we can define monodromy matrices  $\{M_i(\bar{t})\}$ . For all values of  $t$  in a sufficiently small open set, the entries of  $M_i$  are analytic functions of  $t$  and one can show that these matrices belong to the PPV-group. If the equation  $\partial_x Y = AY$  has only regular singular points, then one can further show that these matrices generate a Kolchin dense subgroup of the PPV group.

(2) One can consider parameterized families of linear partial differential equations as well. Let  $k$  be a  $\Delta = \{\partial_0, \dots, \partial_s, \partial_{s+1}, \dots, \partial_m\}$  and let

$$\partial_i Y = A_i Y, \quad A_i \in \mathfrak{gl}_n(k), \quad i = 0, \dots, s$$

be a system of differential equations with  $\partial_i(A_j) + A_i A_j = \partial_j(A_i) + A_j A_i$ . One can develop a parameterized differential Galois theory for these as well.

(3) One can develop a theory of parameterized liouvillian functions and show that one can solve parameterized linear differential in terms of these if and only if the PPV group is solvable-by-finite.

To clarify and apply the PPV theory, we will describe some facts about linear differential algebraic groups.

Linear differential algebraic groups. These groups were introduced by Cassidy [Cas72]. A general theory of differential algebraic groups was initiated by Kolchin [Kol85] and developed by Buium, Hrushovsky, Pillay, Sit and others (see [CS06] for references). We give here some examples and results. Let  $k_0$  be a differentially closed  $\Delta_0 = \{\partial_1, \dots, \partial_m\}$  differential field.

**Examples 1.7.18** (1) *All linear algebraic groups are linear differential algebraic groups.*

(2) *Let  $C = \cap_{i=1}^m \ker \partial_i$  and let  $G(k_0)$  be a linear algebraic group defined over  $k_0$ . Then  $G(C)$  is a linear differential algebraic group (just add  $\{\partial_1 y_{i,j} = \dots = \partial_m y_{i,j} = 0\}_{i,j=1}^n$  to the defining equations!).*

(3) *Differential subgroups of*

$$G_a(k_0) = (k_0, +) = \left\{ \begin{pmatrix} 1 & z \\ 0 & 1 \end{pmatrix} \mid z \in k \right\}.$$

*The linear differential subgroups of this group are all of the form*

$$G_a^{(L_1, \dots, L_s)} = \{z \in k_0 \mid L_1(z) = \dots = L_s(z) = 0\}$$

*where the  $L_i$  are linear differential operators in  $k_0[\partial_1, \dots, \partial_m]$ . When  $m = 1$  we may always take  $s = 1$ .*

*For example, if  $m = 1$ ,*

$$G_a^\partial = \{z \in k_0 \mid \partial z = 0\} = G_a(C)$$

(4) *Differential subgroups of  $G_m(k_0) = (k_0^*, \times) = \text{GL}_1(k_0)$ . Assume  $m = 1, \Delta_0 = \{\partial\}$  (for the general case, see [Cas72]). The linear differential subgroups are:*

- *finite and cyclic, or*
- $G_m^L = \{z \in k_0^* \mid L(\frac{\partial z}{z}) = 0\}$  *where the  $L$  is a linear differential operator in  $k_0[\partial]$ .*

*For example the PPV group of Example 1.7.13,  $G_m^\partial = \{z \in k_0 \mid \partial(\frac{\partial z}{z}) = 0\}$  is of this form. The above characterization follows from the exactness of*

$$(1) \longrightarrow G_m(C) \longrightarrow G_m(k_0) \xrightarrow{z \mapsto \frac{\partial z}{z}} G_a(k_0) \longrightarrow (0)$$

and the characterization of differential subgroups of  $G_a$ . Note that in the usual theory of linear algebraic groups there is no surjective homomorphism from  $G_m$  to  $G_a$ .

(5) A result of Cassidy [Cas72] states: if  $m = 1$ ,  $\Delta_0 = \{\partial\}$  and  $H$  is a Zariski dense proper differential subgroup of  $\mathrm{SL}_n(k_0)$ , then there is an element  $g \in \mathrm{SL}_n(k_0)$  such that  $gHg^{-1} = \mathrm{SL}_n(C)$ , where  $C = \ker \partial$ . This has been generalized by Cassidy and, later, Buium to the following. Let  $H$  be a Zariski dense proper differential subgroup of  $G$ , a simple linear algebraic group, defined over  $\mathbb{Q}$ . Then there exists  $\mathbb{D} = \{D_1, \dots, D_s\} \subset k\Delta_0 = k - \text{span of } \Delta_0$  such that

- (i)  $[D_i, D_j] = 0$  for  $1 \leq i, j \leq s$  and
- (ii) There exists a  $g \in G(k)$  such that  $gHg^{-1} = G(C_{\mathbb{D}})$ , where  $C_{\mathbb{D}} = \bigcap_{i=1}^s \ker D_i$ .

Inverse Problem. In analogy to the Picard-Vessiot theory, the question I will discuss is: Given a differential field  $k$ , which linear differential algebraic groups appear as PPV-groups over  $k$ ? Even in simple cases the complete answer is not known and there are some surprises.

Let  $\Delta = \{\partial_x, \partial_t\}$ ,  $\Delta_0 = \{\partial_x\}$ ,  $k_0$  a  $\partial_t$ -differentially closed  $\partial_t$ -field and  $k = k_0(x)$  with  $\partial_0(k_0) = 0$ ,  $\partial_x(x) = 1$ , and  $\partial_t(x) = 0$ . Let  $C \subset k_0$  be the  $\Delta$ -constants of  $k$ . I will show that all differential subgroups of  $G_a = (k_0, +)$  appear as PPV-groups over this field but that the full group  $G_a$  does not appear as a PPV-group [CS06].

One can show that the task of describing the differential subgroups of  $G_a$  that appear as PPV-groups is equivalent to describing which linear differential algebraic groups appear as PPV-groups for equations of the form

$$\frac{dy}{dx} = a(t, x), \quad a(t, x) \in k_0(x)$$

Using partial fraction decomposition, one sees that a solution of such an equation is of the form

$$y = R(t, x) + \sum_{i=1}^s a_i(t) \log(x - b_i(t)), \quad R(t, x) \in k_0(x), a_i(t), b_i(t) \in k_0$$

and the associated PPV-extension is of the form  $K = k(y)$ . If  $H$  is the PPV-group of  $K$  over  $k$  and  $\sigma \in H$ , then one shows that

$$\sigma(y) = R(t, x) + \sum_{i=1}^s a_i(t) \log(x - b_i(t)) + \sum_{i=1}^s c_i a_i(t)$$

for some  $c_i \in C$ . Let  $L \in k_0[\frac{d}{dt}]$  have solution space spanned by the  $a_i(t)$  (note that  $L \neq 0$ ). For any  $\sigma \in H$ ,  $\sigma$  can be identified with an element of

$$H = \{z \in G_a(k_0) \mid L(z) = 0.\}$$

Conversely, any element of  $H$  can be shown to induce a differential automorphism of  $K$ , so  $H$  is the PPV-group. Therefore  $H$  must be a proper subgroup of  $G_a$  and not all of  $G_a$ . One can furthermore show that all proper differential algebraic subgroups of  $G_a$  can appear in this way.

A distinguishing feature of proper subgroups of  $G_a$  is that they contain a finitely generated Kolchin dense subgroup whereas  $G_a$  does not have such a subgroup. A possible conjecture is that a linear differential algebraic group  $H$  is a PPV-group over  $k$  if and only if  $H$  has a finitely generated Kolchin dense subgroup. If  $H$  has such a finitely generated subgroup, a parameterized solution of Hilbert's 22<sup>nd</sup> shows that one can realize this group as a PPV-group. If one could establish a parameterized version of Ramis's theorem that the formal monodromy, exponential torus and Stokes generate a dense subgroup of the differential Galois group (something of independent interest), the other implication of this conjecture would be true.

I will now construct a field over which one can realize  $G_a$  as a PPV-group. Let  $k$  be as above and let  $F = k(\log x, x^{t-1}e^{-x})$ . The *incomplete Gamma function*

$$\gamma(t, x) = \int_0^x s^{t-1} e^{-s} ds$$

satisfies

$$\frac{d\gamma}{dx} = x^{t-1} e^{-x}$$

over  $F$ . The PPV-group over  $k$  is  $G_a(k_0)$  because  $\gamma, \frac{d\gamma}{dt}, \frac{d^2\gamma}{dt^2}, \dots$  are algebraically independent over  $k$  [JRR95] and so there are no relations to preserve. Over  $k = k_0(x)$ ,  $\gamma(t, x)$  satisfies

$$\frac{d^2\gamma}{dx^2} - \frac{t-1-x}{x} \frac{d\gamma}{dx} = 0$$

and the PPV-group is

$$\begin{aligned} H &= \left\{ \begin{pmatrix} 1 & a \\ 0 & b \end{pmatrix} \mid a \in k_0, b \in k_0^*, \partial_t \left( \frac{\partial_t b}{b} \right) = 0 \right\} \\ &= G_a(k_0) \rtimes G_m^{\partial_t}, \quad G_m^{\partial_t} = \{ b \in k_0^* \mid \partial_t \left( \frac{\partial_t b}{b} \right) = 0 \} \end{aligned}$$

*Isomonodromic families.* The parameterized Picard-Vessiot theory can be used to characterize isomonodromic families of linear differential equations with regular singular points, that is, families of such equations where the monodromy is independent of the parameters.

**Definition 1.7.19** *Let  $k$  be a  $\Delta = \{\partial_0, \dots, \partial_m\}$ -differential field and let  $A \in \text{gl}_n(k)$ . We say that  $\partial_0 Y = AY$  is completely integrable if there exist  $A_i \in \text{gl}_n(k)$ ,  $i = 0, 1, \dots, m$  with  $A_0 = A$  such that*

$$\partial_j A_i - \partial_i A_j = A_j A_i - A_i A_j \text{ for all } i, j = 0, \dots, m$$

The nomenclature is motivated by the fact that the latter conditions on the  $A_i$  are the usual integrability conditions on the system of differential equations  $\partial_i Y = AY$ ,  $i = 0, \dots, m$ .

**Proposition 1.7.20** [CS06] *Let  $k$  be a  $\Delta = \{\partial_0, \dots, \partial_m\}$ -differential field and let  $A \in \text{gl}_n(k)$ . Assume that  $k_0 = \ker \partial_0$  is  $\Delta_0 = \{\partial_1, \dots, \partial_m\}$ -differentially closed field and let  $K$  be the PPV-field of  $k$  for  $\partial_0 Y = AY$  with  $A \in \text{gl}_n(k)$ . Let  $C = \cap_{i=0}^m \ker \partial_i$ .*

*There exists a linear algebraic group  $G$ , defined over  $C$ , such that  $\text{DGal}_\Delta(K/k)$  is conjugate to  $G(C)$  over  $k_0$  if and only if  $\partial_0 Y = AY$  is completely integrable. If this is the case, then  $K$  is a Picard-Vessiot extension corresponding to this integrable system.*

Let  $k = \mathbb{C}(x, t_1, \dots, t_n)$  and  $A \in \text{gl}_n(k)$ . For  $t = (t_1, \dots, t_m)$  in some sufficiently small open set  $\mathcal{O} \subset \mathbb{C}^n$ , one can select generators  $\{\gamma_i(t_1, \dots, t_m)\}$  of the fundamental group of the Riemann Sphere minus the (parameterized) singular points such that the monodromy matrices  $\{M_i(t_1, \dots, t_m)\}$  depend analytically on the parameters. One says that the differential equation  $\partial_x Y = AY$  is *isomonodromic* if for some sufficiently small open set  $\mathcal{O}$ , the  $M_i$  are independent of  $t$ . In [CS06] we deduce the following corollary from the above result.

**Corollary 1.7.21** *Let  $k, A, \mathcal{O}$  be as above and assume that  $\partial_x Y = AY$  has only regular singular points for  $t \in \mathcal{O}$ . Then  $\partial_x Y = AY$  is isomon-*

odromic if and only if the PPV-group is conjugate to a linear algebraic group  $G(\mathbb{C}) \subset \mathrm{GL}_n(\mathbb{C})$ .

*Second order parameterized equations.* A classification of the linear differential subgroups of  $\mathrm{SL}_2$  allows one to classify parameterized systems of second order linear differential equations. To simplify the discussion we will consider equations of the form

$$\frac{\partial Y}{\partial x} = A(x, t)Y$$

that depend on only one parameter and satisfy  $A \in \mathfrak{sl}_n(\mathbb{C}(x, t))$ .

Let  $H$  be a linear differential algebraic subgroup of  $\mathrm{SL}_2(k_0)$  (where  $k_0$  is a  $\partial_t$ -differentially closed field containing  $\mathbb{C}(t)$  and  $\ker \partial_t = \mathbb{C}$ ) and  $\bar{H}$  its Zariski closure. If  $\bar{H} \neq \mathrm{SL}_2$ , then one can show that  $\bar{H}$  and therefore  $H$  is solvable-by-finite [Kov86, Cas72]. Furthermore, Cassidy's result (Example 1.7.18(3)) states that if  $\bar{H} = \mathrm{SL}_2$  but  $H \neq \mathrm{SL}_2$ , then  $H$  is conjugate to  $\mathrm{SL}_2(\mathbb{C})$ . From this discussion and the discussion in the previous paragraphs one can conclude the following.

**Proposition 1.7.22** *Let  $k$  and  $\frac{\partial Y}{\partial x} = AY$  be as above. Assume that for some small open set of values of  $t$  that this equation has only regular singular points. Then either*

- *the PPV group is  $\mathrm{SL}_2$ , or*
- *the equation is solvable in terms of parameterized exponentials, integrals and algebraics, or*
- *the equation is isomonodromic.*

The notion of ‘solvable in terms of parameterized exponentials, integrals and algebraics’ is analogous to the similar notion in the usual Picard-Vessiot theory and is given a formal definition in [CS06], where it is also shown that this notion is equivalent to the PPV-group being solvable-by-finite.

## 1.8 Final Comments

I have touched on only a few of the aspects of the Galois theory of linear differential equations. I will indicate here some of the other aspects and give pointers to some of the current literature.



Kolchin himself generalized this theory to a theory of *strongly normal* extension fields [Kol76]. The Galois groups in this theory are arbitrary algebraic groups for which Kolchin. Recently Kovacic [Kov03, Kov06] has recast this theory in terms of groups schemes and differential schemes. Umemura [Ume90, Ume96b, Ume96a, Ume99, Ume, Ume04] has developed a Galois theory of differential equations for general nonlinear equation. Instead of Galois groups, Umemura uses Lie algebras to measure the symmetries of differential fields. Malgrange [Mal02, Mal01] has proposed a Galois theory of differential equations where the role of the Galois group is replaced by certain groupoids. This theory has been expanded and applied by Cassale [Cas06a, Cas06b, Cas07]. Pillay [Pil97, Pil98, MP97, Pil02, Pil04] develops a Galois theory where the Galois groups can be arbitrary *differential* algebraic groups and, together with Bertrand, has used these techniques to generalize Ax's work on Schanuel's conjecture (see [Ber07, Ber06]). Landesman [Lan06] has generalized Kolchin's theory of strongly normal extension to a theory of strongly normal parameterized extensions. The theory of Cassidy and myself presented in Section 1.7 can be viewed as a special case of this theory and many of the results of [CS06] can be derived from the theories of Umemura and Malgrange as well.

A Galois Theory of linear *difference* equations has also been developed, initially by Franke ([Fra63] and several subsequent papers), Białynicki-Birula [BB62] and more fully by van der Put and myself [dPS97]. André [And01] has presented a Galois theory that treats both the difference and differential case in a way that allows one to see the differential case as a limit of the difference case. Chatzidakis, Hrushovski and Kamenisky have used model theoretic tools to develop Galois theory of difference equations (see [CH99], [Kam06] and [CHS07] for a discussion of connections with the other theories). Algorithmic issues have been considered by Abramov, Barkatou, Bronstein, Hendriks, van Hoeij, Kauers, Petkovsek, Paule, Schneider, Wilf, M. Wu and Zeilberger and there is an extensive literature on the subject for which I will refer to MathSciNet and the ArXiv. Recently work by André, Di Vizio, Etingof, Hardouin, van der Put, Ramis, Reversat, Sauloy, and Zhang, has been done on understanding the Galois theory of  $q$ -difference equation (see [DV02, DV04, ADV04, Eti95, dPR07, Sau03, RS07, RSZ06, DVRSZ03, Ram07, Har07b]). Fields with both derivations and automorphisms have been studied from a model theoretic point of view by Bustamente [Med05] and work of Abramov, Bronstein, Li, Petkovsek, Wu, Zheng

and myself [AP94, ABP95, BP96, BP94, BLW05, LSWZ06, Wu05] consider algorithmic questions for mixed differential-difference systems. The Picard-Vessiot theory of linear differential equations has been used by Hardouin [Har06] to study the differential properties of solutions of difference equations. A Galois theory, with linear differential algebraic groups as Galois groups, designed to measure the differential relations among solutions of difference equation, has been recently developed and announced in [Sin07].

Finally, I have not touched on the results and enormous literature concerning the hypergeometric equations and their generalizations as well as arithmetic properties of differential equations. For a taste of the current research, I refer the reader to [SST00] and [HUY07] for the former and [ABB04] for the latter.

### Bibliography

- [AB94] D. V. Anosov and A. A. Bolibruch. *The Riemann-Hilbert Problem*. Vieweg, Braunschweig, Wiesbaden, 1994.
- [ABB04] A. Adolphson, F. Baldassarri, P. Berthelot, N. Katz, and F. Loeser, editors. *Geometric aspects of Dwork theory. Vol. I, II*. Walter de Gruyter GmbH & Co. KG, Berlin, 2004. Lectures from the Dwork trimester dedicated to Bernard Dwork held at the University of Padova, Padova, May–July 2001.
- [ABP95] S.A. Abramov, M. Bronstein, and M. Petkovšek. On polynomial solutions of linear operator equations. In A. H. M. Levelt, editor, *Proceedings of the 1995 International Symposium on Symbolic and Algebraic Computation (ISSAC'95)*, pages 290–296. ACM Press, 1995.
- [ADV04] Y. André and L. Di Vizio.  $q$ -difference equations and  $p$ -adic local monodromy. *Astérisque*, 296:55–111, 2004. Analyse complexe, systèmes dynamiques, sommabilité des séries divergentes et théories galoisiennes. I.
- [And01] Y. André. Différentielles non commutatives et théorie de Galois différentielle ou aux différences. *Ann. Sci. École Norm. Sup. (4)*, 34(5):685–739, 2001.
- [AP94] S.A. Abramov and M. Petkovšek. D'Alembert solutions of linear differential and difference equations. In J. von zur Gathen, editor, *Proceedings of the 1994 International Symposium on Symbolic and Algebraic Computation (ISSAC'94)*, pages 169–180. ACM Press, 1994.
- [Aud01] M. Audin. *Les Systèmes Hamiltoniens et leur Intégrabilité*, volume 8 of *Cours Spécialisés*. Société Mathématique de France, Paris, 2001.
- [Bal81] F. Baldassarri. On algebraic solutions of Lamé's differential equation. *J. Differential Equations*, 41(1):44–58, 1981.
- [Bal87] F. Baldassarri. Algebraic solutions of the Lamé equation and torsion of elliptic curves. In *Proceedings of the Geometry Conference (Milan and Gargnano, 1987)*, volume 57, pages 203–213 (1989), 1987.

- [Bal94] W. Balser. *From Divergent Power Series to Analytic Functions*, volume 1582 of *Lecture Notes in Mathematics*. Springer-Verlag, Heidelberg, 1994.
- [BB62] A. Bialynicki-Birula. On the Galois theory of fields with operators. *Amer. J. Math.*, 84:89–109, 1962.
- [BBH88] F. Beukers, D. Brownawell, and G. Heckman. Siegel normality. *Annals of Mathematics*, 127:279 – 308, 1988.
- [Bea93] A. Beauville. Monodromie des systèmes différentiels à pôles simples sur la sphère de Riemann (d’après A. Bolibruch). *Astérisque*, 216:103–119, 1993. Séminaire Bourbaki, No. 765.
- [Ber02] M. Berkenbosch. Moduli spaces for linear differential equations. In *Differential equations and the Stokes phenomenon*, pages 15–33. World Sci. Publishing, River Edge, NJ, 2002.
- [Ber04] M. Berkenbosch. *Algorithms and Moduli Spaces for Differential Equations*. Ph.d. thesis, University of Groningen, 2004.
- [Ber06] D. Bertrand. Schanuel’s conjecture for non-constant elliptic curves over function fields. to appear in *Model Theory and Applications*, Cambridge University Press, 2006.
- [Ber07] D. Bertrand. Schanuel’s conjecture over function fields and differential galois theory (joint with Anand Pillay). to appear Oberwolfach Reports, 2007.
- [Beu92] F. Beukers. Differential Galois Theory. In M. Waldschmidt et al, editor, *From Number Theory to Physics (Les Houches, 1989)*, pages 413–439. Berline, 1992. Springer-Verlag.
- [BH89] F. Beukers and G. Heckman. Monodromy for the hypergeometric function  ${}_nF_{n-1}$ . *Inventiones Mathematicae*, 95:325–354, 1989.
- [BLW05] M. Bronstein, Z. Li, and M. Wu. Picard-Vessiot extensions for linear functional systems. In Manuel Kauers, editor, *Proceedings of the 2005 International Symposium on Symbolic and Algebraic Computation (ISSAC 2005)*, pages 68–75. ACM Press, 2005.
- [BMM06] A. A. Bolibruch, S. Malek, and C. Mitschi. On the generalized Riemann-Hilbert problem with irregular singularities. *Expo. Math.*, 24(3):235–272, 2006.
- [BMW97] M. Bronstein, T. Mulders, and J.-A. Weil. On symmetric powers of differential operators. In W. Küchlin, editor, *Proceedings of the 1997 International Symposium on Symbolic and Algebraic Computation (ISSAC’97)*, pages 156–163. ACM Press, 1997.
- [Bou98] A. Boulanger. Contribution à l’étude des équations linéaires homogènes intégrables algébriquement. *Journal de l’École Polytechnique, Paris*, 4:1 – 122, 1898.
- [BP94] M. Bronstein and M. Petkovsek. On Ore rings, linear operators and factorization. *Programming and Computer Software*, 20(1):14–26, 1994.
- [BP96] M. Bronstein and M. Petkovsek. An introduction to pseudo-linear algebra. *Theoretical Computer Science*, 157:3–33, 1996.
- [BvdW04] F. Beukers and A. van der Waall. Lamé equations with algebraic solutions. *J. Differential Equations*, 197(1):1–25, 2004.
- [Cas72] P. J. Cassidy. Differential algebraic groups. *Amer. J. Math.*, 94:891–954, 1972.
- [Cas06a] G. Casale. Feuilletages singuliers de codimension un, groupoïde de Galois et intégrales premières. *Ann. Inst. Fourier (Grenoble)*, 56(3):735–779, 2006.
- [Cas06b] G. Casale. Irréductibilité de la première équation de Painlevé. *C. R.*

- Math. Acad. Sci. Paris*, 343(2):95–98, 2006.
- [Cas07] G. Casale. The Galois groupoid of Picard-Painlevé VI equation. In *Algebraic, analytic and geometric aspects of complex differential equations and their deformations. Painlevé hierarchies*, RIMS Kôkyûroku Bessatsu, B2, pages 15–20. Res. Inst. Math. Sci. (RIMS), Kyoto, 2007.
- [CH99] Z. Chatzidakis and E. Hrushovski. Model theory of difference fields. *Trans. Amer. Math. Soc.*, 351(8):2997–3071, 1999.
- [CHS07] Z. Chatzidakis, C. Hardouin, and M. F. Singer. On the Definition of Difference Galois Groups. To appear in the Proceedings of the Newton Institute 2005 semester ‘Model theory and applications to algebra and analysis’; also [arXiv:math.CA/0705.2975](#), 2007.
- [CK02] R. C. Churchill and J. J. Kovacic. Cyclic vectors. In *Differential algebra and related topics (Newark, NJ, 2000)*, pages 191–218. World Sci. Publ., River Edge, NJ, 2002.
- [CMS05] W. J. Cook, C. Mitschi, and M. F. Singer. On the constructive inverse problem in differential Galois theory. *Comm. Algebra*, 33(10):3639–3665, 2005.
- [Cop34] F. Cope. Formal solutions of irregular linear differential equations, I. *American Journal of Mathematics*, 56:411–437, 1934.
- [Cop36] F. Cope. Formal solutions of irregular linear differential equations, II. *American Journal of Mathematics*, 58:130–140, 1936.
- [CS06] P. J. Cassidy and M. F. Singer. Galois theory of parameterized differential equations and linear differential algebraic groups. In D. Bertrand, B. Enriquez, C. Mitschi, C. Sabbah, and R. Schaefer, editors, *Differential Equations and Quantum Groups*, volume 9 of *IRMA Lectures in Mathematics and Theoretical Physics*, pages 113–157. EMS Publishing House, 2006.
- [Del70] P. Deligne. *Equations Différentielles à Points Singuliers Réguliers*, volume 163 of *Lecture Notes in Mathematics*. Springer-Verlag, Heidelberg, 1970.
- [Del90] P. Deligne. Catégories tannakiennes. In P. Cartier et al., editor, *Grothendieck Festschrift, Vol. 2*, pages 111–195. Birkhäuser, 1990. Progress in Mathematics, Vol. 87.
- [DM82] P. Deligne and J. Milne. Tannakian categories. In P. Deligne et al., editor, *Hodge Cycles, Motives and Shimura Varieties*, pages 101–228, 1982. Lecture Notes in Mathematics, Vol. 900.
- [DM89] A. Duval and C. Mitschi. Groupe de Galois des équations hypergéométriques confluentes généralisées. *C. R. Acad. Sci. Paris*, 309(1):217–220, 1989.
- [dDR79] L. van den Dries and P. Ribenboim. Application de la théorie des modèles aux groupes de Galois de corps de fonctions. *C. R. Acad. Sci. Paris Sér. A-B*, 288(17):A789–A792, 1979.
- [DV02] L. Di Vizio. Arithmetic theory of  $q$ -difference equations: the  $q$ -analogue of Grothendieck-Katz’s conjecture on  $p$ -curvatures. *Invent. Math.*, 150(3):517–578, 2002.
- [DV04] L. Di Vizio. Introduction to  $p$ -adic  $q$ -difference equations (weak Frobenius structure and transfer theorems). In *Geometric aspects of Dwork theory. Vol. I, II*, pages 615–675. Walter de Gruyter GmbH & Co. KG, Berlin, 2004.
- [DVRSZ03] L. Di Vizio, J.-P. Ramis, J. Sauloy, and C. Zhang. Équations aux  $q$ -différences. *Gaz. Math.*, 96:20–49, 2003.

- [Dwo90] B. Dwork. Differential operators with nilpotent  $p$ -curvature. *Amer. J. Math.*, 112(5):749–786, 1990.
- [Dyc05] T. Dyckerhoff. Picard-Vessiot Extensions over Number Fields. Diplomarbeit, Fakultät für Mathematik und Informatik der Universität Heidelberg, 2005.
- [Dyc07] T. Dyckerhoff. The inverse problem of differential Galois theory over the field  $\mathbb{C}(z)$ . to appear Oberwohlfach Reports, 2007.
- [Eti95] P. I. Etingof. Galois groups and connection matrices of  $q$ -difference equations. *Electron. Res. Announc. Amer. Math. Soc.*, 1(1):1–9 (electronic), 1995.
- [FH91] W. Fulton and J. Harris. *Representation theory. A first course*, volume 129 of *Graduate Texts in Mathematics*. Springer, New York, 1991.
- [Fra63] C. H. Franke. Picard-Vessiot theory of linear homogeneous difference equations. *Transactions of the AMS*, 108:491–515, 1963.
- [GKZ94] I.M. Gelfand, M.M. Kapranov, and A. V. Zelevinsky. *Discriminants, Resultants and Multidimensional Determinants*. Birkhäuser, Boston, Basel, Stuttgart, 1994.
- [Gra00] J. J. Gray. *Linear Differential Equations and Group Theory from Riemann to Poincaré*. Birkhäuser, Boston, Basel, Stuttgart, second edition, 2000.
- [Har64] P. Hartman. *Ordinary differential equations*. John Wiley & Sons Inc., New York, 1964.
- [Har02] J. Hartmann. *On the inverse problem in differential Galois theory*. Thesis, University of Heidelberg, 2002. (available at [www.ub.uni-heidelberg.de/archiv/3085](http://www.ub.uni-heidelberg.de/archiv/3085)).
- [Har05] J. Hartmann. On the inverse problem in differential Galois theory. *J. Reine Angew. Math.*, 586:21–44, 2005.
- [Har06] Charlotte Hardouin. Hypertranscendence et groupes de galois aux différences. [arXiv:math.RT/0702846v1](https://arxiv.org/abs/math.RT/0702846v1), 2006.
- [Har07a] D. Harbater. Patching over fields (joint with Julia Hartmann). to appear Oberwohlfach Reports, 2007.
- [Har07b] C. Hardouin. Iterative  $q$ -difference Galois theory. to appear Oberwohlfach Reports, 2007.
- [Har07c] J. Hartmann. Differential Galois groups and patching (joint with David Harbater). to appear in Oberwohlfach Reports, 2007.
- [H02] M. van Hoeij. Decomposing a  $4^{th}$  order linear differential equation as a symmetric product. In *Differential Galois Theory*, volume 58 of *Banach Center Publications*, pages 89–96. Institute of Mathematics, Polish Academy of Sciences, Warszawa, 2002.
- [H07] M. van Hoeij. Solving third order linear differential equations in terms of second order equations. In C.W. Brown, editor, *Proceedings of ISSAC2007*, pages 355–360. ACM Press, 2007.
- [HW05] M. van Hoeij and J.-A. Weil. Solving second order linear differential equations with Klein’s theorem. In *ISSAC’05*, pages 340–347 (electronic). ACM, New York, 2005.
- [Hru02] E. Hrushovski. Computing the Galois group of a linear differential equation. In *Differential Galois Theory*, volume 58 of *Banach Center Publications*, pages 97–138. Institute of Mathematics, Polish Academy of Sciences, Warszawa, 2002.
- [HRUW99] M. van Hoeij, J.-F. Ragot, F. Ulmer, and J.-A. Weil. Liouvillian solutions of linear differential equations of order three and higher. *Journal*

- of *Symbolic Computation*, 28(4/5):589–610, 1999.
- [HS85] W.A. Harris and Y. Sibuya. The reciprocals of solutions of linear ordinary differential equations. *Adv. in Math.*, 58:119–132, 1985.
- [Hum75] J. Humphreys. *Linear Algebraic Groups*. Graduate Texts in Mathematics. Springer-Verlag, New York, 1975.
- [HUY07] R.-P. Holzapfel, A.M. Uludağ, and M. Yoshida, editors. *Arithmetic and geometry around hypergeometric functions*, volume 260 of *Progress in Mathematics*. Birkhäuser Verlag, Basel, 2007. Lecture notes of a CIMPA Summer School held at Galatasaray University, Istanbul, June 13–25, 2005.
- [HvdP95] P. A. Hendriks and M. van der Put. Galois action on solutions of a differential equation. *Journal of Symbolic Computation*, 19(6):559 – 576, 1995.
- [HW96] M. van Hoeij and J. A. Weil. An algorithm for computing invariants of differential Galois groups. *Journal of Pure and Applied Algebra*, 117/118:353–379, 1996.
- [Jac37] N. Jacobson. Pseudo-linear transformations. *Annals of Mathematics*, 38:484–507, 1937.
- [JL07a] L. Juan and A. Ledet. Equivariant vector fields on non-trivial  $SO_n$ -torsors and differential galois theory. Technical report, Texas Tech University, 2007. to appear in J. of Algebra.
- [JL07b] L. Juan and A. Ledet. On generic differential  $SO_n$ -extensions. Technical report, Texas Tech University, 2007. to appear in Proc. AMS.
- [JL07c] L. Juan and A. Ledet. On picard-vessiot extensions with group  $PGL_3$ . Technical report, Texas Tech University, 2007.
- [JRR95] J. Johnson, G. M. Reinhart, and L. A. Rubel. Some counterexamples to separation of variables. *J. Differential Equations*, 121(1):42–66, 1995.
- [Jua07] L. Juan. Generic picard-vessiot extensions for connected-by-finite groups. *J. of Algebra*, 312:194–2006, 2007.
- [Kam06] M. Kamensky. Definable groups of partial automorphisms. Preprint; <http://arxiv.org/abs/math.LO/0607718>, 2006.
- [Kap76] I. Kaplansky. *An Introduction to Differential Algebra*. Hermann, Paris, second edition, 1976.
- [Kat87a] N. Katz. A simple algorithm for cyclic vectors. *American Journal of Mathematics*, 109:65–70, 1987.
- [Kat87b] N. Katz. On the calculation of some differential Galois groups. *Inventiones Mathematicae*, 87:13–61, 1987.
- [Kat90] N. Katz. *Exponential Sums and Differential Equations*, volume 124 of *Annals of Mathematics Studies*. Princeton University Press, Princeton, 1990.
- [Kat96] N. Katz. *Rigid Local Systems*, volume 139 of *Annals of Mathematics Studies*. Princeton University Press, Princeton, 1996.
- [Kol48] Ellis R. Kolchin. Algebraic matrix groups and the Picard-Vessiot theory of homogeneous linear ordinary differential equations. *Annals of Mathematics*, 49:1–42, 1948.
- [Kol68] E. R. Kolchin. Algebraic groups and algebraic dependence. *American Journal of Mathematics*, 90:1151–1164, 1968.
- [Kol74] E. R. Kolchin. Constrained extensions of differential fields. *Advances in Math.*, 12:141–170, 1974.
- [Kol76] E. R. Kolchin. *Differential Algebra and Algebraic Groups*. Academic Press, New York, 1976.

- [Kol85] E. R. Kolchin. *Differential algebraic groups*, volume 114 of *Pure and Applied Mathematics*. Academic Press Inc., Orlando, FL, 1985.
- [Kov86] J.J. Kovacic. An algorithm for solving second order linear homogeneous differential equations. *Journal of Symbolic Computation*, 2:3–43, 1986.
- [Kov03] J. J. Kovacic. The differential Galois theory of strongly normal extensions. *Trans. Amer. Math. Soc.*, 355(11):4475–4522 (electronic), 2003.
- [Kov06] J. J. Kovacic. Geometric characterization of strongly normal extensions. *Trans. Amer. Math. Soc.*, 358(9):4135–4157 (electronic), 2006.
- [Lan66] S. Lang. *Introduction to Transcendental Numbers*. Addison Wesley, New York, 1966.
- [Lan06] P. Landesman. *Generalized Differential Galois Theory*. Ph.d. thesis, Graduate Center, The City University of New York, 2006.
- [Liȋ02] R. Liȋcanu. Counting Lamé differential operators. *Rend. Sem. Mat. Univ. Padova*, 107:191–208, 2002.
- [LR90] M. Loday-Richaud. Introduction à la multisommabilité. *Gazette des Mathématiciens, SMF*, 44:41–63, 1990.
- [LR94] M. Loday-Richaud. Stokes phenomenon, multisummability and differential Galois groups. *Annales de l'Institut Fourier*, 44(3):849–906, 1994.
- [LR95] M. Loday-Richaud. Solutions formelles des systèmes différentiels linéaires méromorphes et sommation. *Expositiones Mathematicae*, 13:116–162, 1995.
- [LSWZ06] Z. Li, M. F. Singer, M. Wu, and D. Zheng. A Recursive Method for Determining the One-Dimensional Submodules of Laurent-Ore Modules. In Jean-Guillaume Dumas, editor, *Proceedings of the 2006 International Symposium on Symbolic and Algebraic Computation (ISSAC 2006)*, pages 200–208. ACM Press, 2006.
- [LvdPU05] F. Loray, M. van der Put, and F. Ulmer. The Lamé family of connections on the projective line. Technical report, IRMAR, 2005.
- [Mag76] A. Magid. Finite generation of class groups of rings of invariants. *Proc. Amer. Math. Soc.*, 60:45–48, 1976.
- [Mag94] A. Magid. *Lectures on Differential Galois Theory*. University Lecture Series. American Mathematical Society, 1994. Second Edition.
- [Mai03] E. Maillet. Sur les séries divergentes et les équations différentielles. *Ann. Éc. Norm. Sup.*, pages 487–518, 1903.
- [Mai04] R. S. Maier. Algebraic solutions of the Lamé equation, revisited. *J. Differential Equations*, 198(1):16–34, 2004.
- [Mal01] B. Malgrange. Le groupoïde de Galois d'un feuilletage. In *Essays on geometry and related topics, Vol. 1, 2*, volume 38 of *Monogr. Enseign. Math.*, pages 465–501. Enseignement Math., Geneva, 2001.
- [Mal02] B. Malgrange. On nonlinear differential Galois theory. *Chinese Ann. Math. Ser. B*, 23(2):219–226, 2002. Dedicated to the memory of Jacques-Louis Lions.
- [Mar00] D. Marker. Model theory of differential fields. In *Model theory, algebra, and geometry*, volume 39 of *Math. Sci. Res. Inst. Publ.*, pages 53–63. Cambridge Univ. Press, Cambridge, 2000.
- [Med05] R. Bustamente Medina. *Théorie des modèles des corps différentiellement clos avec un automorphisme générique*. Ph.d. thesis, Université Paris 7, 2005.
- [Mit89] C. Mitschi. Matrices de Stokes et groupe de Galois des équations hypergéométriques confluentes généralisées. *Pacific Journal of Mathematics*,

- 138(1):25–56, 1989.
- [Mit96] C. Mitschi. Differential Galois groups of confluent generalized hypergeometric equations: An approach using Stokes multipliers. *Pacific Journal of Mathematics*, 176(2):365–405, 1996.
- [MP97] D. Marker and A. Pillay. Differential Galois theory. III. Some inverse problems. *Illinois J. Math.*, 41(3):453–461, 1997.
- [MR89] J. Martinet and J.-P. Ramis. Théorie de Galois différentielle et resommation. In E. Tournier, editor, *Computer Algebra and Differential Equations*, pages 115–214. Academic Press, 1989.
- [MR99] J.J. Morales-Ruiz. *Differential Galois Theory and Non-Integrability of Hamiltonian Systems*. Birkhäuser, Basel, 1999.
- [MS96] C. Mitschi and M. F. Singer. Connected linear groups as differential Galois groups. *Journal of Algebra*, 184:333–361, 1996.
- [MS02] C. Mitschi and M. F. Singer. Solvable-by-finite groups as differential Galois groups. *Ann. Fac. Sci. Toulouse*, XI(3):403 – 423, 2002.
- [MvdP03a] B. H. Matzat and M. van der Put. Constructive differential Galois theory. In *Galois groups and fundamental groups*, volume 41 of *Math. Sci. Res. Inst. Publ.*, pages 425–467. Cambridge Univ. Press, Cambridge, 2003.
- [MvdP03b] B. H. Matzat and M. van der Put. Iterative differential equations and the Abhyankar conjecture. *J. Reine Angew. Math.*, 557:1–52, 2003.
- [Ngu07] K.A. Nguyen. On d-solvability for linear differential equations. Preprint, University of Groningen, 2007.
- [NdP07] K.A. Nguyen and M. van der Put. Solving linear differential equations. Preprint, University of Groningen, 2007.
- [Obe03] T. Oberlies. *Embedding problems in differential Galois theory*. PhD thesis, University of Heidelberg, 2003. (available at [www.ub.uni-heidelberg.de/archiv/4550](http://www.ub.uni-heidelberg.de/archiv/4550)).
- [Per02] A. Person. *Solving Homogeneous Linear Differential Equations of Order 4 in Terms of Equations of Lower Order*. PhD thesis, North Carolina State University, 2002. <http://www.lib.ncsu.edu/theses/available/etd-08062002-104315/>.
- [Pil97] A. Pillay. Differential Galois theory. II. *Ann. Pure Appl. Logic*, 88(2-3):181–191, 1997. Joint AILA-KGS Model Theory Meeting (Florence, 1995).
- [Pil98] A. Pillay. Differential Galois theory. I. *Illinois J. Math.*, 42(4):678–699, 1998.
- [Pil02] A. Pillay. Finite-dimensional differential algebraic groups and the Picard-Vessiot theory. In *Differential Galois theory (Bedlewo, 2001)*, volume 58 of *Banach Center Publ.*, pages 189–199. Polish Acad. Sci., Warsaw, 2002.
- [Pil04] A. Pillay. Algebraic  $D$ -groups and differential Galois theory. *Pacific J. Math.*, 216(2):343–360, 2004.
- [Poo60] E.G.C. Poole. *Introduction to the Theory of Linear Differential Equations*. Dover Publications, New York, 1960.
- [dP95] M. van der Put. Differential equations in characteristic  $p$ . *Compositio Mathematica*, 97:227–251, 1995.
- [dP96] M. van der Put. Reduction modulo  $p$  of differential equations. *Indagationes Mathematicae*, 7:367–387, 1996.
- [dPR07] M. van der Put and M. Reversat. Galois Theory of  $q$ -difference equations. *Ann. Fac. Sci. de Toulouse*, XVI(2):1–54, 2007.



- [dPS97] M. van der Put and M. F. Singer. *Galois Theory of Difference Equations*, volume 1666 of *Lecture Notes in Mathematics*. Springer-Verlag, Heidelberg, 1997.
- [dPS03] M. van der Put and M. F. Singer. *Galois Theory of Linear Differential Equations*, volume 328 of *Grundlehren der mathematischen Wissenschaften*. Springer, Heidelberg, 2003.
- [Ram85a] J.-P. Ramis. Filtration Gevrey sur le groupe de Picard-Vessiot d'une équation différentielle irrégulière. *Informes de Matematica*, Preprint IMPA, Series A-045/85, 1985.
- [Ram85b] J.-P. Ramis. Phénomène de Stokes et filtration Gevrey sur le groupe de Picard-Vessiot. *Comptes Rendus de l'Académie des Sciences, Paris*, 301:165–167, 1985.
- [Ram85c] J.-P. Ramis. Phénomène de Stokes et resommation. *Comptes Rendus de l'Académie des Sciences, Paris*, 301:99–102, 1985.
- [Ram93] J.-P. Ramis. *Séries Divergentes et Théories Asymptotiques*. Number 121 in *Panoramas et Synthèses*. Société Mathématique de France, Paris, 1993.
- [Ram96] J.-P. Ramis. About the inverse problem in differential Galois theory: Solutions of the local inverse problem and of the differential Abhyankar conjecture. Technical report, Université Paul Sabatier, Toulouse, 1996.
- [Ram07] J.-P. Ramis. Local classification of linear meromorphic  $q$ -difference equations (joint with Jacques Sauloy). to appear *Oberwolfach Reports*, 2007.
- [Ros61] M. Rosenlicht. Toroidal algebraic groups. *Proc. Amer. Math. Soc.*, 12:984–988, 1961.
- [RS07] J.-P. Ramis and J. Sauloy. The  $q$ -analogue of the wild fundamental group. I. In *Algebraic, analytic and geometric aspects of complex differential equations and their deformations. Painlevé hierarchies*, RIMS Kôkyûroku Bessatsu, B2, pages 167–193. Res. Inst. Math. Sci. (RIMS), Kyoto, 2007.
- [RSZ06] J.-P. Ramis, J. Sauloy, and C. Zhang. Développement asymptotique et sommabilité des solutions des équations linéaires aux  $q$ -différences. *C. R. Math. Acad. Sci. Paris*, 342(7):515–518, 2006.
- [Sau03] J. Sauloy. Galois theory of Fuchsian  $q$ -difference equations. *Ann. Sci. École Norm. Sup. (4)*, 36(6):925–968 (2004), 2003.
- [Sei58] A. Seidenberg. Abstract differential algebra and the analytic case. *Proc. Amer. Math. Soc.*, 9:159–164, 1958.
- [Sei69] A. Seidenberg. Abstract differential algebra and the analytic case. II. *Proc. Amer. Math. Soc.*, 23:689–691, 1969.
- [Ser64] J.-P. Serre. *Cohomologie Galoisienne*. Number 5 in *Lecture Notes in Mathematics*. Springer-Verlag, New York, 1964.
- [Sin81] M. F. Singer. Liouvillian solutions of  $n^{th}$  order homogeneous linear differential equations. *American Journal of Mathematics*, 103:661–681, 1981.
- [Sin85] M. F. Singer. Solving homogeneous linear differential equations in terms of second order linear differential equations. *American Journal of Mathematics*, 107:663–696, 1985.
- [Sin86] M. F. Singer. Algebraic relations among solutions of linear differential equations. *Transactions of the American Mathematical Society*, 295:753–763, 1986.
- [Sin88] M. F. Singer. Algebraic relations among solutions of linear differential

- equations: Fano's theorem. *American Journal of Mathematics*, 110:115–144, 1988.
- [Sin93] M. F. Singer. Moduli of linear differential equations on the Riemann sphere with fixed Galois group. *Pacific Journal of Mathematics*, 106(2):343–395, 1993.
- [Sin07] M. F. Singer. Differential groups and differential relations. to appear in Oberwolfach Reports, 2007.
- [Spr98] T. A. Springer. *Linear Algebraic Groups, Second Edition*, volume 9 of *Progress in Mathematics*. Birkhäuser, Boston, 1998.
- [SST00] M. Sato, B. Sturmfels, and N. Takayama. *Gröbner Deformations of Hypergeometric Differential Equations*, volume 6 of *Algorithms and Computation in Mathematics*. Springer-Verlag, Berlin, 2000.
- [SU97] M. F. Singer and F. Ulmer. Linear differential equations and products of linear forms. *Journal of Pure and Applied Algebra*, 117:549–564, 1997.
- [Tre71] M. Tretkoff. Algebraic extensions of the field of rational functions. *Comm. Pure Appl. Math.*, 24:491–497, 1971.
- [TT79] C. Tretkoff and M. Tretkoff. Solution of the inverse problem in differential Galois theory in the classical case. *American Journal of Mathematics*, 101:1327–1332, 1979.
- [Ulm94] F. Ulmer. Irreducible linear differential equations of prime order. *Journal of Symbolic Computation*, 18(4):385–401, 1994.
- [Ume] H. Umemura. Galois Theory and Painlevé Equations. Preprint, 2004.
- [Ume90] H. Umemura. Birational automorphism groups and differential equations. *Nagoya Mathematics Journal*, 119:1–80, 1990.
- [Ume96a] H. Umemura. Differential Galois theory of infinite dimension. *Nagoya Mathematics Journal*, 144:59–135, 1996.
- [Ume96b] H. Umemura. Galois theory of algebraic and differential equations. *Nagoya Mathematics Journal*, 144:1–58, 1996.
- [Ume99] H. Umemura. Lie-Drach-Vessiot Theory. In *CR-Geometry and Overdetermined Systems*, volume 25 of *Advanced Studies in Pure Mathematics*, pages 364–385. North-Holland, 1999.
- [Ume04] H. Umemura. Monodromy preserving deformation and differential Galois group. I. Analyse complexe, systèmes dynamiques, sommabilité des séries divergentes et théories galoisiennes. I, 2004.
- [UW96] F. Ulmer and J.-A. Weil. A note on Kovacic's algorithm. *Journal of Symbolic Computation*, 22(2):179 – 200, 1996.
- [Var91] V. S. Varadarajan. Meromorphic differential equations. *Expositiones Mathematicae*, 9(2):97–188, 1991.
- [Var96] V. S. Varadarajan. Linear meromorphic differential equations: A modern point of view. *Bulletin (New Series) of the American Mathematical Society*, 33(1):1 – 42, 1996.
- [Wu05] M. Wu. *On Solutions of Linear Functional Systems and Factorization of Modules over Laurent-Ore Algebras*. Ph.d. thesis, Chinese Academy of Sciences and l'Université de Nice-Sophia Antipolis, 2005.