

Progress in Mathematics
203

David E. Blair

Riemannian Geometry of Contact and Symplectic Manifolds

Second Edition



 Birkhäuser

Progress in Mathematics
Volume 203

Series Editors

Hyman Bass

Joseph Oesterlé

Alan Weinstein

David E. Blair

Riemannian Geometry of Contact and Symplectic Manifolds

Second Edition

David E. Blair
Department of Mathematics
Michigan State University
East Lansing, MI 48824-1027
USA
blair@math.msu.edu

ISBN 978-0-8176-4958-6 e-ISBN 978-0-8176-4959-3
DOI 10.1007/978-0-8176-4959-3

Library of Congress Control Number: 2010932450

Mathematics Subject Classification (2010): 53B35, 53C15, 53C25, 53C26, 53C40, 53C42, 53C55, 53C56, 53D05, 53D10, 53D12, 53D15, 53D22, 53D25, 53D35, 58E11

© Springer Science+Business Media, LLC 2002, 2010

All rights reserved. This work may not be translated or copied in whole or in part without the written permission of the publisher (Springer Science+Business Media, LLC, 233 Spring Street, New York, NY 10013, USA), except for brief excerpts in connection with reviews or scholarly analysis. Use in connection with any form of information storage and retrieval, electronic adaptation, computer software, or by similar or dissimilar methodology now known or hereafter developed is forbidden.

The use in this publication of trade names, trademarks, service marks, and similar terms, even if they are not identified as such, is not to be taken as an expression of opinion as to whether or not they are subject to proprietary rights.

Printed on acid-free paper

birkhauser-science.com

To Rebecca
*in appreciation of all her
love and support*

Preface to Second Edition

For this second edition almost every chapter has been revised and updated in some way. The principal changes from the first edition are the following. A second chapter on the geometry of complex contact manifolds, Chapter 13, has been added. Chapter 7 on the curvature of contact metric manifolds has been reorganized and updated extensively. A section on the projectivized tangent bundle has been added to Chapter 9. Additional examples and commentary on further results have been added throughout.

There has been considerable work in recent years on a number of areas related to the subject, and it has been impossible to treat all of this work. The author believes, however, that the text offers a good introduction to and necessary background for the study of these topics.

The author expresses his appreciation to C. Abbas, T. Draghici, B. Foreman and B. Korkmaz for reading parts of the manuscript and offering valuable suggestions. The author also expresses his appreciation to Birkhäuser for suggesting that he write a second edition and especially to Ann Kostant, Jessica Belanger and Tom Grasso for their kind assistance in producing this book.

March, 2010

David E. Blair

Preface to the First Edition

The author's lectures "Contact Manifolds in Riemannian Geometry", volume 509 (1976), in the Springer-Verlag series *Lecture Notes in Mathematics* have been out of print for some time and it seems appropriate that an expanded version of this material should become available. The present text deals with the Riemannian geometry of both symplectic and contact manifolds, although the book is more contact than symplectic. This work is based on the recent research of the author, his students, colleagues, and other scholars, the author's graduate courses at Michigan State University and the earlier lecture notes.

Chapter 1 presents the general theory of symplectic manifolds. Principal circle bundles are then discussed in Chapter 2 as a prelude to the Boothby–Wang fibration of a compact regular contact manifold in Chapter 3, which deals with the general theory of contact manifolds. Chapter 4 focuses on Riemannian metrics associated to symplectic and contact structures. Chapter 5 is devoted to integral submanifolds of the contact subbundle. In Chapter 6 we discuss the normality of almost contact structures, Sasakian manifolds, K-contact manifolds, the relation of contact metric structures and CR-structures, and cosymplectic structures. Chapter 7 deals with the important study of the curvature of a contact metric manifold. In Chapter 8 we give a selection of results on submanifolds of

Kähler and Sasakian manifolds, including an illustration of the technique of A. Ros in a theorem of F. Urbano on compact minimal Lagrangian submanifolds in $\mathbb{C}P^n$. Chapter 9 discusses the symplectic structure of tangent bundles, contact structure of tangent sphere bundles, general vector bundles and normal bundles of Lagrangian and integral submanifolds giving rise to new examples of symplectic and contact manifolds. In Chapter 10 we study a number of curvature functionals on spaces of associated metrics and their critical point conditions; we show also that in the symplectic case, the “total scalar curvature” is a symplectic invariant and in the contact case is a natural functional whose critical points are the metrics for which the characteristic vector field generates isometries. In the presence of a certain amount of negative curvature, special directions appear in the contact subbundle; we discuss these and their relations to Anosov and conformally Anosov flows in Chapter 11. Chapter 12 deals with the subject of complex contact manifolds. We conclude with a brief treatment of 3-Sasakian manifolds in Chapter 13.

The text attempts to strike a balance between giving detailed proofs of basic properties, which will be instructive to the reader, and stating many results whose proofs would take us too far afield. It has been impossible, however, to be encyclopedic and include everything, so that unfortunately some important topics have been omitted or covered only briefly. An extensive bibliography is given.

It is the author’s hope that the reader will find this both a good introduction to the Riemannian geometry of contact and symplectic manifolds and a useful reference to recent research in the area.

The author expresses his appreciation to C. Baikoussis, B.-Y. Chen, D. Chinea, T. Draghici, B. Foreman, Th. Koufogiorgos, Y.-H. P. Pang and D. Perrone for reading parts of the manuscript and offering valuable suggestions. The author also expresses his appreciation to Ann Kostant of Birkhäuser and to Elizabeth Loew of T_EXniques for their kind assistance in the production of this book.

October, 2001

David E. Blair

Contents

Preface to Second Edition	vii
Preface to the First Edition	ix
1 Symplectic Manifolds	1
1.1 Definitions and examples	1
1.2 Lagrangian submanifolds	6
1.3 The Darboux–Weinstein theorems	9
1.4 Symplectomorphisms	11
2 Principal S^1-bundles	15
2.1 The set of principal S^1 -bundles as a group	15
2.2 Connections on a principal bundle	19
3 Contact Manifolds	23
3.1 Definitions	23
3.2 Examples	27
3.2.1 \mathbb{R}^{2n+1}	27
3.2.2 $\mathbb{R}^{n+1} \times P\mathbb{R}^n$	28
3.2.3 $M^{2n+1} \subset \mathbb{R}^{2n+2}$ with $T_m M^{2n+1} \cap \{0\} = \emptyset$	28
3.2.4 Unit and projectivized tangent and cotangent bundles	29
3.2.5 $T^*M \times \mathbb{R}$	30

3.2.6	Tori	30
3.2.7	Overtwisted contact structures	32
3.2.8	$S^2 \times S^1$	34
3.2.9	Contact circles	35
3.3	The Boothby–Wang fibration	36
3.4	The Weinstein conjecture	38
4	Associated Metrics	41
4.1	Almost complex and almost contact structures	41
4.2	Polarization and associated metrics	45
4.3	Polarization of metrics as a projection	49
4.3.1	Some linear algebra	50
4.3.2	Results on the set \mathcal{A}	53
4.4	Action of symplectic and contact transformations	57
4.5	Examples of almost contact metric manifolds	60
4.5.1	\mathbb{R}^{2n+1}	60
4.5.2	$M^{2n+1} \subset \tilde{M}^{2n+2}$ almost complex	61
4.5.3	$S^5 \subset S^6$	63
4.5.4	The Boothby–Wang fibration	64
4.5.5	$M^{2n} \times \mathbb{R}$	66
4.5.6	Parallelizable manifolds	67
5	Integral Submanifolds and Contact Transformations	69
5.1	Integral submanifolds	69
5.2	Contact transformations	71
5.3	Examples of integral submanifolds	74
5.3.1	$S^n \subset S^{2n+1}$	74
5.3.2	$T^2 \subset S^5$	74
5.3.3	Legendre curves and Whitney spheres	75
5.3.4	Lift of a Lagrangian foliation, Legendre foliations	77
6	Sasakian and Cosymplectic Manifolds	79
6.1	Normal almost contact structures	79
6.2	The tensor field h	83
6.3	Definition of a Sasakian manifold	86
6.4	CR-manifolds	89
6.5	Cosymplectic manifolds and remarks on the Sasakian definition	95
6.6	Products of almost contact manifolds	97

6.7	Examples	100
6.7.1	\mathbb{R}^{2n+1}	100
6.7.2	Principal circle bundles	100
6.7.3	A nonnormal almost contact structure on S^5 . . .	102
6.7.4	$M^{2n+1} \subset \tilde{M}^{2n+2}$	104
6.7.5	Brieskorn manifolds	104
6.8	Some early topology	106
7	Curvature of Contact Metric Manifolds	111
7.1	Basic curvature properties	111
7.2	Curvature of contact metric manifolds	116
7.3	The (κ, μ) -manifolds	123
7.4	Sasakian Einstein manifolds	130
7.5	Locally symmetric contact metric manifolds	132
7.6	Conformally flat contact metric manifolds	133
7.7	ϕ -sectional curvature	137
7.8	Examples of Sasakian space forms	141
7.8.1	S^{2n+1}	142
7.8.2	\mathbb{R}^{2n+1}	142
7.8.3	$B^n \times \mathbb{R}$	142
7.9	Locally ϕ -symmetric spaces	143
8	Submanifolds of Kähler and Sasakian Manifolds	151
8.1	Invariant submanifolds	151
8.2	Lagrangian and integral submanifolds	155
9	Tangent Bundles and Tangent Sphere Bundles	169
9.1	Tangent bundles	169
9.2	Tangent sphere bundles	175
9.3	Geometry of vector bundles	183
9.4	Normal bundles	186
9.5	The geodesic flow on the projectivized tangent bundle . .	191
10	Curvature Functionals on Spaces of Associated Metrics	195
10.1	Introduction to critical metric problems	195
10.2	The $*$ -scalar curvature	201
10.3	The integral of $Ric(\xi)$	206
10.3.1	H-contact manifolds	211
10.4	The Webster scalar curvature	212

10.5	A gauge invariant	215
10.6	The Abbena metric as a critical point	217
11	Negative ξ-sectional Curvature	219
11.1	Special directions in the contact subbundle	219
11.2	Anosov flows	221
11.3	Conformally Anosov flows	227
12	Complex Contact Manifolds	233
12.1	Complex contact manifolds and associated metrics	233
12.2	Examples of complex contact manifolds	238
12.2.1	Complex Heisenberg group	238
12.2.2	Odd-dimensional complex projective space	240
12.2.3	Twistor spaces	242
12.2.4	The Complex Boothby–Wang fibration	244
12.2.5	3-dimensional homogeneous examples	246
12.2.6	Complex contact Lie groups	247
12.2.7	$\mathbb{C}^{n+1} \times \mathbb{C}P^n(16)$	248
12.2.8	$\cos z^3 dz^1 + \sin z^3 dz^2$	250
12.3	Normality of complex contact manifolds	250
12.4	GH -sectional curvature	252
12.5	The set of associated metrics and integral functionals	255
12.6	Holomorphic Legendre curves	257
12.7	The Calabi (Veronese) embeddings as integral submanifolds of $\mathbb{C}P^{2n+1}$	260
13	Additional Topics in Complex Geometry	265
13.1	Partial and holomorphic hyperbolicity	265
13.2	Projectivized holomorphic bundles	268
13.3	The complex geodesic flow	271
13.4	Complex almost contact metric structure on $P\tau$	278
13.4.1	A complex contact structure with nonintegrable vertical subbundle	280
13.5	Special directions on complex contact manifolds and the Lie group $SL(2, \mathbb{C})$	283

14 3-Sasakian Manifolds	291
14.1 3-Sasakian manifolds	291
14.2 Integral submanifolds	299
Bibliography	303
Subject Index	335
Author Index	339

1

Symplectic Manifolds

To set the stage for our development we begin this book with a treatment of the basic features of symplectic geometry. In this chapter we discuss symplectic manifolds and make brief mention of “associated metrics”, a topic that will be thoroughly discussed in Chapter 4. Here we treat in detail Lagrangian submanifolds and theorems of Darboux and Weinstein on the local structure of a symplectic manifold. We end this chapter with a brief discussion of symplectomorphisms.

1.1 Definitions and examples

By a *symplectic manifold* we mean an even-dimensional differentiable (C^∞) manifold M^{2n} together with a global 2-form Ω which is closed and of maximal rank, i.e., $d\Omega = 0$, $\Omega^n \neq 0$. By a *symplectomorphism* $f : (M_1, \Omega_1) \longrightarrow (M_2, \Omega_2)$ we mean a diffeomorphism $f : M_1 \longrightarrow M_2$ such that $f^*\Omega_2 = \Omega_1$.

Before continuing with symplectic manifolds we present some basic linear algebra. On a vector space V^{2n} , if $\Omega \in \bigwedge^2 V$ with $\text{rk } \Omega = 2n$, then there exist $\theta^1, \dots, \theta^{2n} \in V^*$, linearly independent and such that

$$\Omega = \theta^1 \wedge \theta^2 + \theta^3 \wedge \theta^4 + \dots + \theta^{2n-1} \wedge \theta^{2n}.$$

To see this, for a basis $\{\omega^i\}$ of V^* write

$$\begin{aligned}\Omega &= \sum_{i < j} a_{ij} \omega^i \wedge \omega^j = \omega^1 \wedge \sum_{1 < j} a_{1j} \omega^j + \omega^2 \wedge \sum_{2 < j} a_{2j} \omega^j \\ &\quad + \text{terms in } \omega^3, \dots, \omega^{2n} \\ &= \omega^1 \wedge \beta^1 + \omega^2 \wedge \beta^2 + \text{terms in } \omega^3, \dots, \omega^{2n},\end{aligned}$$

where β^2 involves only $\omega^3, \dots, \omega^{2n}$, $\beta^1 = a\omega^2 + \beta^3$, and β^3 involves only $\omega^3, \dots, \omega^{2n}$. Therefore

$$\begin{aligned}\Omega &= \omega^1 \wedge \beta^1 + \frac{1}{a} \beta^1 \wedge \beta^2 - \frac{1}{a} \beta^3 \wedge \beta^2 + \text{terms in } \omega^3, \dots, \omega^{2n} \\ &= \left(\omega^1 - \frac{1}{a} \beta^2 \right) \wedge \beta^1 + \text{terms in } \omega^3, \dots, \omega^{2n}\end{aligned}$$

which is of the form $\theta^1 \wedge \theta^2 + \Omega_1$, where Ω_1 involves only $\omega^3, \dots, \omega^{2n}$. Now repeat the process for Ω_1 .

We shall often choose the labeling such that

$$\Omega = \theta^1 \wedge \theta^{n+1} + \dots + \theta^n \wedge \theta^{2n}.$$

As a corollary we see that there exists a basis $\{e_i, e_{n+i}\}$ of V^{2n} such that $\Omega(e_i, e_{n+j}) = \delta_{ij}$, $i, j = 1, \dots, n$. A change of basis that leaves invariant the normal form $\Omega = \sum_{i=1}^n \theta^i \wedge \theta^{n+i}$ is given by a symplectic matrix, i.e., $\begin{pmatrix} A & B \\ C & D \end{pmatrix}^{-1} = \begin{pmatrix} D & -C \\ -B & A \end{pmatrix}^T$ if and only if

$$\begin{pmatrix} A & B \\ C & D \end{pmatrix}^T \begin{pmatrix} 0 & I \\ -I & 0 \end{pmatrix} \begin{pmatrix} A & B \\ C & D \end{pmatrix} = \begin{pmatrix} 0 & I \\ -I & 0 \end{pmatrix}.$$

In particular, the structural group of the tangent bundle of a symplectic manifold is reducible to $Sp(2n, \mathbb{R})$. Further, using the fact that M^{2n} may be given a Riemannian metric and is orientable, the structural group is reducible to $SO(2n)$ and hence in turn to $U(n)$. Thus in particular, M^{2n} carries an almost complex structure; this will be discussed in greater detail below and in Chapter 4. The name *symplectic* is due to H. Weyl [1939, p. 165] changing the Latin *com/plex* to the Greek *sym/plectic*.

Two canonical examples of symplectic manifolds are the following:

1. \mathbb{R}^{2n} with coordinates $(x^1, \dots, x^n, y^1, \dots, y^n)$ admits the symplectic form $\Omega = \sum dx^i \wedge dy^i$. The classical theorem of Darboux states

that on any symplectic manifold there exist local coordinates with respect to which the symplectic form can be written in this way. We will give a modern proof of this result in Section 1.3.

2. Let M be a differentiable manifold. Then its cotangent bundle T^*M has a natural symplectic structure. For $z \in T^*M$ and V in the tangent space of T^*M at z , $T_z T^*M$, define a 1-form β , often called the *Liouville form*, by $\beta(V)_z = z(\pi_* V)$, where $\pi : T^*M \rightarrow M$ is the projection map. If x^1, \dots, x^n are local coordinates on M , then $q^i = x^i \circ \pi$ together with fiber coordinates p^1, \dots, p^n give local coordinates on T^*M . In these coordinates β has the local expression $\sum_{i=1}^n p^i dq^i$. The natural symplectic structure on T^*M is given by $\Omega = -d\beta$.

The reader may recognize the second example from classical mechanics; indeed, the cotangent bundle of the configuration space may be thought of as the phase space of a dynamical system, and we may obtain Hamilton's equations of motion as follows. Let H be a real-valued function on a symplectic manifold (M, Ω) and define a vector field X_H by $\Omega(X_H, Y) = YH$; X_H is called the *Hamiltonian vector field* generated by H . Two basic properties of X_H are $\mathcal{L}_{X_H} \Omega = 0$, \mathcal{L} being Lie differentiation, and $X_H H = 0$. In fact, the classical Poisson bracket is $\{f_1, f_2\} = \Omega(X_{f_2}, X_{f_1}) = X_{f_1} f_2$. In local coordinates $(q^1, \dots, q^n, p^1, \dots, p^n)$ given by the Darboux theorem, $\Omega = \sum dq^i \wedge dp^i$ and $X_H = \sum \left(-\frac{\partial H}{\partial q^i} \frac{\partial}{\partial p^i} + \frac{\partial H}{\partial p^i} \frac{\partial}{\partial q^i} \right)$. Thus the differential equations for the integral curves of X_H are

$$\dot{p}^i = -\frac{\partial H}{\partial q^i}, \quad \dot{q}^i = \frac{\partial H}{\partial p^i},$$

Hamilton's equations of motion.

Before giving further examples, we mention the relationship with Riemannian geometry that will become central for our study, viz., "associated metrics". Given a symplectic manifold (M, Ω) there exist a Riemannian metric g and an almost complex structure J such that $\Omega(X, Y) = g(X, JY)$. In fact, we shall see in Chapter 4 that there are many such metrics, g and J being created simultaneously by polarization.

On the other hand, given an almost complex structure J , i.e., a tensor field J of type $(1, 1)$ such that $J^2 = -I$, on an almost complex manifold, a Riemannian metric is said to be *Hermitian* if $g(JX, JY) = g(X, Y)$ and

the pair (J, g) is called an *almost Hermitian structure*. Now defining a 2-form Ω by $\Omega(X, Y) = g(X, JY)$, Ω is called the *fundamental 2-form* of the almost Hermitian structure. If $d\Omega = 0$, the almost Hermitian structure is said to be *almost Kähler*, whereas the structure is *Kähler* if J , or equivalently Ω , is parallel with respect to the Levi-Civita connection of g . Thus associated metrics can be thought of as almost Kähler structures whose fundamental 2-form is the given symplectic form.

Since $d\Omega = 0$, $[\Omega] \in H^2(M, \mathbb{R})$, $[\Omega]$ denoting the de Rham cohomology class determined by Ω . Using an associated metric, $\delta\Omega = 0$ and hence Ω is harmonic. To see this, use the fact that an almost Kähler manifold is *quasi-Kähler*, i.e., $(\nabla_k J_{ip})J_j^p = (\nabla_p J_{ij})J_k^p$, sum on the indices i and k , and use $J^2 = -I$. Also $[\Omega]^n = [\Omega^n] \in H^{2n}(M, \mathbb{R})$. In particular, for M compact the following are two necessary conditions for the existence of a symplectic structure:

- (i) M carries an almost complex structure.
- (ii) There exists an element $w \in H^2(M, \mathbb{R})$ such that $w^n \neq 0$.

Thus, for example, from (i) the 4-dimensional sphere S^4 is not symplectic, and from (ii), S^6 is not symplectic.

If M is an open manifold, Gromov in his thesis proved that (i) implies the existence of a 1-form ω such that $d\omega$ is symplectic (see A. Haefliger [1971, p. 133]). Also, Kähler manifolds are symplectic, so there are plenty of compact ones, e.g., complex projective space, $S^2 \times S^2$, algebraic varieties. Note that the even-dimensional Betti numbers of a compact almost Kähler manifold are nonzero. It is also well known that the odd-dimensional Betti numbers of a compact Kähler manifold are even, but this is not true in the almost Kähler case. We now give two descriptions of an example of Thurston [1976] of a compact symplectic manifold with no Kähler structure; this manifold is known as the *Thurston manifold* or as the *Kodaira–Thurston manifold* (Kodaira [1964]).

Briefly, first take the product of a torus T^2 , as a unit square with opposite sides identified, and an interval and glue the ends together by the diffeomorphism of T^2 given by the matrix $\begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}$. This gives a compact 3-manifold whose first Betti number is 2. Now taking the product with S^1 , we have a 4-manifold M with first Betti number 3, and hence M cannot have a Kähler structure. Let θ_1, θ_2 be coordinates on T^2 ; then $d\theta_1 \wedge d\theta_2$ exists after the twisting on the 3-manifold. Thus if ϕ_1 is the

coordinate on the interval and ϕ_2 the coordinate on the final circle, $d\theta_1 \wedge d\theta_2 + d\phi_1 \wedge d\phi_2$ is a symplectic form.

A second version of this example was given by E. Abbena [1984], who also gave a natural associated metric for this symplectic structure, computed its curvature, and showed that the first Betti number is 3 using harmonic forms.

Let G be the closed connected subgroup of $GL(4, \mathbb{C})$ defined by

$$\left\{ \begin{pmatrix} 1 & a_{12} & a_{13} & 0 \\ 0 & 1 & a_{23} & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & e^{2\pi i a} \end{pmatrix} \mid a_{12}, a_{13}, a_{23}, a \in \mathbb{R} \right\},$$

i.e., G is the product of the Heisenberg group and S^1 . Let Γ be the discrete subgroup of G with integer entries and $M = G/\Gamma$. Denote by x, y, z, t coordinates on G , say for $A \in G$, $x(A) = a_{12}$, $y(A) = a_{23}$, $z(A) = a_{13}$, $t(A) = a$. If L_B is left translation by $B \in G$, then $L_B^* dx = dx$, $L_B^* dy = dy$, $L_B^*(dz - xdy) = dz - xdy$, $L_B^* dt = dt$. In particular, these forms are invariant under the action of Γ . Let $\pi : G \rightarrow M$ denote projection. Then there exist 1-forms $\alpha_1, \alpha_2, \alpha_3, \alpha_4$ on M such that $dx = \pi^* \alpha_1$, $dy = \pi^* \alpha_2$, $dz - xdy = \pi^* \alpha_3$, $dt = \pi^* \alpha_4$. Setting $\Omega = \alpha_4 \wedge \alpha_1 + \alpha_2 \wedge \alpha_3$, we see that $\Omega \wedge \Omega \neq 0$ and $d\Omega = 0$ on M giving M a symplectic structure.

The vector fields

$$\mathbf{e}_1 = \frac{\partial}{\partial x}, \quad \mathbf{e}_2 = \frac{\partial}{\partial y} + x \frac{\partial}{\partial z}, \quad \mathbf{e}_3 = \frac{\partial}{\partial z}, \quad \mathbf{e}_4 = \frac{\partial}{\partial t}$$

are dual to $dx, dy, dz - xdy, dt$ and are left-invariant. Moreover, $\{\mathbf{e}_i\}$ is orthonormal with respect to the left-invariant metric on G given by

$$ds^2 = dx^2 + dy^2 + (dz - xdy)^2 + dt^2.$$

On M the corresponding metric is $g = \sum \alpha_i \otimes \alpha_i$ and is called the *Abbena metric*.

Moreover, M carries an almost complex structure defined by

$$J\mathbf{e}_1 = \mathbf{e}_4, \quad J\mathbf{e}_2 = -\mathbf{e}_3, \quad J\mathbf{e}_3 = \mathbf{e}_2, \quad J\mathbf{e}_4 = -\mathbf{e}_1.$$

Then noting that $\Omega(X, Y) = g(X, JY)$, we see that g is an associated metric.

At the time of Thurston's example there was no known example of a compact symplectic manifold with no Kähler structure. Since the appearance of this example there have been many others, e.g., Watson [1983], McDuff [1984], Cordero, Fernández and de Leon [1985], Cordero, Fernández and Gray [1986], Benson and Gordon [1988], [1990], Fernández and Gray [1990], Yamato [1990], McCarthy and Wolfson [1994], Gompf [1994], [1995], Fernández, de Leon and Saralegi [1996], Jelonek [1996], Holubowicz and Mozgawa [1998]. These examples are of several types: nil-manifolds, solvmanifolds, simply connected examples obtained by symplectic blowup, symplectic sums, and total spaces of fiber bundles. A survey of these topics can be found in the book of A. Tralle and J. Oprea [1997]. We also mention that Guan [1994] gave examples of complex symplectic manifolds (also known as holomorphic symplectic manifolds) that are not Kähler including the complexification of the Kodaira–Thurston manifold. A *complex symplectic manifold* is a complex manifold of complex dimension $2n$ together with a closed holomorphic 2-form Ω such that $\Omega^n \neq 0$. Further examples of complex symplectic manifolds were given by Yamada [2005].

1.2 Lagrangian submanifolds

Let $\iota : L \longrightarrow M^{2n}$ be an immersion into a symplectic manifold (M^{2n}, Ω) . We say that L is a *Lagrangian submanifold* if the dimension of L is n and $\iota^*\Omega = 0$. Two simple examples are the following:

1. The fibers of the cotangent bundle T^*M as discussed in the previous section are Lagrangian submanifolds with respect to the symplectic structure $d\beta$. Also suppose that ϕ is a section of T^*M . Then $\phi^*\beta$ is equal to ϕ as a 1-form on M ; in particular,

$$(\phi^*\beta)(X)_m = \beta(\phi_*X)_{\phi(m)} = \phi(m)(\pi_*\phi_*X) = \phi(X)_m.$$

Therefore $\phi^*\Omega = \phi^*(-d\beta) = -d\phi$; thus a section $\phi : M \longrightarrow T^*M$ is a Lagrangian submanifold if and only if ϕ is closed. When ϕ is exact, say dS , S is said to be a *generating function* for the submanifold.

2. Let (M_1, Ω_1) and (M_2, Ω_2) be symplectic manifolds and $f : M_1 \longrightarrow M_2$ a diffeomorphism. Then $(M_1, \Omega_1) \times (M_2, -\Omega_2)$ is symplectic, say $M = M_1 \times M_2$ with projections π_1 and π_2 and $\Omega = \pi_1^*\Omega_1 - \pi_2^*\Omega_2$.

Let Γ_f denote the graph of f . Then f is a symplectomorphism if and only if Γ_f is a Lagrangian submanifold of (M, Ω) .

There are several difficulties in studying Lagrangian submanifolds. First of all, since $\iota^*\Omega = 0$, there is no induced structure, so in a sense the geometry is transverse to the submanifold. From the standpoint of submanifold theory the codimension is high, so that the theory is more complicated. Another difficulty is that Lagrangian submanifolds are very abundant. For example, given any vector X at a point $m \in M$, there exists a Lagrangian submanifold through the point tangent to X . We shall see this as a corollary to the Darboux theorem in the next section. Also, Lagrangian submanifolds tend to get in the way of each other; loosely speaking, two Lagrangian submanifolds that are C^1 -close tend to intersect more than one would expect of two arbitrary C^1 -close submanifolds. Going into this point in some detail, let M be a compact manifold and, identifying M with the zero section of T^*M , view M as a Lagrangian submanifold of T^*M . Now let L be a Lagrangian submanifold of T^*M near M . Regarding L as the image of a closed 1-form ϕ , the question of when L and M intersect reduces to the question of when ϕ has a zero. So to perturb M to a disjoint Lagrangian submanifold L , M must admit a closed 1-form without zeros. An obstruction to this was given by Tischler [1970] in the following theorem.

Theorem 1.1 *If a compact manifold admits a closed 1-form without zeros, then the manifold fibers over the circle and conversely.*

In contrast, the problem of perturbing M to an arbitrary disjoint submanifold is equivalent to finding a nonvanishing 1-form, which is equivalent to finding a nonvanishing vector field, and the obstruction to this is the Euler characteristic.

If M is simply connected, the situation is “worse”. For now ϕ is exact and hence given by a function on M , but since M is compact, such a function must have at least two critical points. So a perturbation of S^2 to an arbitrary submanifold in T^*S^2 may intersect in only one point, but a perturbation to a Lagrangian submanifold must have at least two intersection points.

For L_1 and L_2 , C^1 -close Lagrangian submanifolds of a symplectic manifold (M, Ω) , the same situation holds by virtue of the following theorem of Weinstein [1971] which we will prove in the next section.

Theorem (Weinstein) *If L is a Lagrangian submanifold of a symplectic manifold (M, Ω) , then there exists a neighborhood of L in M that is symplectomorphic to a neighborhood of the zero section in T^*L .*

More general than the notion of a Lagrangian submanifold are the notions of isotropic and coisotropic submanifolds. A submanifold $\iota : N \rightarrow M^{2n}$ is *isotropic* if $\iota^*\Omega = 0$, so in particular the dimension of N is $\leq n$. In the subject of almost Hermitian manifolds (M^{2n}, J, g) these submanifolds are called *totally real* submanifolds; see Yau [1974], Chen and Ogiue [1974a]. The key point is that since $\Omega(X, Y) = g(X, JY)$, J maps the tangent space into the normal space. We remark that one sometimes sees another notion of totally real submanifold in the literature, namely a submanifold for which no tangent space contains a nonzero complex subspace; however, we will use only the stronger notion in this text.

The isotropic or totally real condition at a point $m \in N$ can be written as $\iota_*T_mN \subseteq \{V \mid \Omega(V, \iota_*T_mN) = 0\}$. A submanifold $\iota : N \rightarrow M^{2n}$ is *coisotropic* if $\iota_*T_mN \supseteq \{V \mid \Omega(V, \iota_*T_mN) = 0\}$; in terms of (J, g) , J maps the normal space into the tangent space and hence the dimension of N is $\geq n$.

In particular, for a Lagrangian submanifold N^n in \mathbb{C}^n , J maps the tangent spaces onto the normal spaces; therefore $T\mathbb{C}^n|_N = TN^n \oplus iTN^n = TN \otimes \mathbb{C}$ and hence the complexified tangent bundle is trivial. Gromov [1971] (see also Weinstein [1977]) proved that if N^n is compact, then N^n admits a Lagrangian immersion into \mathbb{C}^n if and only if the complexified tangent bundle is trivial.

The question of embeddings is a different matter. It is known that the sphere S^n cannot be embedded in \mathbb{C}^n as a Lagrangian submanifold. This is a consequence of a more general result of Gromov [1985] that a compact embedded Lagrangian submanifold in \mathbb{C}^n cannot be simply connected (see also Sikorav [1986]). For an immersed sphere as a Lagrangian submanifold with only one double point, see Example 5.3.3, Weinstein [1977, p. 26], or Morvan [1983].

Our discussion also has the following application to the problem of fixed points of symplectomorphisms; see Weinstein [1977, p. 29]. Let (M, Ω) be a compact simply connected symplectic manifold. Then a symplectomorphism f sufficiently C^1 close to the identity has at least two fixed points. To see this, let Δ be the diagonal of $(M, \Omega) \times (M, -\Omega)$ and Γ_f the graph of f . Then Δ and Γ_f intersect at least twice, and hence

f has at least two fixed points. For example, let M be complex projective space $\mathbb{C}P^n$ and f an automorphism of the Kähler structure that is C^1 -close to the identity. Any function on $\mathbb{C}P^n$ has at least $n + 1$ critical points, so that f must have at least $n + 1$ fixed points.

1.3 The Darboux–Weinstein theorems

We have mentioned the theorems of Darboux and Weinstein already; in this section we present a modern proof of both theorems using a technique of Moser [1965]. The use of this idea to prove the classical Darboux theorem is due to Weinstein [1971, 1977] and independently to J. Martinet [1970]. In addition to the papers mentioned, a general reference is the book by P. Libermann and C.-M. Marle [1987, Chapter III, Section 15] which we follow here; for the Darboux theorem see also N. Woodhouse [1980, pp. 7–9]. We begin with the following theorem of Weinstein. As a matter of notation, for a submanifold $\iota : N \rightarrow M$ and a differential form Φ on M , $\Phi|_N$ denotes the form acting on $T_N M$, the restriction of TM to N , and not the pullback, $\iota^* \Phi$, of Φ to N (see, e.g., Libermann and Marle [1987, p. 360]).

Theorem 1.2 *Let Ω_0 and Ω_1 be symplectic forms on a symplectic manifold M , and N a submanifold (possibly a point) on which $\Omega_0|_N = \Omega_1|_N$. Then there exist tubular neighborhoods U and V of N and a symplectomorphism $\rho : U \rightarrow V$ such that $\rho|_N$ is the identity.*

Proof. Since $d(\Omega_1 - \Omega_0) = 0$, by the generalized Poincaré lemma (see, e.g., Libermann and Marle [1987, p. 361]) there exists a tubular neighborhood W of N and a 1-form α on W such that $\Omega_1 - \Omega_0 = d\alpha$ and $\alpha|_N = 0$. Now for $t \in \mathbb{R}$ set $\Omega_t = \Omega_0 + t(\Omega_1 - \Omega_0)$; Ω_t is nondegenerate on an open subset W_1 of $W \times \mathbb{R}$ containing $N \times \mathbb{R}$. Let X be the vector field on W_1 defined by

$$X(m, t) \lrcorner \Omega_t(m) = -\alpha(m),$$

where \lrcorner denotes the left interior product. For a point m , consider the integral curve $t \rightarrow \phi_m(t)$ of X through $(m, 0)$ and regard the domain of $\phi : (m, t) \rightarrow \phi_m(t)$ as an open subset W_2 of $W \times \mathbb{R}$ with $W_2 \subset W_1$. Since $\alpha|_N = 0$, X restricted to $N \times \mathbb{R}$ vanishes and hence $N \times \mathbb{R} \subset W_2$. Now since $[0, 1]$ is compact, any point $m \in N$ has a neighborhood $U_m \subset M$

such that $U_m \times [0, 1] \subset W_2$. Let $U = \cup_{m \in N} U_m$. For $(m, t_0) \in U \times [0, 1]$ we compute

$$\frac{d}{dt}(\phi_m(t)^* \Omega_t)(m) \Big|_{t=t_0} = \phi_m(t_0)^* \left(\mathcal{L}_X \Omega_{t_0} + \frac{d}{dt} \Omega_t \Big|_{t=t_0} \right) (m) = 0$$

since

$$\mathcal{L}_X \Omega_t = X \lrcorner d\Omega_t + d(X \lrcorner \Omega_t) = -d\alpha = \Omega_0 - \Omega_1, \quad \text{and} \quad \frac{d}{dt} \Omega_t = \Omega_1 - \Omega_0.$$

Thus if $\rho : U \longrightarrow V = \rho(U) \subset W$ is the diffeomorphism determined by $(\rho(m), 1) = \phi_m(1)$, then $\rho^* \Omega_1 = \Omega_0$ and $\rho(m) = m$ for $m \in N$. ■

As a corollary we now have the classical theorem of Darboux.

Theorem 1.3 *Given (M^{2n}, Ω) symplectic and $m \in M$, there exist a neighborhood U of m and local coordinates $(x^1, \dots, x^n, y^1, \dots, y^n)$ on U such that $\Omega = \sum dx^i \wedge dy^i$.*

Proof. Let $(u^1, \dots, u^n, v^1, \dots, v^n)$ be local coordinates on a neighborhood of m such that $\frac{\partial}{\partial u^i}(m)$ and $\frac{\partial}{\partial v^i}(m)$ form a symplectic frame at the point m . Set $\Omega_0 = \Omega$ and $\Omega_1 = \sum du^i \wedge dv^i$. Then Ω_0 and Ω_1 agree at m . Now constructing ρ as in the previous theorem, $x^i = u^i \circ \rho$ and $y^i = v^i \circ \rho$ form the desired coordinates. ■

We remark that one can easily choose the “Darboux” coordinates such that $\frac{\partial}{\partial y^1}(m)$ is any preassigned vector X at m and that $x^i = \text{const}$ defines a Lagrangian submanifold. Thus we see that given a point m and a tangent vector X at m there exists a Lagrangian submanifold through the point and tangent to X as we remarked in the last section.

There are more general versions of the Darboux theorem, and it seems worthwhile to state the following two theorems here. We refer the reader to the book of S. Sternberg [1983, Chapter III, Section 6] for proofs and as a more classical reference to these results.

Theorem 1.4 *Let Ω be a closed 2-form such that $\Omega^p \neq 0$ but $\Omega^{p+1} \equiv 0$. Then about every point there exist local coordinates $(x^1, \dots, x^{n-p}, y^1, \dots, y^p)$ such that $\Omega = dx^1 \wedge dy^1 + \dots + dx^p \wedge dy^p$.*

Theorem 1.5 *Let $\omega \neq 0$ be a 1-form such that $d\omega^p \neq 0$ but $\omega \wedge (d\omega)^p \equiv 0$. Then there exist local coordinates $(x^1, \dots, x^{n-p}, y^1, \dots, y^p)$ such that $\omega = x^1 dy^1 + \dots + x^p dy^p$. If $\omega \wedge (d\omega)^p \neq 0$ but $d\omega^{p+1} \equiv 0$, then there exist local coordinates $(x^1, \dots, x^{n-p}, y^1, \dots, y^p)$ such that $\omega = x^1 dy^1 + \dots + x^p dy^p + dx^{p+1}$.*

We now prove the theorem of Weinstein that locally a symplectic manifold is the cotangent bundle of a Lagrangian submanifold.

Theorem 1.6 *If L is a Lagrangian submanifold of a symplectic manifold (M, Ω) , then there exists a neighborhood of L in M that is symplectomorphic to a neighborhood of the zero section in T^*L .*

Proof. Let $T_L M$ be the restriction of the tangent bundle TM to L and \mathbf{E} a Lagrangian complement of TL in $T_L M$. Such a vector bundle \mathbf{E} exists but is by no means unique; e.g., relative to an associated metric as described above, \mathbf{E} could be taken as the normal bundle of L . Define $j : \mathbf{E} \rightarrow T^*L$ by

$$j(\zeta)(X) = \Omega(\zeta, X),$$

where $\zeta \in \mathbf{E}_m$ and $X \in T_m L$, $m \in L$. Moreover there exist a tubular neighborhood \mathcal{U} of L in M and a diffeomorphism ϕ of \mathcal{U} onto $\phi(\mathcal{U}) \subset \mathbf{E}$ such that $\phi|_L$ is the zero section and identifying $T_{\phi(m)}\mathbf{E}_m$ with \mathbf{E}_m ,

$$\phi_*(m)|_{\mathbf{E}_m} = id|_{\mathbf{E}_m}, \quad m \in L$$

(see e.g., Libermann and Marle [1987, p. 358], or in the Riemannian case use the inverse of the exponential map). Then $j \circ \phi$ is a diffeomorphism of \mathcal{U} onto the open subset $j(\phi(\mathcal{U}))$ of T^*L whose restriction to L is the zero section, $s_0 : L \rightarrow L' \subset T^*L$. Moreover, $(j \circ \phi)_*(m)$ maps the complementary Lagrangian subspaces $T_m L$ and \mathbf{E}_m onto $T_{s_0(m)}L'$ and $T_{s_0(m)}(T_m^*L)$ respectively. But $T_{s_0(m)}L'$ and $T_{s_0(m)}(T_m^*L)$ are complementary Lagrangian subspaces with respect to the symplectic form $d\beta$ on T^*L . Now identifying L and L' , the restriction of $(j \circ \phi)_*(m)$ to $T_m L$ is just the identity and since $\phi_*(m)|_{\mathbf{E}_m} = id|_{\mathbf{E}_m}$, $(j \circ \phi)_*(m)$ restricted to \mathbf{E}_m is j . In particular, $(j \circ \phi)_*(m)\zeta$ is vertical and $X \in T_m L$, so using the local expression $\sum dp^i \wedge dq^i$ of $d\beta$ on T^*L ,

$$d\beta((j \circ \phi)_*(m)\zeta, (j \circ \phi)_*(m)X) = j(\zeta)(X) = \Omega(\zeta, X).$$

The result now follows from Theorem 1.2. ■

1.4 Symplectomorphisms

Recall that a diffeomorphism $f : M \rightarrow M$ is a symplectomorphism if $f^*\Omega = \Omega$. A vector field X which generates a 1-parameter group of symplectomorphisms is called a *symplectic vector field*. Clearly $\mathcal{L}_X \Omega = 0$.

Theorem 1.7 *Let X be a symplectic vector field on (M, Ω) , g an associated metric, and J the corresponding almost complex structure. Then $X^i = J^{ik}\theta_k$ for some closed 1-form θ . Conversely, given a closed 1-form θ , $X^i = J^{ik}\theta_k$ defines a symplectic vector field.*

Proof. First note that $0 = \mathcal{L}_X \Omega = d(X \lrcorner \Omega)$ implies that $\theta = \frac{1}{2}(X \lrcorner \Omega)$ is a closed 1-form. Let T be the vector field given by $g(T, Y) = \theta(Y)$. Then $g(JT, Y) = -\theta(JY) = -\Omega(X, JY) = g(X, Y)$. Therefore $X^i = J^i_k T^k$ or $X^i = J^{ik}\theta_k$ as desired. Conversely, given θ closed, define X by $X^i = J^{ik}\theta_k$ from which $-\theta_l = J_{li}X^i = -J_{il}X^i$, i.e., $\theta = \frac{1}{2}(X \lrcorner \Omega)$. Therefore $\mathcal{L}_X \Omega = d(X \lrcorner \Omega) = 2d\theta = 0$. ■

Corollary 1.1 *For $f \in C^\infty(M)$, $J\nabla f$ is symplectic, where ∇f is the gradient of f . Conversely, given X symplectic, X is locally $J\nabla f$.*

In particular, X is locally the Hamiltonian vector field X_f . Compare this with the following classical treatment. Suppose that N is a level hypersurface of the function H on (M, Ω) , on which $dH \neq 0$. Then X_H is a nonzero tangent vector field that is in the direction of J of the normal direction: $g(Y, \nabla H) = YH = \Omega(X_H, Y) = g(X_H, JY) = -g(JX_H, Y)$, giving $X_H = J\nabla H$.

Finally, we prove a result of Hatakeyama [1966] that the group of symplectomorphisms acts transitively on a compact symplectic manifold.

Theorem 1.8 *The group of symplectomorphisms acts transitively on a compact symplectic manifold (M, Ω) .*

Proof. We first prove the result for a Darboux neighborhood \mathcal{U} about $p \in M$, i.e., we have local coordinates $(x^1, \dots, x^n, y^1, \dots, y^n)$ such that $\Omega = \sum dx^i \wedge dy^i$ and $x^i(p) = y^i(p) = 0$. Let $q(\neq p) \in \mathcal{U}$ with coordinates (a^i, b^i) and define a function f on \mathcal{U} by $f = \frac{1}{2} \sum (a^i y^i - b^i x^i)$. Then the vector field X defined by $X \lrcorner \Omega = 2df$ generates a 1-parameter group ϕ_t such that $\phi_1(p) = q$. Writing X as $X^i \frac{\partial}{\partial x^i} + X^{i*} \frac{\partial}{\partial y^i}$, we have $X \lrcorner \Omega = X^i dy^i - X^{i*} dx^i = a^i dy^i - b^i dx^i$. Thus $X = a^i \frac{\partial}{\partial x^i} + b^i \frac{\partial}{\partial y^i}$, and its integral curves have the form $x^i = a^i t, y^i = b^i t$. Strictly speaking, X is determined by $f \in C^\infty(M)$, where f equals $\frac{1}{2} \sum (a^i y^i - b^i x^i)$ on \mathcal{U} and vanishes outside some larger neighborhood; M compact then implies that ϕ_t is a diffeomorphism of M . Thus any two points in \mathcal{U} may be joined by a symplectomorphism. Now for $p, q \in M$ join them by a curve and cover it

by a finite number of Darboux neighborhoods $\mathcal{U}_\alpha, \alpha = 1, \dots, k$. Choose a sequence of points p_α such that $p_0 = p$, $p_k = q$ and $p_\alpha \in \mathcal{U}_\alpha \cap \mathcal{U}_{\alpha+1}$, and apply the above result. ■

For a generalization to symplectomorphisms mapping k points to k points, see Boothby [1969] or Kriegl–Michor [1997, p. 472].

2

Principal S^1 -bundles

A very important theorem in the geometry of contact manifolds, and the start of the modern theory, is the Boothby–Wang theorem, which states that a compact regular contact manifold is a principal circle bundle over a symplectic manifold of integral class. We will prove this result in Section 3.3. In preparation for this we review principal circle bundles in this chapter.

2.1 The set of principal S^1 -bundles as a group

Let P and M be C^∞ manifolds, $\pi : P \longrightarrow M$ a C^∞ map of P onto M , and G a Lie group acting on P to the right. Then (P, G, M) is called a *principal G -bundle* if

1. G acts freely on P ,
2. $\pi(p_1) = \pi(p_2)$ if and only if there exists $g \in G$ such that $p_1g = p_2$,
3. P is locally trivial over M , i.e., for every $m \in M$ there exists a neighborhood \mathcal{U} of m and a map $F_{\mathcal{U}} : \pi^{-1}(\mathcal{U}) \longrightarrow G$ such that for every $p \in \pi^{-1}(\mathcal{U})$ and $g \in G$, $F_{\mathcal{U}}(pg) = (F_{\mathcal{U}}(p))g$, and such

that the map $\psi : \pi^{-1}(\mathcal{U}) \longrightarrow \mathcal{U} \times G$ taking p to $(\pi(p), F_{\mathcal{U}}(p))$ is a diffeomorphism.

For a general reference to the theory of principal fiber bundles see Bishop and Crittenden [1964, Chapters 3 and 5], Kobayashi and Nomizu [1963–69, Chapter II].

We now turn to the case where $G = S^1$, in which case we say that P is a *principal circle bundle* over M and we study the group structure of the set $\mathcal{P}(M, S^1)$ of all principal circle bundles over M . Our treatment is based on Kobayashi [1956].

Given $P, P' \in \mathcal{P}(M, S^1)$ with projections π, π' , let $\Delta(P \times P') = \{(u, u') \in P \times P' \mid \pi(u) = \pi'(u')\}$. We say $(u_1, u'_1) \sim (u_2, u'_2)$ if there exists $s \in S^1$ such that $u_1 s = u_2$ and $u'_1 s^{-1} = u'_2$. Note that since S^1 is abelian, $u_3 = u_2 t = u_1 s t$, $u'_3 = u'_2 t^{-1} = u'_1 s^{-1} t^{-1} = u'_1 (st)^{-1}$.

Let $P + P' = \Delta(P \times P') / \sim$ and $\pi'' : P + P' \longrightarrow M$ the induced projection. S^1 acts on $\Delta(P \times P')$ by $(u, u')s = (us, u')$. Now if $(u_1, u'_1) \sim (u_2, u'_2)$, $u_1 t = u_2$ and $u'_1 t^{-1} = u'_2$, we have $u_2 s = u_1 t s = (u_1 s)t$. Therefore $(u_1 s, u'_1) \sim (u_2 s, u'_2)$ and hence S^1 acts on $P + P'$.

S^1 acts freely: Suppose $u''s = u'', u'' \in P + P'$ and suppose (u, u') represents u'' . Then $(u, u') \sim (us, u')$, so that $u's^{-1} = u'$ and hence $s = 1 \in S^1$.

S^1 acts transitively on fibers: Suppose $u''_1, u''_2 \in \pi''^{-1}(m)$ and $(u_1, u'_1), (u_2, u'_2)$ are representatives. Then $u_2 = u_1 s$, $u'_2 = u'_1 s'$, $s, s' \in S^1$. Now $(u_2, u'_2) \sim (u_2 s', u'_1) = (u_1 s s', u'_1) = (u_1, u'_1) s s'$ and hence $u''_2 = u''_1 s s'$.

$P + P'$ is locally trivial: If $F_{\mathcal{U}}(u) = g, F'_{\mathcal{U}}(u') = g'$, set $F''_{\mathcal{U}}(u, u') = gg'$. Then $F''_{\mathcal{U}}(us, u') = gsg' = gg's$.

Theorem 2.1 *Under the operation $+$, $\mathcal{P}(M, S^1)$ is an abelian group.*

Proof. Let P_0 be the trivial bundle and $\alpha : P \longrightarrow P + P_0$ defined by $\alpha(u) = [(u, (\pi(u), 1))]$. Then α is a bundle isomorphism:

$$\begin{aligned} \alpha(us) &= [(us, (\pi(u), 1))] = [(u, (\pi(u), 1))s] \\ &= [(u, (\pi(u), 1))]s = \alpha(u)s; \\ \alpha^{-1}([(u, (\pi(u), g))]) &= \alpha^{-1}([(ug^{-1}, (\pi(u), 1))]) = ug^{-1}. \end{aligned}$$

Let $-P$ be a manifold diffeomorphic to P and $-u$ the point corresponding to u . Define $-\pi : -P \rightarrow M$ by $-\pi(-u) = \pi(u)$. S^1 acts on $-P$ by $(-u)s = -(us^{-1})$. Then $-P \in \mathcal{P}(M, S^1)$. Now let $(u_1, -u_2) \in \Delta(P \times -P)$; then there exists a unique $s \in S^1$ such that $u_1 = u_2s$. Let $\alpha : P + (-P) \rightarrow P_0$ be defined by $\alpha([(u_1, -u_2)]) = (\pi(u_1), s)$. Then α is a bundle isomorphism.

Let $\Delta(P \times P' \times P'') = \{(u, u', u'') | \pi(u) = \pi'(u') = \pi''(u'')\}$ and define the equivalence \sim by $(u, u', u'') \sim (us, u's'^{-1}s', u''s'^{-1})$. Then $\Delta(P \times P' \times P'') / \sim$ is naturally isomorphic to $(P + P') + P''$, $((u's'^{-1}, us)s', u''s'^{-1})$, and to $P + (P' + P'')$, $(us, (u's', u''s'^{-1})s^{-1})$. S^1 acts on $\Delta(P \times P' \times P'')$ by $(u, u', u'')s = (us, u's', u'')$. Now if $(u_1, u'_1, u''_1) \sim (u_2, u'_2, u''_2)$, then $u_2 = u_1t, u'_2 = u'_1t^{-1}t', u''_2 = u''_1t'^{-1}$. Then $u_2s = u_1ts = (u_1s)t$ so that the right action preserves \sim .

Finally, $P + P'$ is isomorphic to $P' + P$ by $[(u, u')] \longleftrightarrow [(u', u)]$, $(us, u') \sim (u, u's)$. ■

Let G_m be the cyclic subgroup of S^1 of order m and $P \in \mathcal{P}(M, S^1)$. Since S^1 acts on P on the right, so does G_m . Then P/G_m is a principal bundle over M with group S^1/G_m . But $S^1/G_m \cong S^1$ and hence we can consider $P/G_m \in \mathcal{P}(M, S^1)$. More precisely: Let $[u]$ be an element of P/G_m that is represented by $u \in P$. Define the action of S^1 on P/G_m by setting $[u]s = [us']$, where $s = s'^m$. This definition is independent of the choice of u and s' . For if $g \in G_m$, then $[ug]s = [ugs'] = [us'g] = [us'] = [u]s$, and if $s''^m = s$, then $(s'^{-1}s'')^m = 1$ so that $s'^{-1}s'' \in G_m$ and hence $[us''] = [us's'^{-1}s''] = [us']$.

Theorem 2.2 *Let P , G_m and P/G_m be as above. Then $P/G_m \cong m \cdot P$.*

Proof. From the definition above it follows by induction that $m \cdot P$ can be defined directly by

$$\Delta(P \times \cdots \times P) = \{(u_1, \dots, u_m) \in P \times \cdots \times P | \pi(u_1) = \cdots = \pi(u_m)\},$$

two elements of which, say (u_1, \dots, u_m) and (u_1s_1, \dots, u_ms_m) , are equivalent if and only if $s_1 \cdots s_m = 1$. The quotient space of $\Delta(P \times \cdots \times P)$ by this relation is $m \cdot P$. The action of S^1 on $m \cdot P$ is given by $[(u_1, \dots, u_m)]s = [(u_1s, u_2, \dots, u_m)]$. Define $\phi : P/G_m \rightarrow m \cdot P$ by $\phi([u]) = [(u, \dots, u)]$, which is independent of the choice of u , for if $g \in G_m$, $g^m = 1$, then $\phi([ug]) = [(ug, \dots, ug)] = [(u, \dots, u)]$. Now take $s \in S^1$ and s' such that $s'^m = s$. Then $\phi([u]s) = \phi([us']) =$

$[(us', \dots, us')] = [(us, u, \dots, u)] = [(u, \dots, u)]s = (\phi([u]))s$. Therefore ϕ is a bundle isomorphism of P/G_m onto $m \cdot P$ \blacksquare

Corollary 2.1 *If P is simply connected and $m > 1$, then there is no bundle $P' \in \mathcal{P}(M, S^1)$ such that $P = m \cdot P'$.*

Proof. Suppose that P' exists. Then $P \cong P'/G_m$ and so P' is a covering space of P . Since P is simply connected, this can happen only if $m = 1$. \blacksquare

A principal bundle may also be thought of as an equivalence class of principal coordinate bundles that are given by their transition functions. Let $\{\mathcal{U}_i\}$ be a differentiably simple open cover of M (i.e., $\{\mathcal{U}_i\}$ is locally finite, each \mathcal{U}_i has compact closure and any nonempty finite intersection is diffeomorphic to an open cell of \mathbb{R}^n). With respect to this cover let $f_{ij} : \mathcal{U}_i \cap \mathcal{U}_j \rightarrow S^1$ be the transition functions of a bundle $P \in \mathcal{P}(M, S^1)$. The f_{ij} are defined by $f_{ij}(\pi(p)) = F_{U_i}(p)(F_{U_j}(p))^{-1}$. Then $f_{ij} \in \Gamma(\mathcal{U}_i \cap \mathcal{U}_j, \mathcal{S}^1)$, the set of all sections over $\mathcal{U}_i \cap \mathcal{U}_j$ with coefficients in \mathcal{S}^1 , the sheaf of germs of local C^∞ maps from M into S^1 . Thus $f = \{f_{ij}\}$ is a cochain of M . Now $f_{ik} = f_{ij}f_{jk}$. Thus $f_{i_0i_1i_2} = \delta f_{i_0i_1} = f_{i_0i_1}f_{i_0i_2}^{-1}f_{i_1i_2} = f_{i_0i_2}f_{i_0i_1}^{-1} = 1$ and f is a cocycle. Now P and P' are equivalent if and only if $P - P'$ is the trivial bundle, so P and P' are the same here if and only if $f - f'$ is a coboundary, where f' is the cocycle of P' . Therefore $\mathcal{P}(M, S^1) \cong H^1(M, S^1)$.

The natural short exact sequence $0 \rightarrow \mathbb{Z} \rightarrow \mathbb{R} \rightarrow S^1 \rightarrow 0$ induces a short exact sequence of the corresponding sheaves $0 \rightarrow \mathcal{Z} \rightarrow \mathcal{R} \rightarrow S^1 \rightarrow 0$. From this we get the cohomology sequence

$$\dots \rightarrow H^1(M, \mathcal{R}) \rightarrow H^1(M, \mathcal{S}^1) \rightarrow H^2(M, \mathcal{Z}) \rightarrow H^2(M, \mathcal{R}) \rightarrow \dots$$

Now let $\{\phi_i\}$ be a partition of unity subordinate to $\{\mathcal{U}_i\}$ with $\phi_i|_{M-\mathcal{V}_i} = 0$, $\bar{\mathcal{V}}_i \subset \mathcal{U}_i$. Let $\{\beta_{ijk}\} \in Z^2(M, \mathcal{R})$. Consider $\alpha_{ij} = \sum_k \phi_k \beta_{ijk}$. If in a neighborhood of $m \in \mathcal{U}_i \cap \mathcal{U}_j$, β_{ijk} is not defined, we have $\phi_k(m') = 0$ for every m' in the neighborhood. Hence we see that α_{ij} is defined on $\mathcal{U}_i \cap \mathcal{U}_j$. Now

$$\begin{aligned} \delta\{\alpha_{ij}\} &= \{\alpha_{ij} - \alpha_{ik} + \alpha_{jk}\} = \left\{ \sum_l \phi_l (\beta_{ijl} - \beta_{ikl} + \beta_{jkl}) \right\} \\ &= \left\{ \sum_l \phi_l \beta_{ijk} \right\} = \{\beta_{ijk}\}. \end{aligned}$$

This can be done for other-dimensional cocycles as well. Hence $Z^i(M, \mathcal{R}) = B^i(M, \mathcal{R})$, $i > 0$. Thus in particular we have that

$$H^1(M, \mathcal{R}) = H^2(M, \mathcal{R}) = 0.$$

On the other hand, the C^∞ maps of M into \mathbb{Z} are just the constant integer functions and hence the corresponding cohomology groups must be isomorphic, in particular $H^2(M, \mathcal{Z}) = H^2(M, \mathbb{Z})$. Thus we finally have the isomorphisms

$$H^1(M, \mathcal{S}^1) \cong H^2(M, \mathbb{Z}), \quad \mathcal{P}(M, \mathcal{S}^1) \cong H^2(M, \mathbb{Z}).$$

For example, it is well known that for complex projective space $\mathbb{C}P^n$, $H^2(\mathbb{C}P^n, \mathbb{Z}) \cong \mathbb{Z}$. Thus by the above isomorphism, $\mathcal{P}(\mathbb{C}P^n, \mathcal{S}^1) \cong \mathbb{Z}$. The most famous example of a principal circle bundle over $\mathbb{C}P^n$ is the Hopf fibration of an odd-dimensional sphere S^{2n+1} . Suppose that $S^{2n+1} \in \mathcal{P}(\mathbb{C}P^n, \mathcal{S}^1)$ corresponds to $k \in \mathbb{Z}$. Then since S^{2n+1} is simply connected, $k = \pm 1$ by the corollary. Therefore we finally have the following result.

Theorem 2.3 $\mathcal{P}(\mathbb{C}P^n, \mathcal{S}^1) \cong \mathbb{Z}$ with S^{2n+1} corresponding to 1 for a proper orientation of $\mathbb{C}P^n$.

2.2 Connections on a principal bundle

A *connection* on a principal G -bundle (P, G, M) is a C^∞ distribution H on P such that

1. $T_p P = H_p \oplus V_p$, $V_p = \ker \pi_*$,
2. $R_{g*}(H_p) = H_{pg}$, R_g being right translation.

Vectors in H_p are said to be *horizontal*, and for $t \in T_p P$ we denote its horizontal part by Ht . The map $\pi_*|_{H_p}$ is one-to-one and hence $\pi_*(H_p) = T_{\pi(p)} M$. Thus given a vector field X on M there exists a unique vector field \tilde{X} on P such that $\tilde{X}(p) \in H_p$ and $\pi_* \tilde{X}(p) = X(\pi(p))$, i.e., \tilde{X} is π -related to X . \tilde{X} is called the *horizontal lift* of X . The following two properties of horizontal lifts follow easily from the definitions:

$$\widetilde{[X, Y]} = H[\tilde{X}, \tilde{Y}], \quad R_{g*} \tilde{X} = \tilde{X}.$$

A p -form ω on P is *vertical* (resp. *horizontal*) if $\omega(t_1, \dots, t_p) = 0$ when one or more of the t_i 's is horizontal (resp. vertical).

Now regard $p \in P$ as a map of $G \longrightarrow P$ by $p(g) = pg$. Let \mathfrak{g} denote the Lie algebra of G and define a Lie algebra homomorphism of \mathfrak{g} into a Lie algebra $\bar{\mathfrak{g}}$ of vector fields on P by $\bar{X} = (p_*X)(e)$. The map $p : G \longrightarrow \pi^{-1}(\pi(p))$ is a diffeomorphism, so given $t \in V_p$, let $X(e) = p_*^{-1}t$. Then there exists $\bar{X} \in \bar{\mathfrak{g}}$ such that $\bar{X}(p) = t$. On the other hand, given $\bar{X} \in \bar{\mathfrak{g}}$, $\pi_*\bar{X}(p) = \pi_*(p_*X)(e) = (\pi \circ p)_*X(e) = 0$, since $\pi \circ p$ is a constant map. Thus \bar{X} is vertical.

Given a connection H on (P, G, M) define a \mathfrak{g} -valued 1-form ϕ on P by $\phi(t) = X \in \mathfrak{g}$ where $\bar{X}(p)$ is the vertical part of t . ϕ is called the *connection form* of H . The following lemmas are well known and their proofs can be found in the references (e.g., Bishop and Crittenden [1964, pp. 76–77], Kobayashi and Nomizu [1963–69, Volume I, p. 64]).

Lemma 2.1 $\phi \circ R_{g*} = \text{Ad } g^{-1} \circ \phi$, i.e., ϕ is equivariant.

Lemma 2.2 If ϕ is a \mathfrak{g} -valued C^∞ equivariant 1-form such that $\phi(\bar{X}(p)) = X$, then there exists a unique connection H whose 1-form is ϕ .

Given a \mathfrak{g} -valued p -form σ and a \mathfrak{g} -valued q -form ω , their bracket is defined by

$$\begin{aligned} & [\sigma, \omega](X_1, \dots, X_{p+q}) \\ &= \frac{1}{(p+q)!} \sum_{(i_1, \dots, i_{p+q})} \text{sgn}(i_1, \dots, i_{p+q}) [\sigma(X_{i_1}, \dots, X_{i_p}), \omega(X_{i_{p+1}}, \dots, X_{i_{p+q}})]. \end{aligned}$$

Let ω be a p -form on P and define a $(p+1)$ -form $D\omega$ by

$$D\omega(t_1, \dots, t_{p+1}) = d\omega(Ht_1, \dots, Ht_p).$$

Clearly $D\omega$ is horizontal. If ϕ is the connection form of H , $\Phi = D\phi$ is called the *curvature form* of H . Φ is equivariant as can be seen as follows:

$$\begin{aligned} \Phi(R_{g*}t_1, R_{g*}t_2) &= d\phi(HR_{g*}t_1, HR_{g*}t_2) = d\phi(R_{g*}Ht_1, R_{g*}Ht_2) \\ &= -\frac{1}{2}\phi(R_{g*}[Ht_1, Ht_2]) = -\frac{1}{2}\text{Ad } g^{-1}\phi([Ht_1, Ht_2]) \\ &= \text{Ad } g^{-1}\Phi(t_1, t_2). \end{aligned}$$

We now have the structural equation; again for the proof see the references (e.g., Bishop and Crittenden [1964, p. 81], Kobayashi and Nomizu [1963–69, Volume I, p. 77]).

Theorem 2.4 $d\phi = -\frac{1}{2}[\phi, \phi] + \Phi$.

Let $P \in \mathcal{P}(M, S^1)$ and note that the Lie algebra of S^1 is \mathbb{R} with the trivial bracket operation. Thus if η is a connection form on P , then $[\eta, \eta] = 0$, and if Φ is the curvature, the structural equation is simply $d\eta = \Phi$.

Again since S^1 is abelian, for $X, Y \in T_u P$ and $s \in S^1$ we have

$$\Phi(R_{s*}X, R_{s*}Y) = Ad s^{-1}\Phi(X, Y) = \Phi(X, Y).$$

Therefore there exists a unique 2-form Ω on M such that $\Phi = \pi^*\Omega$. Now $\pi^*(d\Omega) = d\Phi = 0$ and hence Ω is a closed 2-form on M .

If now η' is another connection, then as before, $(\eta - \eta') \circ R_{s*} = \eta - \eta'$. Therefore there exists a unique 1-form β on M such that $\pi^*\beta = \eta - \eta'$. Now $\pi^*d\beta = d(\eta - \eta') = \Phi - \Phi' = \pi^*\Omega - \pi^*\Omega'$ and hence $d\beta = \Omega - \Omega'$. Thus the cohomology class of Ω is independent of the choice of the connection form and is called the *characteristic class* of P ; again see Kobayashi [1956]. Since the transition functions are mappings from $\mathcal{U}_i \cap \mathcal{U}_j$ into S^1 , they can be considered as real-valued functions (mod 1), and it can then be shown that Ω is integral, i.e., $\int_c \Omega = \text{integer}$ for any finite singular cocycle c with integer coefficients. This gives a homomorphism of $\mathcal{P}(M, S^1)$ onto the integral classes of the second de Rham cohomology.

We end this chapter with the following theorem of Kobayashi [1963], which should again be compared with the isomorphism $\mathcal{P}(M, S^1) \cong H^2(M, \mathbb{Z})$.

Theorem 2.5 *Let Ω be a 2-form on M representing an element of $H^2(M, \mathbb{Z})$. Then there exist a principal circle bundle P and a connection form η on P such that $d\eta = \pi^*\Omega$.*

Proof. Let P be the principal bundle corresponding to Ω and η' a connection form on P such that the closed 2-form Ω' defined by $d\eta' = \pi^*\Omega'$ is cohomologous to Ω . Let β be a 1-form on M such that $\Omega - \Omega' = d\beta$. Now set $\eta = \eta' + \pi^*\beta$. Then $\pi^*\beta$ is horizontal and equivariant, and hence η is a connection form on P and $d\eta = \pi^*\Omega$ as desired. ■

3

Contact Manifolds

In this chapter we give the basic definitions and properties concerning contact manifolds both as given by a global contact form and as a contact structure in the wider sense. We then give many examples of contact manifolds, a discussion of the celebrated Boothby–Wang fibration, and a discussion of the Weinstein conjecture.

3.1 Definitions

By a *contact manifold* we mean a C^∞ manifold M^{2n+1} together with a 1-form η such that $\eta \wedge (d\eta)^n \neq 0$. In particular, $\eta \wedge (d\eta)^n \neq 0$ is a volume element on M , so that a contact manifold is orientable. Also $d\eta$ has rank $2n$ on the Grassmann algebra $\bigwedge T_m^*M$ at each point $m \in M$, and thus we have a 1-dimensional subspace, $\{X \in T_m M \mid d\eta(X, T_m M) = 0\}$, on which $\eta \neq 0$ and which is complementary to the subspace defined by $\eta = 0$. Therefore choosing ξ_m in this subspace normalized by $\eta(\xi_m) = 1$, we have a global vector field ξ satisfying

$$d\eta(\xi, X) = 0, \quad \eta(\xi) = 1.$$

ξ is called the *characteristic vector field* or *Reeb vector field* (Reeb [1952]) of the contact structure η . Computing Lie derivatives by the formula

$\mathcal{L}_\xi = d \circ (\xi \lrcorner) + (\xi \lrcorner) \circ d$, we have immediately that

$$\mathcal{L}_\xi \eta = 0, \quad \mathcal{L}_\xi d\eta = 0.$$

We denote by \mathcal{D} the *contact distribution* or *subbundle* defined by the subspaces $\mathcal{D}_m = \{X \in T_m M : \eta(X) = 0\}$. Roughly speaking, the meaning of the contact condition, $\eta \wedge (d\eta)^n \neq 0$, is that the contact subbundle is as far from being integrable as possible; in particular, \mathcal{D} rotates as one moves around on the manifold. For a subbundle defined by a 1-form η to be integrable it is necessary and sufficient that $\eta \wedge (d\eta) \equiv 0$. In contrast, we shall see in Chapter 5 that for a contact manifold M^{2n+1} , the maximum dimension of an integral submanifold of \mathcal{D} is only n . A one-dimensional integral submanifold of \mathcal{D} will be called a *Legendre curve*, especially to avoid confusion with an integral curve of the vector field ξ .

A contact structure is *regular* if ξ is regular as a vector field, that is, every point of the manifold has a neighborhood such that any integral curve of the vector field passing through the neighborhood passes through only once (cf. Palais [1957]). Two well-known examples of nonregular vector fields on surfaces are the irrational flow on a torus and the flow around a Möbius band.

We now prove the classical theorem of Darboux. It is, of course, a special case of Theorem 1.5, but it is worthwhile to give a short proof here and to note that we will sometimes refer to a *Darboux coordinate system* as one in which the contact form is given locally in the notation of Theorem 3.1.

Theorem 3.1 *About each point of a contact manifold (M^{2n+1}, η) there exist local coordinates $(x^1, \dots, x^n, y^1, \dots, y^n, z)$ with respect to which*

$$\eta = dz - \sum_{i=1}^n y^i dx^i.$$

Proof. In some coordinate neighborhood choose a $2n$ -ball transverse to ξ ; $d\eta$ is symplectic on this ball, and hence there exist local coordinates $(x^1, \dots, x^n, y^1, \dots, y^n, u)$ such that $d\eta = \sum dx^i \wedge dy^i$. Now $d(\eta + \sum y^i dx^i) = 0$ so that $\eta + \sum y^i dx^i = df$ for some function f , i.e., $\eta = df - \sum y^i dx^i$. Now $\eta \wedge (d\eta)^n = df \wedge dx^1 \wedge \dots \wedge dx^n \wedge dy^1 \wedge \dots \wedge dy^n \neq 0$. Therefore df is independent of $dx^1, \dots, dx^n, dy^1, \dots, dy^n$, and hence we can regard x^i, y^i and $z = f$ as a coordinate system. ■

In a Darboux coordinate system, $\xi = \frac{\partial}{\partial z}$. For if $\xi = a \frac{\partial}{\partial z} + b^i \frac{\partial}{\partial x^i} + c^i \frac{\partial}{\partial y^i}$, then $1 = \eta(\xi) = a - b^i y^i$. Also $0 = d\eta(\xi, \frac{\partial}{\partial x^i}) = \sum dx^i \wedge dy^i(\xi, \frac{\partial}{\partial x^i})$ gives $c^i = 0$. Similarly, $0 = d\eta(\xi, \frac{\partial}{\partial y^i})$ gives $b^i = 0$. Thus $a = 1$ and $\xi = \frac{\partial}{\partial z}$.

A diffeomorphism f of M^{2n+1} or between open subsets of \mathbb{R}^{2n+1} with the contact structure of the Darboux form of Theorem 3.1 is called a *contact transformation* if $f^*\eta = \tau\eta$ for some nonvanishing function τ on the domain of f . If $\tau \equiv 1$, f is called a *strict contact transformation*.

There is also the notion of a *contact structure in the wider sense*, often called simply a “contact structure” by many authors, which can be defined in a number of ways. For example, a contact manifold in the wider sense is a manifold with a differentiable structure modeled on the pseudogroup of contact transformations on \mathbb{R}^{2n+1} (J. Gray [1959]), i.e., in the overlap of coordinate neighborhoods the transition functions preserve the Darboux form to within a nonvanishing function multiple. An alternate approach is to put the emphasis on the field of $2n$ -planes \mathcal{D} and to define the structure as a hyperplane field defined locally by a contact form. In the overlap of coordinate neighborhoods $\mathcal{U} \cap \mathcal{U}'$, we have $\eta' = f^*\eta = \tau\eta$ and hence $d\eta' = f^*d\eta = d\tau \wedge \eta + \tau d\eta$ from which

$$\eta' \wedge (d\eta')^n = \tau^{n+1} \eta \wedge (d\eta)^n \neq 0.$$

Let M^{2n+1} be a contact manifold in the wider sense. On a coordinate neighborhood \mathcal{U}_α choose coordinates (x^1, \dots, x^{2n+1}) such that $\eta_\alpha \wedge (d\eta_\alpha)^n = \lambda_\alpha dx^1 \wedge \dots \wedge dx^{2n+1}$ with $\lambda_\alpha > 0$. Similarly, on a neighborhood \mathcal{U}_β choose coordinates (y^1, \dots, y^{2n+1}) such that $\eta_\beta \wedge (d\eta_\beta)^n = \lambda_\beta dy^1 \wedge \dots \wedge dy^{2n+1}$ with $\lambda_\beta > 0$. Now on $\mathcal{U}_\alpha \cap \mathcal{U}_\beta$, $\eta_\alpha = \tau_{\alpha\beta} \eta_\beta$ and hence $\eta_\alpha \wedge (d\eta_\alpha)^n = \tau_{\alpha\beta}^{n+1} \eta_\beta \wedge (d\eta_\beta)^n$. Therefore $\tau_{\alpha\beta}^{n+1} \lambda_\beta |\frac{\partial y^i}{\partial x^j}| = \lambda_\alpha$. From this one may easily obtain the following results (see also J. Gray [1959], Sasaki [1965], Stong [1974]).

Theorem 3.2 *Let M^{2n+1} be a contact manifold in the wider sense. If n is odd, then M^{2n+1} is orientable.*

Theorem 3.3 *Let M^{2n+1} be a contact manifold in the wider sense. If n is even and M^{2n+1} is orientable, then M^{2n+1} is a contact manifold.*

We also have the following theorem of Sasaki [1965].

Theorem 3.4 *Let M^{2n+1} be a contact manifold in the wider sense which is not a contact manifold in the restricted sense. Then its 2-fold covering manifold is a contact manifold in the restricted sense.*

Proof. Let $\{\mathcal{U}_\alpha\}$ be an open cover of M^{2n+1} by coordinate charts. Recall the local contact forms η_α above and their transition $\eta_\alpha = \tau_{\alpha\beta}\eta_\beta$. Consider the 2-fold covering $\pi : \tilde{M}^{2n+1} \longrightarrow M^{2n+1}$ given as follows: \tilde{M}^{2n+1} is the union of the sets $\{\mathcal{U}_\alpha \times \mathbb{Z}_2\}$ with the following equivalence relation, where \mathbb{Z}_2 is the set of integers ± 1 . Let ϵ_α denote $+1$ or -1 . Two elements $(p_\alpha, \epsilon_\alpha) \in \mathcal{U}_\alpha \times \mathbb{Z}_2$ and $(p_\beta, \epsilon_\beta) \in \mathcal{U}_\beta \times \mathbb{Z}_2$ are equivalent if $p_\alpha = p_\beta$ and $\epsilon_\alpha = \text{sgn}(\tau_{\alpha\beta}(p_\beta))\epsilon_\beta$.

Now define local contact forms on $\mathcal{U}_\alpha \times \pm 1$ by $\eta_{(\alpha, \epsilon_\alpha)} = \epsilon_\alpha \pi^* \eta_\alpha$. Then $\eta_{(\alpha, \epsilon_\alpha)} = \epsilon_\alpha \pi^* \eta_\alpha = \epsilon_\alpha \pi^*(\tau_{\alpha\beta}\eta_\beta) = \epsilon_\alpha(\tau_{\alpha\beta} \circ \pi) \pi^* \eta_\beta = \epsilon_\alpha \epsilon_\beta (\tau_{\alpha\beta} \circ \pi) \eta_{(\beta, \epsilon_\beta)}$ and $\epsilon_\alpha \epsilon_\beta (\tau_{\alpha\beta} \circ \pi) > 0$. The local forms $\eta_{(\alpha, \epsilon_\alpha)}$ can now be used to construct a global contact form on \tilde{M}^{2n+1} . ■

In particular, a connected and simply connected contact manifold in the wider sense is a contact manifold in the restricted sense; for an alternate approach to this idea see Monna [1983].

The name contact (Berührungstransformation) seems to be due to Sophus Lie [1890] and is natural in view of the simple example of Huygens' principle (Huygens [1690]; see also Mac Lane [1968, part II, p. 83]). Consider \mathbb{R}^2 with coordinates (x, y) . The classical notion of a "line element" of \mathbb{R}^2 is a point together with a nonvertical line through the point. Thus a line element may be regarded as a point in \mathbb{R}^3 determined by the point and the slope p of the line. Given a smooth curve C in the plane without vertical tangents, say $y = f(x)$, its tangent lines determine a curve in \mathbb{R}^3 with coordinates (x, y, p) which is a Legendre curve of the contact form $\eta = dy - p dx$. If now C is a wave front, by Huygens' principle the new wave front C_t at time t is the envelope of the circular waves centered at all the points of C , say of radius t taking the velocity of propagation to be 1. Corresponding to a point (x, y) on C , the point (\bar{x}, \bar{y}) on C_t lies on both the normal line and the circle of radius t centered at (x, y) , i.e., $\bar{y} - y = -\frac{1}{p}(\bar{x} - x)$ and $(\bar{x} - x)^2 + (\bar{y} - y)^2 = t^2$. Thus $(\bar{x} - x)^2 = \frac{p^2 t^2}{p^2 + 1}$, so depending on the direction of propagation, e.g., choosing the negative root, the transformation of \mathbb{R}^3 mapping (x, y, p) to

$$\bar{x} = x - \frac{pt}{\sqrt{p^2 + 1}}, \quad \bar{y} = y + \frac{t}{\sqrt{p^2 + 1}}, \quad \bar{p} = p,$$

maps C to C_t . A simple calculation shows that $d\bar{y} - \bar{p} d\bar{x} = dy - p dx$, and so the transformation is a contact transformation. Moreover, tangent wave fronts (curves) are mapped to tangent wave fronts (curves) and hence the name “contact”.

Beginning with Chapter 4, the main thrust of this book will be the Riemannian geometry of contact and symplectic manifolds. For the reader interested in purely topological considerations and constructions of contact manifolds (in the wider sense) we refer to the expository writings of H. Geiges [2006], [2008].

3.2 Examples

In [1971] J. Martinet proved that every compact orientable 3-manifold carries a contact structure. However, we now have the following theorem of J. Gonzalo [1987] showing that there are three independent contact structures.

Theorem 3.5 *Every closed orientable 3-manifold has a parallelization by three contact forms.*

Before turning to some detailed examples we should mention that in contrast to Martinet’s result, there exist $(2n+1)$ -dimensional manifolds, $n \geq 2$, with no contact structure even in the wider sense. In particular, Stong [1974] showed that for every $n \geq 2$ there is a closed oriented connected manifold of dimension $2n+1$ with no contact structure in the wider sense. One such manifold is $SU(3)/SO(3)$ for $n = 2$ and $(SU(3)/SO(3)) \times S^{2n-4}$ for $n > 2$.

3.2.1 \mathbb{R}^{2n+1}

In effect, we have already seen that $\mathbb{R}^{2n+1}(x^1, \dots, x^n, y^1, \dots, y^n, z)$ with the Darboux form $\eta = dz - \sum_{i=1}^n y^i dx^i$ is a contact manifold. The characteristic vector field ξ is $\frac{\partial}{\partial z}$ and the contact subbundle \mathcal{D} is spanned by

$$X_i = \frac{\partial}{\partial x^i} + y^i \frac{\partial}{\partial z}, \quad X_{n+i} = \frac{\partial}{\partial y^i},$$

$i = 1, \dots, n$.

3.2.2 $\mathbb{R}^{n+1} \times P\mathbb{R}^n$

We now give an example of a contact manifold in the wider sense which is not a contact manifold in our sense (J. Gray [1959]). Consider \mathbb{R}^{n+1} with coordinates (x^1, \dots, x^{n+1}) and real projective space $P\mathbb{R}^n$ with homogeneous coordinates, (t_1, \dots, t_{n+1}) and let $M^{2n+1} = \mathbb{R}^{n+1} \times P\mathbb{R}^n$. The subsets $\mathcal{U}_i, i = 1, \dots, n+1$, defined by $t_i \neq 0$ form an open cover of M^{2n+1} by coordinate neighborhoods. On \mathcal{U}_i define a 1-form η_i by $\eta_i = \frac{1}{t_i} \sum_{j=1}^{n+1} t_j dx^j$; we then have $\eta_i \wedge (d\eta_i)^n \neq 0$ and $\eta_i = \frac{t_j}{t_i} \eta_j$. Thus, M^{2n+1} has a contact structure in the wider sense, but for n even, M^{2n+1} is nonorientable and hence cannot carry a global contact form.

3.2.3 $M^{2n+1} \subset \mathbb{R}^{2n+2}$ with $T_m M^{2n+1} \cap \{0\} = \emptyset$

Turning to more standard examples, we prove the following theorem (J. Gray [1959]).

Theorem 3.6 *Let $\iota : M^{2n+1} \longrightarrow \mathbb{R}^{2n+2}$ be a smooth hypersurface immersed in \mathbb{R}^{2n+2} and suppose that no tangent space of M^{2n+1} contains the origin of \mathbb{R}^{2n+2} . Then M^{2n+1} has a contact structure.*

Proof. Let $x^A, A = 1, \dots, 2n+2$, be Cartesian coordinates on \mathbb{R}^{2n+2} and consider the 1-form

$$\alpha = x^1 dx^2 - x^2 dx^1 + \dots + x^{2n+1} dx^{2n+2} - x^{2n+2} dx^{2n+1}.$$

Let V_1, \dots, V_{2n+1} be $2n+1$ linearly independent vectors at a point $x_0 = (x_0^1, \dots, x_0^{2n+2})$ and define a vector W at x_0 with components

$$W^A = *dx^A(V_1, \dots, V_{2n+1}),$$

where $*$ is the Hodge star operator of the Euclidean metric on \mathbb{R}^{2n+2} . Then W is normal to the hyperplane spanned by V_1, \dots, V_{2n+1} . Now regard x_0 as a vector with components x_0^A . Then

$$(\alpha \wedge (d\alpha)^n)(V_1, \dots, V_{2n+1}) = \sum x_0^A W^A.$$

Thus if no tangent space of M^{2n+1} regarded as a hyperplane in \mathbb{R}^{2n+2} contains the origin, then $\eta = \iota^* \alpha$ is a contact form on M^{2n+1} . \blacksquare

As a special case we see that an odd-dimensional sphere S^{2n+1} carries a contact structure. Moreover, α on S^{2n+1} is invariant under reflection through the origin, $(x^1, \dots, x^{2n+2}) \rightarrow (-x^1, \dots, -x^{2n+2})$, and hence

the real projective space $P\mathbb{R}^{2n+1}$ is also a contact manifold. J. A. Wolf [1968] then considered more general quotients of S^{2n+1} and proved that a complete connected odd-dimensional Riemannian manifold of positive constant curvature inherits a contact structure from the form α .

Similarly, consider the 1-form $\beta = \sum_{i=1}^{n+1} x^i dx^{n+1+i}$ and denote by \mathbb{R}_1^{n+1} and \mathbb{R}_2^{n+1} the subspaces defined by $x^i = 0$ and $x^{n+1+i} = 0$ respectively, $i = 1, \dots, n+1$. Then β induces a contact form on M^{2n+1} if and only if $M^{2n+1} \cap \mathbb{R}_1^{n+1} = \emptyset$ and $M^{2n+1} \cap \mathbb{R}_2^{n+1}$ is an n -dimensional submanifold and no tangent space of $M^{2n+1} \cap \mathbb{R}_2^{n+1}$ in \mathbb{R}_2^{n+1} contains the origin of \mathbb{R}_2^{n+1} .

More generally, given a symplectic manifold (M, Ω) , a hypersurface $\iota : S \rightarrow M$ is said to be of *contact type* if there exists a contact form η on S such that $d\eta = \iota^*\Omega$.

3.2.4 Unit and projectivized tangent and cotangent bundles

We shall show that the cotangent sphere bundle and the tangent sphere bundle of a Riemannian manifold are contact manifolds (see, e.g., Reeb [1952], Sasaki [1962]). Let M be an $(n+1)$ -dimensional Riemannian manifold and T^*M its cotangent bundle. Also let (x^1, \dots, x^{n+1}) be local coordinates on a neighborhood \mathcal{U} of M and $(\hat{p}^1, \dots, \hat{p}^{n+1})$ coordinates on the fibers over \mathcal{U} . If $\pi : T^*M \rightarrow M$ is the projection map, then as in Chapter 1, $q^i = x^i \circ \pi$ and \hat{p}^i are local coordinates on T^*M . Consider the Liouville form β ; locally it is given by $\beta = \sum_{i=1}^{n+1} \hat{p}^i dq^i$. The bundle T_1^*M of unit cotangent vectors has empty intersection with the zero section of T^*M , its intersection with any fiber of T^*M is an n -dimensional sphere, and no tangent space to this intersection contains the origin of the fiber. Thus, as in the discussion at the end of the last example, β induces a contact structure on the hypersurface T_1^*M of T^*M .

Instead of the unit cotangent bundle one can consider the projectivized cotangent bundle. Each fiber of T^*M is \mathbb{R}^{n+1} , and one can form the corresponding projective space $\mathbb{R}P^n$ giving the projectivized cotangent bundle, PT^*M . The \hat{p}^i 's are homogeneous coordinates for the fibers of PT^*M and we introduce nonhomogeneous coordinates (p^1, \dots, p^n) on the neighborhood defined by $\hat{p}^{n+1} \neq 0$ by $p^i = \frac{\hat{p}^i}{\hat{p}^{n+1}}$. Then

$$\eta = dq^{n+1} + \sum_{i=1}^n p^i dq^i$$

is a local contact form, and taking charts on PT^*M defined by $\hat{p}^i \neq 0$, we obtain a contact structure in the wider sense.

Example 3.2.2 can be thought of as the projectivized tangent bundle of \mathbb{R}^{n+1} , and in Section 9.5 we will briefly consider the geometry of the projectivized tangent bundle of the Beltrami model of the hyperbolic plane. In the complex setting in Chapter 13 we will consider the projectivized holomorphic tangent bundle of a Hermitian manifold.

Similarly one obtains a contact structure on the bundle T_1M of unit tangent vectors. In fact, if G_{ij} denotes the components of the metric on M with respect to the coordinates (x^1, \dots, x^{n+1}) and if (v^1, \dots, v^{n+1}) are the fiber coordinates on TM , define β locally by $\beta = \sum_{i,j} G_{ij} v^j dq^i$, where $q^i = x^i \circ \pi$ and $\pi : TM \rightarrow M$ is the projection. This structure will be discussed in detail in Section 9.2.

3.2.5 $T^*M \times \mathbb{R}$

Let M be an n -dimensional manifold and T^*M its cotangent bundle. As in the previous example we can define a 1-form β by the local expression $\beta = \sum_{i=1}^n p^i dq^i$. Let $M^{2n+1} = T^*M \times \mathbb{R}$, t the coordinate on \mathbb{R} , and $\mu : M^{2n+1} \rightarrow T^*M$ the projection to the first factor. Then $\eta = dt - \mu^*\beta$ is a contact form on M^{2n+1} .

3.2.6 *Tori*

We have mentioned that Martinet proved that every compact orientable 3-manifold carries a contact structure. Here we first give explicitly a contact structure on the 3-dimensional torus T^3 . Consider \mathbb{R}^3 with the contact form

$$\eta = \sin y \, dx + \cos y \, dz; \quad \eta \wedge d\eta = -dx \wedge dy \wedge dz,$$

$\xi = \sin y \frac{\partial}{\partial x} + \cos y \frac{\partial}{\partial z}$ and \mathcal{D} is spanned by $\{\frac{\partial}{\partial y}, \cos y \frac{\partial}{\partial x} - \sin y \frac{\partial}{\partial z}\}$. The rotation of \mathcal{D} in the direction of the y -axis is dramatically clear in this example. Thus one sees the nonintegrability of \mathcal{D} as one moves around on the manifold, especially along the y -axis.

Now η is invariant under translation by 2π in each coordinate, and hence the 3-dimensional torus also carries this structure. For each value of y , ξ induces a flow on the 2-torus defined by $y = \text{const.}$ Depending on the value of y , the flow is a rational or irrational flow on T^2 . Thus the

contact structure on T^3 is not regular. Note here in particular, though, that some of the integral curves of ξ are closed and some are not.

Concerning contact structures on higher dimensional tori, we will discuss briefly the fact that all odd-dimensional tori carry contact structures. However, we will prove in Theorem 4.14 that no torus carries a regular contact structure.

In [1979] R. Lutz proved the existence of contact structures on principal T^2 -bundles over 3-manifolds. In particular, the 5-dimensional torus admits a contact structure given by the form

$$\begin{aligned}\eta = & \sin \theta_2 \cos \theta_2 d\theta_1 - \sin \theta_1 \cos \theta_1 d\theta_2 + \cos \theta_1 \cos \theta_2 d\theta_3 \\ & + (\sin \theta_1 \cos \theta_3 - \sin \theta_2 \sin \theta_3) d\theta_4 + (\sin \theta_1 \sin \theta_3 + \sin \theta_2 \cos \theta_3) d\theta_5.\end{aligned}$$

The characteristic vector field for this structure is

$$\begin{aligned}\xi = & \frac{1}{(\sin^2 \theta_1 + \sin^2 \theta_2)^2 + \cos^2 \theta_1 \cos^2 \theta_2} \\ & \times \left(\sin \theta_2 \cos \theta_2 \frac{\partial}{\partial \theta_1} - \sin \theta_1 \cos \theta_1 \frac{\partial}{\partial \theta_2} + \cos \theta_1 \cos \theta_2 \frac{\partial}{\partial \theta_3} \right. \\ & + (\sin \theta_1 \cos \theta_3 (\sin^2 \theta_1 + \sin^2 \theta_2 - \cos^2 \theta_1) \\ & - \sin \theta_2 \sin \theta_3 (\sin^2 \theta_1 + \sin^2 \theta_2 - \cos^2 \theta_2)) \frac{\partial}{\partial \theta_4} \\ & + (\sin \theta_1 \sin \theta_3 (\sin^2 \theta_1 + \sin^2 \theta_2 - \cos^2 \theta_1) \\ & \left. + \sin \theta_2 \cos \theta_3 (\sin^2 \theta_1 + \sin^2 \theta_2 - \cos^2 \theta_2)) \frac{\partial}{\partial \theta_5} \right)\end{aligned}$$

and the contact subbundle is spanned by

$$\begin{aligned}\left\{ \frac{\partial}{\partial \theta_1} + \cos \theta_2 \sin \theta_3 \frac{\partial}{\partial \theta_4} - \cos \theta_2 \cos \theta_3 \frac{\partial}{\partial \theta_5}, \right. \\ \frac{\partial}{\partial \theta_2} + \cos \theta_1 \cos \theta_3 \frac{\partial}{\partial \theta_4} + \cos \theta_1 \sin \theta_3 \frac{\partial}{\partial \theta_5}, \\ \left. - \cos \theta_1 \frac{\partial}{\partial \theta_1} + \sin \theta_2 \frac{\partial}{\partial \theta_3}, \cos \theta_2 \frac{\partial}{\partial \theta_2} + \sin \theta_1 \frac{\partial}{\partial \theta_3} \right\}.\end{aligned}$$

Hadjar [1998] obtains other explicit contact forms on the 5-torus, especially ones for which the contact subbundle is transverse to the trivial fibration of T^5 over T^4 by circles, answering a question of Eliashberg and Thurston, [1998, p. 20].

There has been also been recent interest in constructing contact manifolds using other surfaces as fiber. In dimension 5 both Geiges [1997a] and Altschuler and Wu [2000] have proved that if M is a compact orientable 3-manifold and Σ a compact orientable surface, then $M \times \Sigma$ carries a contact form.

Altschuler and Wu also show the existence of contact forms on the products $S^{2p+k+3} \times M^k \times \Sigma$ where M^k is compact, orientable and parallelizable, and Σ is a compact orientable surface as before.

Finally, Bourgeois [2002] proved the following theorem.

Theorem 3.7 *Let M^{2n-3} be a closed contact manifold of dimension $2n - 3 \geq 3$ and let Σ_g be a Riemann surface of genus $g \geq 1$. Then the manifold $\Sigma_g \times M^{2n-3}$ also admits a contact structure.*

As a consequence we see that all odd-dimensional tori have contact structures.

3.2.7 Overtwisted contact structures

Examples 3.2.1 and 3.2.6 on \mathbb{R}^3 , namely $\eta = (dz - ydx)$ and $\eta = (\sin y dx + \cos y dz)$, have cylindrical coordinate versions (see also Douady [1982/83, p. 131], Bennequin [1983, p. 93]). Let (r, θ, z) be the usual cylindrical coordinates on $\mathbb{R}^3 \setminus \{x = y = 0\}$. Making the naive substitutions $y \rightarrow r$, $dx \rightarrow r d\theta$, $dz \rightarrow dz$, these examples become

$$\begin{aligned} \eta &= dz - r^2 d\theta, \quad \eta \wedge d\eta = -2r dr \wedge d\theta \wedge dz, \\ \xi &= \frac{\partial}{\partial z}, \quad \mathcal{D} = \left\{ \frac{\partial}{\partial r}, \frac{\partial}{\partial \theta} + r^2 \frac{\partial}{\partial z} \right\}, \end{aligned}$$

and

$$\begin{aligned} \eta &= \cos r dz + r \sin r d\theta, \quad \eta \wedge d\eta = (r + \sin r \cos r) dr \wedge d\theta \wedge dz, \\ \xi &= \frac{\sin r}{r + \sin r \cos r} \frac{\partial}{\partial \theta} + \frac{r \cos r + \sin r}{r + \sin r \cos r} \frac{\partial}{\partial z}, \\ \mathcal{D} &= \left\{ \frac{\partial}{\partial r}, \cos r \frac{\partial}{\partial \theta} - r \sin r \frac{\partial}{\partial z} \right\}. \end{aligned}$$

Note that in both examples the integral curves of $\frac{\partial}{\partial r}$ are Legendre curves and in the second example that the curve $r = \pi, z = \text{const.}$ is also a Legendre curve. Thus in the second example \mathcal{D} is tangent to the

disk $\Delta = \{z = 0, r \leq \pi\} \subset \mathbb{R}^3$ along the boundary. Now consider the topological disk $\Delta^\epsilon = \{z = \epsilon r^2, r \leq \pi\}$. \mathcal{D} is tangent to Δ^ϵ only at the origin. On $\Delta^\epsilon \setminus \{(0, 0, 0)\}$ define a line field by the intersection of the tangent plane to the paraboloid Δ^ϵ and \mathcal{D} at each point. These fields can be expressed by the vector fields

$$r \frac{\partial}{\partial r} + 2\epsilon \frac{\partial}{\partial \theta} + 2\epsilon r^2 \frac{\partial}{\partial z}$$

in the first case and

$$r \sin r \frac{\partial}{\partial r} - 2\epsilon r \cos r \frac{\partial}{\partial \theta} + 2\epsilon r^2 \sin r \frac{\partial}{\partial z}$$

in the second. For simplicity take the projection of these vector fields to the xy -plane. The integral curves in the first case are the logarithmic spirals $r = Ae^{\frac{\theta}{2\epsilon}}$ and in the second case are the curves $\theta = -2\epsilon \ln \sin r + C$, which near the origin spiral indefinitely and approach a limit cycle on the boundary of the disk. Thus we have the following diagrams; a diagram of this type for the second example was introduced by Douady [1982/83] (see also Bennequin [1983, p. 94]), and such a diagram is called the *Douady portrait* of the contact structure.



A 3-dimensional contact manifold is said to be *overtwisted* (Eliashberg [1989]) if there exists a contact embedding of a neighborhood of the disk Δ with the contact structure $\eta = \cos r \, dz + r \sin r \, d\theta$.

Roughly speaking, the meaning of overtwisted is that as one moves radially from the image of the origin, the plane \mathcal{D} turns over in a finite distance. For the radial Legendre curves in the disk Δ , \mathcal{D} turns over as

one goes from $r = 0$ to $r = \pi$. This means that for some r between 0 and π the vector field ξ is tangent to the disk Δ , and in particular for r the solution of $r \cos r + \sin r = 0$ in $(0, \pi)$, $r \approx 2.02876$, the circle of this radius in Δ is an integral curve of ξ .

As a higher dimensional analogue of being overtwisted, Niederkrüger [2006] introduced the notion of a Plastikstufe (see also Albers and Hofer [2009]). Here let Δ denote the unit disk with coordinates (x, y) . A contact manifold of dimension $2n + 1$ with contact subbundle \mathcal{D} contains a *Plastikstufe with singular set* S if it admits a closed submanifold S of dimension $n - 1$ and an embedding $\iota : \Delta \times S \longrightarrow M$ with $\iota(\{0\} \times S) = S$ having the following properties:

- (1) There exists a contact form η_{PS} inducing \mathcal{D} such that the 1-form $\beta = \iota^* \eta_{\text{PS}}$ satisfies $\beta \wedge d\beta = 0$ and $\beta \neq 0$ on $(\Delta \setminus \{0\}) \times S$. Near $\{0\} \times S$, $\beta = xdy - ydx$ and the pullback of β to $\partial\Delta \times S$ vanishes.
- (2) The complement of $\{0\} \times S$ in $(\Delta \setminus \partial\Delta) \times S$ is smoothly foliated by β via an S^1 -family of leaves diffeomorphic to $(0, 1) \times S$, where one of the ends converges to the singular set $\{0\} \times S$ and the other is asymptotic to the leaf $\partial\Delta \times S$. The set $\iota(\Delta \times S)$ is called the *Plastikstufe*, and a closed contact manifold is said to be *PS-overtwisted* if it admits a contact form η_{PS} inducing \mathcal{D} and containing a Plastikstufe.

3.2.8 $S^2 \times S^1$

Let Θ and Ψ be the azimuth and zenith on S^2 respectively ($0 \leq \Theta < 2\pi$, $0 \leq \Psi \leq \pi$) and α the coordinate on S^1 ($0 \leq \alpha < 2\pi$). Then the form

$$\eta = \frac{1}{2}(\sin \Psi d\Theta + \cos \Psi d\alpha)$$

is a contact form on $S^2 \times S^1$. The characteristic vector field and contact subbundle are given by

$$\xi = \frac{1}{2} \left(\sin \Psi \frac{\partial}{\partial \Theta} + \cos \Psi \frac{\partial}{\partial \alpha} \right), \quad \mathcal{D} = \left\{ \frac{\partial}{\partial \Psi}, \cos \Psi \frac{\partial}{\partial \Theta} - \sin \Psi \frac{\partial}{\partial \alpha} \right\}.$$

Viewing $S^2 \times S^1$ as a thick spherical shell with the interior and exterior surfaces identified, one can visualize the geometry of this structure. At the north and south poles of the 2-spheres, one sees ξ pointing upward, and at the equatorial level ($\Psi = \frac{\pi}{2}$), ξ is tangent to the equators.

An interesting variation of this structure arises if we consider the form

$$\eta = \frac{1}{2}(\sin 2\Psi d\Theta + \cos 2\Psi d\alpha).$$

Introduce new local coordinates (r, θ, z) by

$$r = 2\Psi, \quad \theta = \frac{\Theta}{2\Psi + \sin 2\Psi \cos 2\Psi}, \quad z = \alpha + \frac{\Theta \sin^2 \Psi}{2\Psi + \sin 2\Psi \cos 2\Psi}.$$

In these coordinates $\eta = r \sin r d\theta + \cos r dz$, and we see that as one moves radially, i.e., in the direction Ψ , from a north pole, the contact subbundle turns over as we reach the equator $r = \pi$ ($\Psi = \frac{\pi}{2}$). Thus we have an embedded overtwisted disk and $S^2 \times S^1$ becomes an overtwisted contact manifold.

3.2.9 Contact circles

At the beginning of this section we mentioned the result of Gonzalo [1987] that a compact orientable 3-manifold has three independent contact structures. Geiges and Gonzalo [1995] introduce the notion of a contact circle: A 3-manifold admits a *contact circle* if it admits a pair of contact forms (η_1, η_2) such that for any $(\lambda_1, \lambda_2) \in S^1$, $\lambda_1 \eta_1 + \lambda_2 \eta_2$ is also a contact form. This circle is a *taut contact circle* if the contact forms $\lambda_1 \eta_1 + \lambda_2 \eta_2$ define the same volume form for all $(\lambda_1, \lambda_2) \in S^1$; this is equivalent to the following two conditions:

$$\eta_1 \wedge d\eta_1 = \eta_2 \wedge d\eta_2, \quad \eta_1 \wedge d\eta_2 = -\eta_2 \wedge d\eta_1.$$

Geiges and Gonzalo then prove the following classification theorem.

Theorem 3.8 *A compact orientable 3-manifold admits a taut contact circle if and only if it is diffeomorphic to the quotient of a Lie group G by a discrete subgroup, acting by left multiplication, where G is either $SU(2)$, the universal cover of $PSL(2, \mathbb{R})$, or the universal cover of the group of Euclidean motions $E(2)$.*

In contrast to the overtwisted contact structures in Example 3.2.7 (and in contrast to taut contact circles), a contact structure is said to be *tight* if it is not overtwisted (see, e.g., Eliashberg and Thurston [1998]). Geiges and Gonzalo [1995] also prove that the connected sum of any number

of copies of the manifolds listed in Theorem 3.8, T^2 -bundles over S^1 , or $S^2 \times S^1$, admits a contact circle consisting of tight contact structures.

In further work Geiges and Gonzalo [1997] show that in fact on every closed, orientable 3-manifold there are contact circles realizing any of the two orientations. This paper also contains a number of explicit examples of contact circles.

3.3 The Boothby–Wang fibration

We now give an important class of examples, namely principal circle bundles over symplectic manifolds of integral class, and we will prove the celebrated theorem of Boothby and Wang [1958] that a compact regular contact manifold is of this type. An example of this type is often referred to as a *Boothby–Wang fibration*.

In Chapter 2 we saw that the set of principal circle bundles over a manifold M has a group structure isomorphic to the cohomology group $H^2(M, \mathbb{Z})$. Now let (M^{2n}, Ω) be a symplectic manifold such that $[\Omega] \in H^2(M^{2n}, \mathbb{Z})$, and $\pi : M^{2n+1} \rightarrow M^{2n}$ the corresponding circle bundle. By Theorem 2.5 there exists a connection form η on M^{2n+1} such that $d\eta = \pi^*\Omega$. Now if ξ is a vertical vector field, say with $\eta(\xi) = 1$, and X_1, \dots, X_{2n} linearly independent horizontal vector fields, then $(\eta \wedge (d\eta)^n)(\xi, X_1, \dots, X_{2n})$ is nonzero. Thus regarding the Lie-algebra-valued form η as a real-valued form, we see that η is a contact structure on M^{2n+1} .

The most well known special case of a Boothby–Wang fibration is the Hopf fibration of an odd-dimensional unit sphere S^{2n+1} over complex projective space $\mathbb{C}P^n$ of constant holomorphic curvature equal to 4. The standard contact structure on S^{2n+1} obtained in Example 3.2.3 can also be obtained by the above construction. Additional details of the geometry of Boothby–Wang fibrations will be given from time to time, particularly in Examples 4.5.4 and 6.7.2.

Theorem 3.9 *Let (M^{2n+1}, η') be a compact regular contact manifold. Then there exists a contact form $\eta = \tau\eta'$ for some nonvanishing function τ whose characteristic vector field ξ generates a free effective S^1 action on M^{2n+1} . Moreover, M^{2n+1} is the bundle space of a principal circle bundle $\pi : M^{2n+1} \rightarrow M^{2n}$ over a symplectic manifold M^{2n} whose symplectic*

form Ω determines an integral cocycle on M^{2n} and η is a connection form on the bundle with curvature form $d\eta = \pi^*\Omega$.

Proof. Since η' is regular, its characteristic vector field ξ' is a regular vector field, and hence its maximal integral curves or orbits are closed subsets of M^{2n+1} ; but M^{2n+1} is compact, so these integral curves are homeomorphic to circles. Moreover, since ξ' is regular, M^{2n+1} is a fiber bundle over a manifold M^{2n} (the set of maximal integral curves with the quotient topology; see, e.g., Palais [1957]) and we denote the projection by π .

Now let $f'_t : M^{2n+1} \rightarrow M^{2n+1}$ denote the 1-parameter group of diffeomorphisms generated by ξ' and define the period λ' of ξ' at $m \in M^{2n+1}$ by $\lambda'(m) = \inf\{t \mid t > 0, f'_t(m) = m\}$. Then λ' is constant on each orbit and since there are no fixed points, λ' is never zero. Also, since the orbits are circles, λ' is not infinite. We will show that λ' is constant on all of M^{2n+1} . Our argument is due to Tanno [1965]. Let k be a Riemannian metric on M^{2n} and let $g = \pi^*k + \eta' \otimes \eta'$. Then g is a Riemannian metric on M^{2n+1} and ξ' is a unit Killing vector field with respect to g since $\eta'(\xi') = 1$ and $\mathcal{L}_{\xi'}\eta' = 0$. If ∇ denotes the Levi-Civita connection of g , then $g(\nabla_{\xi'}\xi', X) = -g(\nabla_X\xi', \xi') = 0$ and hence the orbits of ξ' are geodesics. If γ is an orbit through m , let γ' be an orbit sufficiently near to γ that there exists a unique minimal geodesic from m to γ' meeting γ' orthogonally at m' . Then since f'_t is an isometry for all t , the image of the geodesic arc $\widehat{mm'}$ is orthogonal to γ and γ' for all t . Thus, as a point m on γ moves through one period along γ , the corresponding point on γ' moves through one period and hence λ' is constant on M^{2n+1} .

Now define η and ξ by $\eta = \frac{1}{\lambda'}\eta'$ and $\xi = \lambda'\xi'$. Since λ' is constant, ξ is the characteristic vector field of the contact form η . Moreover, ξ has the same orbits as ξ' and its period function $\lambda = 1$. Thus the one-parameter group f_t of ξ depends only on the equivalence class modulo 1 of t and the action of S^1 is effective and free.

Since ξ is regular, we may cover M^{2n+1} by coordinate neighborhoods with coordinates (x^1, \dots, x^{2n+1}) such that the integral curves of ξ are given by $x^1 = \text{const.}, \dots, x^{2n} = \text{const.}$ Projecting such neighborhoods, we obtain an open cover $\{\mathcal{U}_i\}$ of M^{2n} , and on each \mathcal{U}_i we define a local cross section s_i by setting $x^{2n+1} = \text{const.}$ Then define $F_i : \mathcal{U}_i \times S^1 \rightarrow S^1$ by $F_i(p, t) = s_i(p)t$. The transition functions for the bundle structure are then given by $f_{ij}(p) = F_i(p, t)F_j(p, t)^{-1}$.

We have already seen that $\mathcal{L}_\xi \eta = 0$ and $\mathcal{L}_\xi d\eta = 0$, so that η and $d\eta$ are invariant under the action of S^1 . Now take $A = \frac{d}{dt}$ as a basis of $\mathfrak{S}^1 = \mathbb{R}$, the Lie algebra of S^1 , and set $\tilde{\eta} = \eta A$ so that η may be regarded as a Lie-algebra-valued 1-form. For an element $B \in \mathfrak{S}^1$ denote by B^* the induced vector field on M^{2n+1} . In particular, $A^* = \xi$, so that $\tilde{\eta}(A^*) = A$. Moreover, right translation by $t \in S^1$ is just f_t so that $R_t^* \tilde{\eta} = \tilde{\eta}$ by the invariance of η under the S^1 action. Thus, η (precisely $\tilde{\eta}$) is a connection form on M^{2n+1} .

If $\tilde{\Omega}$ is the curvature form of η , then the structural equation is $d\eta = -\frac{1}{2}[\eta, \eta] + \tilde{\Omega} = \tilde{\Omega}$ since S^1 is abelian. On the other hand, $d\eta$ is horizontal and invariant, so there exists a 2-form Ω on M^{2n} such that $d\eta = \pi^* \Omega$. Now $\pi^* d\Omega = d\pi^* \Omega = d^2 \eta = 0$ so that $d\Omega = 0$ and $\pi^*(\Omega)^n = (\pi^* \Omega)^n = (d\eta)^n \neq 0$ giving $\Omega^n \neq 0$. Therefore M^{2n} is symplectic. Finally, since the transition functions f_{ij} are real (mod 1)-valued, $[\Omega] \in H^2(M^{2n}, \mathbb{Z})$ (see, e.g., Kobayashi [1956]). ■

3.4 The Weinstein conjecture

In Example 3.2.6 we studied a contact structure on the 3-dimensional torus and observed that some of the orbits of ξ are closed and some are not. It is a well-known conjecture of Weinstein [1979] that on a compact contact manifold M satisfying $H^1(M, \mathbb{R}) = 0$, ξ must have a closed orbit. The present author knows of no example of a compact contact manifold not satisfying $H^1(M, \mathbb{R}) = 0$ for which ξ does not have a closed orbit and believes the conjecture is true without the assumption of $H^1(M, \mathbb{R}) = 0$. In view of the example on the torus, the following result of Petkov and Popov [1995] is interesting. Let (M^{2n+1}, η) be an analytic, connected contact manifold with complete characteristic vector field ξ . Since $\eta \wedge (d\eta)^n$ is invariant under the action of ξ , it induces an invariant Lebesgue measure on M^{2n+1} . A point $m \in M^{2n+1}$ is a *periodic point* if the integral curve of ξ through m is periodic. Petkov and Popov prove that either the set of periodic points has measure 0 or there exists $T > 0$ such that $\exp(T\xi)(m) = m$ for every point m . Note that for some m , T could be a multiple of the period for that m (cf. the notion of an almost regular contact structure below).

Interest in the Weinstein conjecture has often been phrased in terms of the question of the existence of periodic orbits of Hamiltonian systems.

In Section 1.1 we considered briefly a real-valued function H on a symplectic manifold (M, Ω) and its Hamiltonian vector field X_H . Since $X_H H = 0$, X_H is tangent to the level (energy) surfaces of H . Thus if a level surface S is of contact type, the Hamiltonian vector field X_H is collinear with the characteristic vector field ξ .

For hypersurfaces in \mathbb{R}^{2n+2} with the standard symplectic structure, the Weinstein conjecture is known to be true if the hypersurface is convex (Weinstein [1978]), star-shaped (Rabinowitz [1978]), and, more recently, if the hypersurface is of contact type and without the assumption $H^1(M, \mathbb{R}) = 0$ (Viterbo [1987], see also Hofer and Zehnder [1987]). As an aside, it is interesting to note that Ginzburg [1995] showed that if one gives up on the hypersurface being of contact type, then there exist embeddings of S^{2n+1} , $n \geq 3$, in \mathbb{R}^{2n+2} whose Hamiltonian vector field has no closed orbits.

In cotangent bundles T^*M , first note that if S is a compact connected hypersurface, then $T^*M \setminus S$ has exactly two components, one of which is bounded; Hofer and Viterbo [1988] prove that if the bounded component of $T^*M \setminus S$ contains the zero section and if S is of contact type, then ξ has a closed orbit.

For overtwisted compact contact manifolds the Weinstein conjecture is intuitive, as alluded to in Example 3.2.7. On the standard overtwisted disk we noted that as one moves radially from the origin, the plane \mathcal{D} turns over as one goes from $r = 0$ to $r = \pi$. In particular, the vector field ξ is tangent to the disk for r the solution of $r \cos r + \sin r = 0$ in $(0, \pi)$, giving a periodic orbit of ξ . In [1993] Hofer showed that indeed the Weinstein conjecture is true on a compact overtwisted contact manifold. In the same paper he proved the conjecture for any contact form on the 3-sphere and on a closed orientable 3-manifold with $\pi_2 \neq 0$.

In [2005], Abbas, Cieliebak and Hofer proved the 3-dimensional Weinstein conjecture for planar contact structures. A contact structure is said to be *planar* if the pages of a supporting open-book decomposition have genus zero. These include overtwisted contact structures.

Finally, for the 3-dimensional case, C. H. Taubes [2007] proved the Weinstein conjecture in full generality. His proof uses a perturbed version of the Seiberg–Witten equations on 3-dimensional manifolds.

In higher dimension Albers and Hofer [2009] proved the Weinstein conjecture for closed PS-overtwisted contact manifolds, showing that every

Reeb vector field associated to a contact form inducing the contact subbundle \mathcal{D} has a contractible periodic orbit.

Banyaga [1990] showed that the Weinstein conjecture is true if the contact form is C^2 -close to a regular contact form (again without assuming $H^1(M, \mathbb{R}) = 0$). C. B. Thomas [1976] introduced the notion of an almost regular contact structure. A contact structure is said to be *almost regular*, sometimes called *quasi-regular*, if there exists a positive integer N such that every point has a neighborhood such that any integral curve of ξ passing through the neighborhood passes through at most N times. With this idea in mind, Banyaga and Rukimbira [1994] showed that the Weinstein conjecture is true if the contact form is C^1 -close to an almost regular contact form.

A contact manifold is called an *R-contact manifold* (Rukimbira [1993]) if ξ is Killing with respect to some (not necessarily associated (Chapter 4)) Riemannian metric g for which $\eta(X) = g(X, \xi)$. For compact contact manifolds admitting such a metric the Weinstein conjecture is true (Rukimbira [1993]). Also on a compact K-contact manifold M^{2n+1} (see Subsection 4.5.4 or Section 6.2), ξ has at least $n+1$ closed orbits (Rukimbira [1995a]). If a compact K-contact manifold M^{2n+1} has exactly $n+1$ closed orbits, then it is finitely covered by a sphere (Rukimbira [2000]).

4

Associated Metrics

The main topic of this chapter is that of Riemannian metrics associated to symplectic and contact structures, and their construction by means of polarization. We also discuss the action of symplectic and contact transformations on associated metrics. Some of our discussion is broader, dealing with almost Hermitian and almost contact metric structures. The chapter closes with several examples.

4.1 Almost complex and almost contact structures

We will generally regard the theory of almost Hermitian structures as well known and give here only definitions and a few properties that will be important for our study; many of these were already mentioned in Chapter 1. For more detail the reader is referred to Gray and Hervella [1980], Kobayashi–Nomizu [1963–69, Chapter IX] and Kobayashi–Wu [1983]; also, despite its classical nature, the book of Yano [1965] contains helpful information on many of these structures.

An *almost complex structure* is a tensor field J of type $(1, 1)$ such that $J^2 = -I$. A *Hermitian metric* on an almost complex manifold (M, J) is a Riemannian metric that is invariant by J , i.e.,

$$g(JX, JY) = g(X, Y).$$

Noting that J is negative self-adjoint with respect to g , i.e., $g(X, JY) = -g(JX, Y)$, therefore $\Omega(X, Y) = g(X, JY)$ defines a 2-form called the *fundamental 2-form* of the *almost Hermitian structure* (M, J, g) . If $d\Omega = 0$, the structure is *almost Kähler*. If M is a complex manifold and J the corresponding almost complex structure, we say that (M, J, g) is a *Hermitian manifold*. For geometers working strictly over the complex domain, a Hermitian metric is a Hermitian quadratic form and hence complex-valued. It takes its nonzero values, as appropriate, when one argument is of type $(1, 0)$ and the other of type $(0, 1)$. In particular, our metric g becomes half the real part of $g(X - iJX, Y + iJY)$. For geometers concerned with a variety of structures presented for the most part in terms of real tensor fields, as we will be, it has become quite standard to use the word Hermitian as we have done. A reader interested in this point may want to see Kobayashi–Wu [1983, pp. 80–81] for commentary.

Note also that every almost complex manifold admits a Hermitian metric, for if k is any Riemannian metric, then g defined by

$$g(X, Y) = k(X, Y) + k(JX, JY)$$

is Hermitian.

Given (M, J, g) we can construct a particular local orthonormal basis as follows. Let \mathcal{U} be a coordinate neighborhood on M and X_1 any unit vector field on \mathcal{U} . Let $X_{1*} = JX_1$. Now choose a unit vector field X_2 orthogonal to both X_1 and X_{1*} . Then JX_2 is also orthogonal to X_1 and X_{1*} . Continuing in this manner we have a local orthonormal basis of the form $\{X_1, \dots, X_n, JX_1, \dots, JX_n\}$. Such a basis is called a *J-basis*. Note in particular that an almost complex manifold is even-dimensional.

Again given (M, J) , choose g Hermitian. Let $\{\mathcal{U}_\alpha\}$ be an open cover with J -bases $\{X_i, X_{i*}\}$, $\{\bar{X}_i, \bar{X}_{i*}\}$ on \mathcal{U}_α and \mathcal{U}_β respectively. With respect to these bases J is given by

$$\begin{pmatrix} 0 & -I \\ I & 0 \end{pmatrix}.$$

If now $X \in T_m M$, $m \in \mathcal{U}_\alpha \cap \mathcal{U}_\beta$, then for the column vectors of components (X) and (\bar{X}) with respect to these bases, we have

$$(\bar{X}) = \begin{pmatrix} A & B \\ C & D \end{pmatrix} (X),$$

where A, B, C, D are $n \times n$ matrices and $\begin{pmatrix} A & B \\ C & D \end{pmatrix} \in O(2n)$. Now

$$\begin{aligned} \begin{pmatrix} 0 & -I \\ I & 0 \end{pmatrix} \begin{pmatrix} A & B \\ C & D \end{pmatrix} (X) &= (\overline{JX}) = \begin{pmatrix} A & B \\ C & D \end{pmatrix} (JX) \\ &= \begin{pmatrix} A & B \\ C & D \end{pmatrix} \begin{pmatrix} 0 & -I \\ I & 0 \end{pmatrix} (X), \end{aligned}$$

i.e. $\begin{pmatrix} A & B \\ C & D \end{pmatrix}$ and $\begin{pmatrix} 0 & -I \\ I & 0 \end{pmatrix}$ commute. Therefore $D = A$ and $C = -B$ and hence $\begin{pmatrix} A & B \\ C & D \end{pmatrix} \in U(n)$. In particular, the structural group of the tangent bundle of an almost complex manifold is reducible to $U(n)$. Recall also that $\det \begin{pmatrix} A & B \\ -B & A \end{pmatrix} = |\det(A + iB)|^2 > 0$ and therefore an almost complex manifold is orientable.

Conversely, suppose that we are given M such that the structural group of TM can be reduced to $U(n)$. Let $\{\mathcal{U}_\alpha\}$ be an open cover such that we can choose local orthonormal bases which transform in the overlaps of neighborhoods by the action of $U(n)$. In each $\{\mathcal{U}_\alpha\}$ define J_α by $\begin{pmatrix} 0 & -I \\ I & 0 \end{pmatrix}$; this matrix commutes with $U(n)$, and hence the set $\{J_\alpha\}$ determines a global tensor field J such that $J^2 = -I$. Thus an almost complex (almost Hermitian) structure on M can be thought of as a reduction of the structural group to $U(n)$.

As we will see in our discussion of associated metrics below, the structural group of a symplectic manifold is reducible to $U(n)$ and that of a contact manifold to $U(n) \times 1$ (Chern [1953]). For an odd-dimensional manifold M^{2n+1} , J. Gray [1959] defined an *almost contact structure* as a reduction of the structural group to $U(n) \times 1$. In terms of structure tensors we say that M^{2n+1} has an *almost contact structure* or sometimes (ϕ, ξ, η) -*structure* if it admits a tensor field ϕ of type $(1, 1)$, a vector field ξ , and a 1-form η satisfying

$$\phi^2 = -I + \eta \otimes \xi, \quad \eta(\xi) = 1.$$

Many authors include also that $\phi\xi = 0$ and $\eta \circ \phi = 0$. However, these are deducible from the other conditions, as we now show.

Theorem 4.1 *Suppose M^{2n+1} has a (ϕ, ξ, η) -structure. Then $\phi\xi = 0$ and $\eta \circ \phi = 0$. Moreover, the endomorphism ϕ has rank $2n$.*

Proof. First note that $\phi^2 = -I + \eta \otimes \xi$ and $\eta(\xi) = 1$ give $\phi^2\xi = -\xi + \eta(\xi)\xi = 0$ and hence either $\phi\xi = 0$ or $\phi\xi$ is a nontrivial eigenvector of ϕ corresponding to the eigenvalue 0. Using $\phi^2 = -I + \eta \otimes \xi$ again,

we have $0 = \phi^2 \phi \xi = -\phi \xi + \eta(\phi \xi) \xi$ or $\phi \xi = \eta(\phi \xi) \xi$. Now if $\phi \xi$ is a nontrivial eigenvector of the eigenvalue 0, $\eta(\phi \xi) \neq 0$, and therefore $0 = \phi^2 \xi = \eta(\phi \xi) \phi \xi = (\eta(\phi \xi))^2 \xi \neq 0$, a contradiction. Thus, $\phi \xi = 0$.

Now since $\phi \xi = 0$, we also have that $\eta(\phi X) \xi = \phi^3 X + \phi X = -\phi X + \phi(\eta(X) \xi) + \phi X = 0$ for any vector field X and hence $\eta \circ \phi = 0$.

Finally, since $\phi \xi = 0$, $\xi \neq 0$ everywhere, $\text{rank } \phi < 2n + 1$. If a vector field $\bar{\xi}$ satisfies $\phi \bar{\xi} = 0$, then $\phi^2 = -I + \eta \otimes \xi$ gives $0 = -\bar{\xi} + \eta(\bar{\xi}) \xi$; thus $\bar{\xi}$ is collinear with ξ and so $\text{rank } \phi = 2n$. ■

If a manifold M^{2n+1} with a (ϕ, ξ, η) -structure admits a Riemannian metric g such that

$$g(\phi X, \phi Y) = g(X, Y) - \eta(X)\eta(Y),$$

we say that M^{2n+1} has an *almost contact metric structure* and g is called a *compatible* metric. Setting $Y = \xi$ we have immediately that

$$\eta(X) = g(X, \xi).$$

As in the almost Hermitian case, we define a 2-form, called the *fundamental 2-form* of the almost contact metric structure, by $\Phi(X, Y) = g(X, \phi Y)$. Also as in the case of an almost complex structure, the existence of the compatible metric is easy. For if k' is any metric, first set $k(X, Y) = k'(\phi^2 X, \phi^2 Y) + \eta(X)\eta(Y)$; then $\eta(X) = k(X, \xi)$. Now define g by

$$g(X, Y) = \frac{1}{2}(k(X, Y) + k(\phi X, \phi Y) + \eta(X)\eta(Y))$$

and check the details.

For a manifold M^{2n+1} with an almost contact metric structure (ϕ, ξ, η, g) we can also construct a useful local orthonormal basis. Let \mathcal{U} be a coordinate neighborhood on M and X_1 any unit vector field on \mathcal{U} orthogonal to ξ . Then $X_{1*} = \phi X_1$ is a unit vector field orthogonal to both X_1 and ξ . Now choose a unit vector field X_2 orthogonal to ξ , X_1 and X_{1*} . Then ϕX_2 is also a unit vector field orthogonal to ξ , X_1 , X_{1*} and X_2 . Proceeding in this way we obtain a local orthonormal basis $\{X_i, X_{i*} = \phi X_i, \xi\}$, called a *ϕ -basis*.

Now given a manifold M^{2n+1} with a (ϕ, ξ, η) -structure, let g be a compatible metric and $\{\mathcal{U}_\alpha\}$ an open cover with ϕ -bases $\{X_i, X_{i*}, \xi\}$. With respect to such a basis ϕ is given by the matrix

$$\begin{pmatrix} 0 & -I & 0 \\ I & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}.$$

Proceeding as in the almost complex case, we see that the structural group of M^{2n+1} is reducible to $U(n) \times 1$. Conversely, given an almost contact structure as defined by this reduction of the structural group and an open cover $\{\mathcal{U}_\alpha\}$ respecting the action of $U(n) \times 1$, define ϕ_α on \mathcal{U}_α by the above matrix and define η_α and ξ_α by row and column vectors of $2n$ zeros and last entry 1. Then, again as in the almost complex case, this defines global structure tensors (ϕ, ξ, η) satisfying $\phi^2 = -I + \eta \otimes \xi$ and $\eta(\xi) = 1$. We shall subsequently speak of an almost contact structure (ϕ, ξ, η) and suppress the terminology “ (ϕ, ξ, η) -structure”.

4.2 Polarization and associated metrics

We begin with a discussion of the well-known decomposition, called “polarization”, of a nonsingular matrix A into the product of an orthogonal matrix F and a positive definite symmetric matrix G . Let $H(n)$ denote the set of positive definite symmetric $n \times n$ matrices and as usual $O(n)$ the orthogonal group. In treating the subject of constructing Riemannian metrics associated to 2-forms of rank $2r$, Y. Hatakeyama [1962] proved the analyticity of the polar decomposition; that the decomposition is continuous can be found in Chevalley [1946, pp. 14–16]. We prove Hatakeyama’s result by a sequence of lemmas.

Lemma 4.1 *For $G \in H(n)$, let $\lambda_i > 0$ be the eigenvalues of G . Then the map $\sigma_G : \mathfrak{gl}(n, \mathbb{R}) \longrightarrow \mathfrak{gl}(n, \mathbb{R})$ given by $\sigma_G(A) = GAG^{-1}$ has positive eigenvalues $\frac{\lambda_i}{\lambda_l}$.*

Proof. There exists $P \in O(n)$ such that PGP^{-1} is diagonal, say Δ . Then $\sigma_P^{-1}\sigma_\Delta\sigma_P = \sigma_G$ and hence σ_Δ and σ_G have the same eigenvalues. Now $\sigma_\Delta(A)_{il} = (\Delta A \Delta^{-1})_{il} = \sum_{jk} \lambda_i \delta_{ij} a_{jk} \delta_{kl} \frac{1}{\lambda_l} = \frac{\lambda_i}{\lambda_l} a_{il}$. Thus the n^2 eigenvalues of σ_G are the positive numbers $\frac{\lambda_i}{\lambda_l}$. ■

Lemma 4.2 *For $G \in H(n)$ and A skew-symmetric, AG symmetric implies that $A = 0$.*

Proof. $AG = G^T A^T = -GA$. Therefore $\sigma_G(A) = -A$, and so by the previous lemma $A = 0$. ■

Now $O(n)$ and $H(n)$ are analytic submanifolds of $GL(n, \mathbb{R})$; thus $\varphi : O(n) \times H(n) \longrightarrow GL(n, \mathbb{R})$ defined by $\varphi(F, G) = FG$ is analytic.

Lemma 4.3 *$d\varphi$ is one-to-one and hence φ^{-1} given by polarization is analytic.*

Proof. Let $X \in T_{(F,G)}O(n) \times H(n)$ and consider the curve $(Fe^{tA}, G + tB)$, where A is skew-symmetric and B is symmetric and which has tangent X at (F, G) :

$$d\varphi(X) = \lim_{t \rightarrow 0} \frac{Fe^{tA}(G + tB) - FG}{t} = FAG + FB.$$

If now $d\varphi(X) = 0$, then $F(AG + B) = 0$. Therefore $AG = -B$, which is symmetric. Thus by Lemma 4.2, $A = 0$ and hence also $B = 0$. ■

Theorem 4.2 *Polarization as a map from $GL(n, \mathbb{R}) \longrightarrow O(n) \times H(n)$ gives an analytic diffeomorphism between these manifolds with respect to the usual analytic structures.*

We now prove the existence of associated metrics.

Theorem 4.3 *Let (M^{2n}, Ω) be a symplectic manifold. Then there exist a Riemannian metric g and an almost complex structure J such that*

$$g(X, JY) = \Omega(X, Y).$$

Proof. Let k be any Riemannian metric on M and let $\{X_1, \dots, X_{2n}\}$ be a local k -orthonormal basis. Let $A_{ij} = \Omega(X_i, X_j)$. A is a $2n \times 2n$ nonsingular skew-symmetric matrix. By polarization we have $A = FG$ for some orthogonal matrix F and positive definite symmetric matrix G . Now define g and J by

$$g(X_i, X_j) = G_{ij}, \quad JX_i = F_i^j X_j.$$

g is independent of the choice of k -orthonormal basis. For if $\{Y_1, \dots, Y_{2n}\}$ is another k -orthonormal basis, there is an orthogonal matrix P such that

$$B_{ij} = \Omega(Y_i, Y_j) = \Omega(P^k_i X_k, P^l_j X_l) = P^k_i P^l_j A_{kl} = (PAP^{-1})_{ij}.$$

If $B = \Phi\Gamma$ is the polarization of B , then $\Phi\Gamma = PFP^{-1}PGP^{-1}$, and so by the uniqueness of the polar decomposition, $\Phi = PFP^{-1}$ and $\Gamma = PGP^{-1}$. Thus, in particular, we see that g and J are globally defined. Also since A is skew-symmetric, $F^2 = -I$ and F is skew-symmetric. To see this, note that $A^T = GF^T = -FG$ and hence applying F on

the right, $G = -FFF^TGF$. But F^TGF is positive definite symmetric, and so the uniqueness of the decomposition gives $-F^2 = I$. Finally, $F = -F^{-1} = -F^T$. ■

In particular, given a symplectic manifold (M^{2n}, Ω) , we say that a Riemannian metric g is an *associated metric* if there exists an almost complex structure J such that $g(X, JY) = \Omega(X, Y)$. We remark that if g is an associated metric and one uses it as the starting metric k in the above polarization process, the process yields the metric g back again.

Theorem 4.4 *Let (M^{2n+1}, η) be a contact manifold and ξ its characteristic vector field. Then there exists an almost contact metric structure such that $g(X, \phi Y) = d\eta(X, Y)$.*

Proof. This time the proof is a two-step process. First let k' be any Riemannian metric and define a new metric k by

$$k(X, Y) = k'(-X + \eta(X)\xi, -Y + \eta(Y)\xi) + \eta(X)\eta(Y).$$

Then $k(X, \xi) = \eta(X)$. Now polarize $d\eta$ on the contact subbundle \mathcal{D} as in the symplectic case. This gives a metric g' and almost complex structure ϕ' on \mathcal{D} such that $g'(X, \phi'Y) = d\eta(X, Y)$. Extending g' to a metric g agreeing with k in the direction ξ and extending ϕ' to a field of endomorphisms ϕ by requiring $\phi\xi = 0$, we have an almost contact metric structure (ϕ, ξ, η, g) such that $g(X, \phi Y) = d\eta(X, Y)$. ■

As in the symplectic case, given a contact manifold (M^{2n+1}, η) , we say that a Riemannian metric g is an *associated metric* if there exists an almost contact metric structure such that $g(X, \phi Y) = d\eta(X, Y)$. In this case we also speak of a *contact metric structure*; other authors often use the phrase *contact Riemannian structure*. Working strictly with structure tensors, one may avoid the polarization process by defining an associated metric as follows: Given a contact manifold (M^{2n+1}, η) with characteristic vector field ξ , a Riemannian metric g is an associated metric if, first of all,

$$\eta(X) = g(X, \xi)$$

and secondly, there exists a tensor field ϕ of type $(1, 1)$ such that

$$\phi^2 = -I + \eta \otimes \xi, \quad d\eta(X, Y) = g(X, \phi Y).$$

Finally, we caution that it is possible to have a contact manifold (M^{2n+1}, η) with characteristic vector field ξ and an almost contact metric structure

(ϕ, ξ, η, g) , same ξ and η , without $g(X, \phi Y) = d\eta(X, Y)$; see Example 4.5.3 for an example.

In the course of this book we will give many properties of associated metrics; we give one simple property here, since it is discussed in Example 4.5.5 and used periodically.

Theorem 4.5 *On a contact metric manifold the integral curves of ξ are geodesics.*

Proof. For a contact metric structure we have

$$0 = (\mathcal{L}_\xi \eta)(X) = \xi g(X, \xi) - g(\nabla_\xi X - \nabla_X \xi, \xi) = g(X, \nabla_\xi \xi),$$

so the integral curves of ξ are geodesics. ■

In our discussion above in both the symplectic and contact cases we started with an arbitrary Riemannian metric and obtained an associated metric. Thus we are led to believe that there are many associated metrics for a given symplectic or contact form. Indeed, this is the case, and we now show that the set \mathcal{A} of all associated metrics is infinite-dimensional by exhibiting a path of metrics in \mathcal{A} determined by a C^∞ function with compact support. Such paths of associated metrics will be useful to us in the study of critical points of curvature functionals on \mathcal{A} . We give the construction in the symplectic case and then remark on the similarity with the contact case.

Let f be a C^∞ function with compact support contained in a neighborhood \mathcal{U} of (M^{2n}, Ω) and $\{X_1, \dots, X_n, X_{1*}, \dots, X_{n*}\}$ a local J -basis. Let g be an associated metric. Make no change in g outside \mathcal{U} , and on \mathcal{U} change g only in the planes spanned by $\{X_1, X_{1*}\}$ by the matrix

$$\begin{pmatrix} 1 + tf + \frac{1}{2}t^2 f^2 & \frac{1}{2}t^2 f^2 \\ \frac{1}{2}t^2 f^2 & 1 - tf + \frac{1}{2}t^2 f^2 \end{pmatrix}.$$

This defines a path of metrics g_t , and it is easy to check that each g_t is an associated metric for the symplectic form Ω . In the contact case simply begin with a ϕ -basis and make the same construction.

Also, as already remarked in Section 1.1, it is evident that in the symplectic case, \mathcal{A} may be thought of as the set of all almost Kähler metrics that have Ω as their fundamental 2-form.

On the other hand, it is possible for a Riemannian metric g to be an associated metric for more than one symplectic structure. For example,

on a *hyper-Kähler manifold* one has, by definition, three independent global Kähler structures (J_a, g) , $a = 1, 2, 3$, satisfying $J_1 J_2 + J_2 J_1 = 0$, $J_3 = J_1 J_2$. Thus the three fundamental 2-forms give three symplectic structures with g an associated metric for each of them.

We close this section by noting that all associated metrics have the same volume element and give the proof only in the symplectic case.

Theorem 4.6 *Let (M^{2n}, Ω) be a symplectic manifold (resp. (M^{2n+1}, η) a contact manifold) and g an associated metric. Then*

$$dV = \frac{(-1)^n}{2^n n!} \Omega^n \text{ (resp. } dV = \frac{(-1)^n}{2^n n!} \eta \wedge (d\eta)^n \text{)}.$$

Proof. Let $X_1, \dots, X_n, X_1^*, \dots, X_n^*$ be a J -basis and $\theta^1, \dots, \theta^n, \theta^{1*}, \dots, \theta^{n*}$ the dual basis. Then with respect to this dual basis,

$$dV = \theta^1 \wedge \theta^{1*} \wedge \theta^2 \wedge \theta^{2*} \wedge \dots \wedge \theta^n \wedge \theta^{n*}$$

and

$$\Omega = \sum_{i=1}^n (\theta^{i*} \wedge \theta^i - \theta^i \wedge \theta^{i*}) = -2 \sum_{i=1}^n \theta^i \wedge \theta^{i*}.$$

Therefore

$$\begin{aligned} \Omega^n &= (-2)^n ((\theta^1 \wedge \theta^{1*}) + \dots + (\theta^n \wedge \theta^{n*}))^n \\ &= (-2)^n n! (\theta^1 \wedge \theta^{1*}) \wedge \dots \wedge (\theta^n \wedge \theta^{n*}) = (-2)^n n! dV. \end{aligned}$$

■

4.3 Polarization of metrics as a projection

In the previous section we created associated metrics from an arbitrary metric by polarization. In this section we give some further properties of the set \mathcal{A} and discuss how \mathcal{A} sits in the set \mathcal{N} of all Riemannian metrics with the same volume element. Restricting ourselves to \mathcal{N} , we will interpret the polarization process of constructing associated metrics from a given metric as a projection from \mathcal{N} onto \mathcal{A} .

On a compact manifold M the set \mathcal{M} of all Riemannian metrics may be given a Riemannian metric (Ebin [1970]): The tangent space, $T_g \mathcal{M}$, of \mathcal{M} at a metric g is the space of symmetric tensor fields of type $(0, 2)$.

For symmetric tensor fields S and T of type $(0, 2)$ we define a Riemannian metric (\cdot, \cdot) by

$$(S, T)_g = \int_M S_{ij} T_{kl} g^{ik} g^{jl} dV_g.$$

For the set \mathcal{N} of metrics with the same volume element, the geodesics in \mathcal{N} were found by Ebin [1970] and are curves of the form ge^{St} , where S is symmetric with $\text{tr } S = 0$. $g_t = ge^{St}$ is computed by

$$g_t(X, Y) = g(X, e^{St}Y),$$

where here e^{St} acts on Y as a tensor field of type $(1, 1)$. Again and throughout we shall often denote a symmetric tensor field of type $(0, 2)$ and the corresponding tensor field of type $(1, 1)$ by the same letter. In fact, in most of our computations we begin with a local orthonormal basis for some $g \in \mathcal{A}$ and regard S simply as its matrix of components.

As an aside, we remark that one can think of metrics with the same total volume on an n -dimensional manifold as being at a fixed distance from the zero tensor. Consider the path tg , $t \in [0, 1]$, from the zero tensor to the metric g ; then since g may be considered as the tangent to the path at each point, we have the following entertaining computation:

$$|g| \equiv \int_0^1 (g, g)_{tg}^{1/2} dt = \int_0^1 \left(\frac{n}{t^2} \int_M \sqrt{t^n \det g} dx^1 \cdots dx^n \right)^{1/2} dt = 4\sqrt{\frac{\text{vol}_g M}{n}}.$$

4.3.1 Some linear algebra

For the projection result below, Theorem 4.8, we will need the polarization of a particular path in $GL(2n, \mathbb{R})$. Let \mathcal{J} denote the matrix $\begin{pmatrix} 0 & -I \\ I & 0 \end{pmatrix}$ and let S be any symmetric $2n \times 2n$ matrix. First diagonalize S , say $Q^{-1}SQ = \Lambda$, $Q \in O(2n)$, Λ diagonal, and set $P = Qe^{-\frac{1}{2}\Lambda t}$. Our problem will be to polarize $P^T \mathcal{J} P$, i.e., find $F(t) \in O(2n)$, $G(t) \in H(2n)$ such that $P^T \mathcal{J} P = F(t)G(t)$.

Lemma 4.4 *Any symmetric $2n \times 2n$ matrix S can be uniquely written as $B + D$, where B and D are symmetric and $\mathcal{J}B - B\mathcal{J} = 0$ and $\mathcal{J}D + D\mathcal{J} = 0$.*

Proof. Setting $B = \frac{1}{2}(-\mathcal{J}S\mathcal{J} + S)$ and $D = \frac{1}{2}(\mathcal{J}S\mathcal{J} + S)$, the decomposition is immediate. If now $B + D = B' + D'$, then $B - B' = D' - D$

and hence $\mathcal{J}(B - B') = (B - B')\mathcal{J}$ gives $\mathcal{J}(D' - D) = (D' - D)\mathcal{J} = -\mathcal{J}(D' - D)$. Therefore $\mathcal{J}(D' - D) = 0$, and so since \mathcal{J} is nonsingular, $D' = D$ and in turn $B' = B$. ■

Of course B and D need not commute with each other. We will see below that if $[B, D] = 0$, then $F(t) = Q^{-1}\mathcal{J}Q$, and $G(t) = Q^{-1}e^{-Bt}Q$, and conversely, if either $F(t)$ or $G(t)$ has this simple form, B and D commute.

As a matter of notation, using the analyticity of the polar decomposition, we write

$$F(t) = \sum_{k=0}^{\infty} F^{(k)} t^k, \quad G(t) = \sum_{k=0}^{\infty} G^{(k)} t^k.$$

At $t = 0$, $P^T \mathcal{J} P = Q^{-1} \mathcal{J} Q$, which we denote by M . Since $\mathcal{J} \in O(2n)$, we have $F(0) = M$ and $G(0) = I$. The main lemma of this section is the following; due to its complexity we refer to the author's paper [1983] for its proof.

Lemma 4.5

$$\begin{aligned} F^{(k)} &= \frac{1}{2} \sum_{j=1}^{k-1} \left(\frac{(-1)^k}{k! 2^k} \binom{k}{j} \right) Q^{-1} \mathcal{J} [(B - D)^{k-j}, (B + D)^j] Q \\ &\quad - F^{(k-j)} G^{(j)} - M G^{(j)} F^{(k-j)} M + M F^{(k-j)} F^{(j)}, \\ G^{(k)} &= M F^{(k)} + M \sum_{j=1}^{k-1} F^{(k-j)} G^{(j)} \\ &\quad + \frac{(-1)^k}{k! 2^k} Q^{-1} \left(\sum_{j=0}^k \binom{k}{j} (B - D)^{k-j} (B + D)^j \right) Q. \end{aligned}$$

From this lemma we see easily that

$$F^{(1)} = 0, \quad F^{(2)} = \frac{1}{4} Q^{-1} \mathcal{J} [B, D] Q.$$

Continuing, we can find $F^{(k)}$ and $G^{(k)}$ as far as desired; in particular,

$$\begin{aligned} G^{(1)} &= -Q^{-1} B Q, \quad G^{(2)} = \frac{1}{2} Q^{-1} B^2 Q, \\ G^{(3)} &= Q^{-1} \left(-\frac{1}{6} B^3 - \frac{1}{24} (B D^2 - 2 D B D + D^2 B) \right. \\ &\quad \left. - \frac{1}{8} (B^2 D - 2 B D B + D B^2) \right) Q, \end{aligned}$$

$$\begin{aligned}
G^{(4)} = Q^{-1} & \left(\frac{1}{24}B^4 + \frac{1}{16}(B^3D - B^2DB - BDB^2 + DB^3) \right. \\
& + \frac{1}{96}(2B^2D^2 - 7BDBD + 3DB^2D + 7BD^2B \\
& \left. - 7DBDB + 2D^2B^2) \right) Q.
\end{aligned}$$

Corollary 4.1 $[B, D] = 0$ implies $F(t) = Q^{-1}\mathcal{J}Q$ and $G(t) = Q^{-1}e^{-Bt}Q$. Conversely, either of these implies the commutativity.

Proof. $[B, D] = 0$ implies that the bracket in $F^{(k)}$ vanishes, and hence by induction, $F^{(k)} = 0$ for $k > 0$. Again if B and D commute,

$$\sum_{j=0}^k \binom{k}{j} (B - D)^{k-j} (B + D)^j = 2^k B^k$$

and hence $G(t) = Q^{-1}e^{-Bt}Q$.

Clearly $F(t) = Q^{-1}\mathcal{J}Q$ gives $F^{(2)} = \frac{1}{4}Q^{-1}\mathcal{J}[B, D]Q = 0$ and hence $[B, D] = 0$. Finally, if $G(t) = Q^{-1}e^{-Bt}Q$, then

$$QG^{(3)}Q^{-1} + \frac{1}{6}B^3 = 0, \quad QG^{(4)}Q^{-1} - \frac{1}{24}B^4 = 0.$$

Thus if we multiply the rest of the expression for $G^{(3)}$ on the left by $48B$ and separately on the right by $48B$ and add these two to 96 times the rest of $G^{(4)}$, we have

$$-3BDBD + 3DB^2D + 3BD^2B - 3DBDB = 0.$$

Thus $[B, D]^2 = 0$, but $[B, D]$ is skew-symmetric and hence $[B, D] = 0$. ■

Lemma 4.6 $\mathcal{J}e^S = e^{-S}\mathcal{J}$ if and only if $S\mathcal{J} + \mathcal{J}S = 0$.

Proof. The sufficiency is clear, so we prove only the necessity. Suppose $SX = \lambda X$. Then $\mathcal{J}e^SX = e^\lambda\mathcal{J}X = e^{-S}\mathcal{J}X$. Also let $\{e_i\}$ be an orthonormal eigenvector basis of S with $Se_i = \lambda_i e_i$. If now $\mathcal{J}X = Y^i e_i$, then

$$e^{-S}\mathcal{J}X = e^\lambda\mathcal{J}X = e^\lambda Y^i e_i$$

and

$$e^{-S}\mathcal{J}X = e^{-S}Y^i e_i = \sum_i Y^i e^{-\lambda_i} e_i.$$

Thus for each i , either $\lambda = -\lambda_i$ or $Y^i = 0$ and hence

$$S\mathcal{J}X = \sum_i Y^i \lambda_i e_i = -\lambda Y^i e_i = -\lambda \mathcal{J}X,$$

but $\mathcal{J}SX = \lambda \mathcal{J}X$, giving $(S\mathcal{J} + \mathcal{J}S)X = 0$ for any eigenvector X and hence $S\mathcal{J} + \mathcal{J}S = 0$. ■

4.3.2 Results on the set \mathcal{A}

First we will give a remark about general curves in \mathcal{A} . Let g be a metric in \mathcal{A} . If $g_t = g + D^{(1)}t + D^{(2)}t^2 + \dots$ is a path of metrics, we can easily obtain a sequence of necessary conditions for g_t to lie in \mathcal{A} . Again note that we use the convention that D will signify a tensor field of type $(0, 2)$ or type $(1, 1)$, related by the metric g , and it being clear from context which is meant. We give the details in the symplectic case; the reader can easily read ϕ instead of J for the contact case and follow the computation, showing also that $D^{(k)}\xi = 0$ by a small amount of further computation:

$$g(X, JY) = \Omega(X, Y) = g_t(X, J_t Y) = g(X, J_t Y) + \sum_{k=1}^{\infty} D^{(k)}(X, J_t Y)t^k,$$

from which

$$J = J_t + D^{(1)}J_t t + D^{(2)}J_t t^2 + \dots$$

Applying J_t on the right and J on the left, we have

$$J_t = J(I + D^{(1)}t + D^{(2)}t^2 + \dots).$$

Squaring this and comparing coefficients gives

$$JD^{(k)} + D^{(k)}J = -\sum_{j=1}^{k-1} D^{(j)}JD^{(k-j)}.$$

Using this last result repeatedly, we see that

$$JD^{(k)} + D^{(k)}J = J(\text{polynomial in the } D^{(j)}, j < k).$$

In particular, we have the following:

$$JD^{(1)} + D^{(1)}J = 0, \quad JD^{(2)} + D^{(2)}J = JD^{(1)2}.$$

These equations yield immediately the following corollary.

Corollary 4.2 *\mathcal{A} contains no line of metrics.*

Recall that in the contact case the construction of associated metrics was a two-step process, the first being the creation of a metric k such that $k(X, \xi) = \eta(X)$ from an arbitrary metric k' and the second the polarization of $d\eta$ restricted to the contact subbundle. The first step leads to a linear subspace \mathcal{L} of \mathcal{M} . If k_0 and k_1 are two Riemannian metrics for which $k_i(X, \xi) = \eta(X)$, $i = 0, 1$, then $k = (1 - t)k_0 + tk_1$ for all t for which k is positive definite is also a Riemannian metric with this property. Moreover, if k'_0 and k'_1 yield k_0 and k_1 respectively under the first step of the process, $(1 - t)k'_0 + tk'_1$ yields $k = (1 - t)k_0 + tk_1$.

We have also seen that all associated metrics have the same volume element. On a symplectic manifold of dimension 2, $\mathcal{A} = \mathcal{N}$, and on a contact manifold of dimension 3, $\mathcal{A} = \mathcal{N} \cap \mathcal{L}$. In higher dimensions \mathcal{A} is a proper subset of \mathcal{N} ($\mathcal{N} \cap \mathcal{L}$). We shall now show that \mathcal{A} is totally geodesic in \mathcal{N} and then study how polarization of a path $k_t \in \mathcal{N}$ acts as a projection onto $g_t \in \mathcal{A}$.

Theorem 4.7 *\mathcal{A} is totally geodesic in \mathcal{N} ($\mathcal{N} \cap \mathcal{L}$) in the sense that if $g \in \mathcal{A}$ and D is a symmetric tensor field satisfying $DJ + JD = 0$ ($D\phi + \phi D = 0$ and $D\xi = 0$), then $g_t = ge^{Dt}$ lies in \mathcal{A} .*

Proof. Let $J_t = Je^{Dt}$ ($\phi_t = \phi e^{Dt}$) and note that $Je^{Dt} = e^{-Dt}J$. Then

$$g_t(X, J_t Y) = g(X, e^{Dt} J e^{Dt} Y) = g(X, J Y) = \Omega(X, Y) (= d\eta(X, Y))$$

and

$$J_t^2 = Je^{Dt} J e^{Dt} = J^2 = -I \quad (\phi_t^2 = -I + \eta \otimes \xi).$$

■

In particular, the tangent space $T_g \mathcal{A}$ of \mathcal{A} at g is the set of all symmetric tensor fields that anticommute with J (ϕ and annihilate ξ).

We now turn to the problem of understanding how the polarization of a geodesic in \mathcal{N} gives rise to path $g_t \in \mathcal{A}$ and hence of viewing polarization as a projection of \mathcal{N} onto \mathcal{A} (the author [1983]). We suppress mentioning the contact case and for that case all tensors T should be understood as also satisfying $T\xi = 0$.

Let us start with a geodesic $k_t = ge^{St}$ in \mathcal{N} where $g \in \mathcal{A}$ and S any symmetric tensor field of vanishing trace. As we have seen, if S and J anticommute, k_t is already the path g_t in \mathcal{A} . We shall see as a corollary

that if S and J commute, k_t collapses to the metric g . (One can think of a matrix B that commutes with $\mathcal{J} = \begin{pmatrix} 0 & -I \\ I & 0 \end{pmatrix}$ as being orthogonal to a matrix D that anticommutes with \mathcal{J} by showing that $\text{tr}BD = 0$.) Writing S as $B + D$, we will see that the construction process takes k_t to ge^{Dt} if and only if B and D commute, but that g_t and ge^{Dt} agree through second order in general.

Let $\{X_i\}$ be a local g -orthonormal basis with respect to which J is given by the matrix $\begin{pmatrix} 0 & -I \\ I & 0 \end{pmatrix}$. The first problem is to construct a k_t -orthonormal basis. As remarked above, we also denote by S the matrix of S with respect to the initial basis $\{X_i\}$. If $SX = \lambda X$, then $e^{St}X = e^{\lambda t}X$, and hence the eigenvalues of e^{St} are analytic in t . Let Q be an orthogonal matrix diagonalizing S as in Subsection 4.3.1, i.e., $Q^{-1}SQ = \Lambda$, and set $\Delta = e^{-\frac{1}{2}\Lambda t}$ and $P = Q\Delta$ so that $P^Te^{St}P = I$. Let $X_i(t) = P_{ki}(t)X_k$. Then

$$k_t(X_i(t), X_j(t)) = g(P_{ki}X_k, e^{St}P_{lj}X_l) = P_{ki}P_{lj}(e^{St})^m_l g_{km} = P^Te^{St}P = I.$$

Thus $\{X_i(t)\}$ is a k_t -orthonormal basis, but note that $X_i(0) = Q_{ki}X_k$. Now our job is to polarize $A(t) = \Omega(X_i(t), X_j(t)) = P^TJP$, giving $F(t)$ and $G(t)$ as in Lemma 4.5.

If we are to have an expression for g_t that we can compare more easily for each t , we should express $g(t)$ with respect to the original basis $\{X_i\}$. Now $G(t) = g_t(X_i(t), X_j(t)) = P_{ki}P_{lj}g_t(X_k, X_l) = P^T(g_t(X_k, X_l))P$, but since $P = Q\Delta$, we have

$$g_t(X_k, X_l) = Q\Delta^{-1}G(t)\Delta^{-1}Q^{-1}.$$

Theorem 4.8 *If $k_t = ge^{St}$ is a geodesic in \mathcal{N} through $g \in \mathcal{A}$, then the path $g_t = g + D^{(1)}t + D^{(2)}t^2 + \dots$ in \mathcal{A} obtained by polarization is given with respect to the basis $\{X_i\}$ by*

$$D^{(l)} = \sum_{j=0}^l \frac{1}{j!2^j} \sum_{k=0}^j \binom{j}{k} (B + D)^{j-k} QG^{(l-j)}Q^{-1}(B + D)^k,$$

$G^{(l-j)}$ being given by Lemma 4.5.

The proof is by expansion of $g_t(X_k, X_l) = Q\Delta^{-1}G(t)\Delta^{-1}Q^{-1}$ using the series expansions of $\Delta^{-1} = e^{\frac{1}{2}\Lambda t}$ and $G(t)$ and noting that $\Lambda = Q^{-1}SQ$ (again see the author [1983]).

We remark that Q need not be unique in the above argument (e.g., if S does not have distinct eigenvalues), but each $G^{(k)}$ is of the form

$$Q^{-1} (\text{polynomial in } B \text{ and } D) Q$$

and therefore $D^{(k)}$ is independent of the orthogonal matrix Q diagonalizing S .

Again we list the first few terms:

$$\begin{aligned} D^{(1)} &= D, \quad D^{(2)} = \frac{D^2}{2}, \quad D^{(3)} = \frac{D^3}{6} - \frac{1}{12}(B^2D - 2BDB + DB^2), \\ D^{(4)} &= \frac{D^4}{24} + \frac{1}{96}(-4B^2D^2 + 5BDBD + 3BD^2B - 5DB^2D \\ &\quad + 5DBDB - 4D^2B^2). \end{aligned}$$

Corollary 4.3 $[B, D] = 0$ if and only if $g_t = ge^{Dt}$.

Proof. If $[B, D] = 0$, then as in Corollary 4.1,

$$QG^{(k-j)}Q^{-1} = \frac{(-1)^{k-j}}{(k-j)!} B^{k-j}.$$

Thus

$$\begin{aligned} D^{(k)} &= \sum_{j=0}^k \frac{1}{j!} (B + D)^j \frac{(-1)^{k-j}}{(k-j)!} B^{k-j} \\ &= \frac{1}{k!} \sum_{j=0}^k \binom{k}{j} (-1)^{k-j} (B + D)^j B^{k-j} = \frac{1}{k!} D^k. \end{aligned}$$

Conversely, if $D^{(k)} = \frac{1}{k!} D^k$, multiply the rest of the expression for $D^{(3)}$ on both the left and right by $-48D$ and add these two to 96 times the rest of $D^{(4)}$ to yield $-3BDBD + 3DB^2D + 3BD^2B - 3DBDB = 0$ and in turn the conclusion as in Corollary 4.1. ■

Corollary 4.4 If $k_t = ge^{St}$ and S commutes with J , then $g_t = g$.

Proof. In this case $D = 0$, and the result follows from the previous corollary. ■

Theorem 4.9 Two metrics in \mathcal{A} may be joined by a unique geodesic.

Proof. Let g_0 and g_1 be two metrics in \mathcal{A} . From what has been said so far, the problem is to find S such that $g_1 = g_0 e^S$ and $SJ_0 + J_0S = 0$. As before, let $\{X_i\}$ be a local g_0 -orthonormal basis with respect to which J_0 is given by the matrix $\mathcal{J} = \begin{pmatrix} 0 & -I \\ I & 0 \end{pmatrix}$. Then

$$\mathcal{J} = \Omega(X_i, X_j) = g_1(X_i, J_1 X_j),$$

from which $g_1 = -\mathcal{J}J_1$, where here J_1 and g_1 are regarded as the matrices of J_1 and g_1 with respect to the basis $\{X_i\}$. In particular, $-\mathcal{J}J_1$ is positive definite symmetric and hence there exists a unique real symmetric matrix S satisfying $e^S = -\mathcal{J}J_1$. Then $g_1 = g_0 e^S$, and since $J_1^2 = -I$, we have $\mathcal{J}e^S \mathcal{J}e^S = -I$, from which $e^{-S} \mathcal{J} = \mathcal{J}e^S$. Lemma 4.6 then gives $S\mathcal{J} + \mathcal{J}S = 0$. ■

4.4 Action of symplectic and contact transformations

We begin by showing that if f is a symplectomorphism or strict contact transformation and g an associated metric, then f^*g is also an associated metric.

Theorem 4.10 *Let (M, Ω) be a symplectic manifold, or respectively (M, η) a contact manifold, and f a diffeomorphism satisfying $f^*\Omega = \Omega$, respectively $f^*\eta = \eta$. Then for any associated metric g , f^*g is also an associated metric.*

Proof. In the symplectic case define J^* by $(f^*g)(X, J^*Y) = \Omega(X, Y)$. Then

$$g(f_*X, f_*J^*Y) = \Omega(X, Y) = \Omega(f_*X, f_*Y) = g(f_*X, Jf_*Y)$$

and therefore $f_*J^* = Jf_*$. Now $f_*J^{*2}X = J^2f_*X = -f_*X$, so that J^* is an almost complex structure satisfying $(f^*g)(X, J^*Y) = \Omega(X, Y)$ and hence f^*g is an associated metric.

In the case of a strict contact transformation, first note that

$$\begin{aligned} \eta(f_*\xi) &= (f^*\eta)(\xi) = \eta(\xi) = 1, \\ d\eta(f_*\xi, f_*X) &= (f^*d\eta)(\xi, X) = d\eta(\xi, X) = 0 \end{aligned}$$

and therefore $f_*\xi = \xi$. Now define ϕ^* by $(f^*g)(X, \phi^*Y) = d\eta(X, Y)$ and proceed as in the symplectic case to get $f_*\phi^* = \phi f_*$ and in turn $\phi^{*2} = -I + \eta \otimes \xi$. Also it is easy to check that $(f^*g)(\xi, X) = \eta(X)$. ■

There is a partial converse for the case where M is a compact symplectic manifold and the diffeomorphism belongs to the connected component of the identity of the diffeomorphism group; in general, the converse is not true and we will also give a couple of counterexamples (cf. Apostolov and Draghici [1999]). The diffeomorphism group of a compact manifold will be denoted by $Diff$ and the connected component of the identity by $Diff_0$. The group of symplectomorphisms will be denoted by \mathcal{S} .

Theorem 4.11 *Let (M, Ω) be a compact symplectic manifold and g an associated metric. If for a diffeomorphism $f \in Diff_0$, f^*g is also an associated metric, then $f \in \mathcal{S}$.*

Proof. For the associated metrics g and f^*g let J and \tilde{J} be the corresponding almost complex structures. Thus (J, g, Ω) and $(\tilde{J}, f^*g, \Omega)$ are almost Kähler structures, and consider the action of f^{-1} on the second structure, i.e., setting $\tilde{J}^* = f_*\tilde{J}f^{-1}_*$. Then $(\tilde{J}^*, g, f^{-1*}\Omega)$ is again an almost Kähler structure with the same metric g . Since the fundamental 2-form of a compact almost Kähler manifold is harmonic, both Ω and $f^{-1*}\Omega$ are harmonic with respect to the metric g . Now $f \in Diff_0$ and so Ω and $f^{-1*}\Omega$ represent the same cohomology class in $H^2(M, \mathbb{R})$. Therefore $\Omega - f^{-1*}\Omega$ is exact, but since it is also harmonic, it must be zero by the Hodge decomposition. Thus $f^{-1*}\Omega = \Omega$, giving $f \in \mathcal{S}$. ■

In general, the above theorem is not true and we give two counterexamples. First consider the diffeomorphism f of \mathbb{R}^4 given by $\bar{x}^1 = \frac{1}{2}(x^1 + x^2 + x^3 + x^4)$, $\bar{x}^2 = \frac{1}{2}(-x^1 + x^2 - x^3 + x^4)$, $\bar{x}^3 = \frac{1}{2}(x^1 + x^2 - x^3 - x^4)$, $\bar{x}^4 = \frac{1}{2}(-x^1 + x^2 + x^3 - x^4)$ and the symplectic form $\Omega = 2(d\bar{x}^1 \wedge d\bar{x}^3 + d\bar{x}^2 \wedge d\bar{x}^4)$. Let g be the standard Euclidean metric on \mathbb{R}^4 , which is clearly an associated metric. Now it is easy to see that $\left(\frac{\partial \bar{x}^j}{\partial x^i}\right) \in SO(4)$ but not in $Sp(4, \mathbb{R})$. Thus $f^*g = g$ but $f^*\Omega \neq \Omega$; in fact $f^*\Omega = -\Omega$.

For a compact counterexample consider almost Kähler structures of the form $(M = M_1 \times M_2, g = g_1 + g_2, \Omega = \Omega_1 + \Omega_2)$, where (M_1, g_1, Ω_1) is any almost Kähler manifold and $(M_2 = S^1 \times S^1, g_2, \Omega_2)$ is the standard product Kähler structure on $S^1 \times S^1$. Let f be the diffeomorphism $id_{M_1} \times \psi$, where ψ is the map on $S^1 \times S^1$ that interchanges the two factors. Clearly, f is an isometry of g , but it is not a \pm -symplectomorphism, since $f^*\Omega = \Omega_1 - \Omega_2$.

In [1970], Ebin proved a “slice theorem” for the set of Riemannian metrics \mathcal{M} , i.e., given a metric $g \in \mathcal{M}$ there is a neighborhood of g in

\mathcal{M} that is the product of a neighborhood of g in the orbit of g under the action of the diffeomorphism group and a submanifold orthogonal to the orbit with respect to the inner product on \mathcal{M} . The tangent space to this submanifold or “slice” at g is the kernel of the codifferential δ of g acting on second-order symmetric tensor fields (Berger–Ebin [1969]). In Theorem 4.10 we saw that if $f \in \mathcal{S}$ and $g \in \mathcal{A}$, then $f^*g \in \mathcal{A}$ and we may consider the quotient space \mathcal{A}/\mathcal{S} . Considering the group of isometries that are also symplectic transformations, Smolentsev [1995] shows that the slice theorem of Ebin can be restricted to give a slice theorem for \mathcal{A} .

This is a good point to give a technical lemma for use in Chapter 10, to show the Berger–Ebin result that a symmetric tensor field D orthogonal to an orbit of $Diff$ is in the kernel of the codifferential and to show at least that a symmetric tensor field D is orthogonal to an orbit of \mathcal{S} if and only if there exists a 2-form Ψ such that $(\delta D) \circ J = \delta \Psi$.

Lemma 4.7 *Let (M, g) be a compact orientable Riemannian manifold and for a vector field V let $v(X) = g(V, X)$. Then $\int_M V^i \theta_i dV_g = 0$ for every closed 1-form θ if and only if $v = \delta \Psi$ for some 2-form Ψ . $\int_M v_i X^i dV_g = 0$ for every vector field X if and only if $v = 0$.*

Proof. Taking θ exact, say df , we have $(v, df) = 0$ and hence $(\delta v, f) = 0$ for every smooth function f , where (\cdot, \cdot) denotes the global inner product of differential forms. Therefore $\delta v = 0$, so that $v = \omega + \delta \Psi$ for some 2-form Ψ and harmonic 1-form ω . Now taking $\theta = \omega$, $(v, \theta) = 0$ gives $(\omega, \omega) = 0$. Thus $\omega = 0$ and $v = \delta \Psi$. Conversely, for $v = \delta \Psi$ and θ closed, $(v, \theta) = (\delta \Psi, \theta) = (\Psi, d\theta) = 0$. For the second statement we already have $v = \delta \Psi$, but now let X be the contravariant form of $\delta \Psi$, then $0 = \int_M v_i X^i dV_g = (\delta \Psi, \delta \Psi)$ giving $v = 0$. ■

We now look again at the diffeomorphism group, $Diff$, of M . For a vector field X on M , let f_t be its 1-parameter subgroup in $Diff$. Then $f_t(m)$ is the integral curve of X starting at $m \in M$, so in particular,

$$\left. \frac{d}{dt} f_t(m) \right|_{t=0} = X(m).$$

Conversely, given a path f_t in $Diff$ with $f_0 = id$, we have for every $m \in M$, $\left. \frac{d}{dt} f_t(m) \right|_{t=0} \in T_m M$. Thus the tangent space to $Diff$ at the identity may be viewed as the Lie algebra \mathfrak{X} of vectors fields on M .

Now consider the orbit of \mathcal{O}_g of $g \in \mathcal{M}$ under $Diff$. We have remarked (Section 4.3) that the tangent space $T_g \mathcal{M}$ is the set of symmetric tensor

fields of type $(0, 2)$ on M and ask what are the symmetric tensor fields that are tangent to the orbit. Let $\psi_g : Diff \rightarrow \mathcal{M}$ be defined by $\psi_g(f) = f^*g$. Then $\psi_{g*} : \mathfrak{X} \rightarrow T_g\mathcal{O}_g$. Given $X \in \mathfrak{X}$, let f_t be its 1-parameter subgroup. Therefore

$$\psi_{g*}(X) = \frac{d}{dt}f_t^*g = \mathcal{L}_Xg.$$

Suppose now that a symmetric tensor field $D \in T_g\mathcal{M}$ is orthogonal to \mathcal{O}_g at g . Then

$$\begin{aligned} 0 &= (\mathcal{L}_Xg, D) = \int_M (\nabla_i X_j + \nabla_j X_i) D^{ij} dV_g \\ &= 2 \int_M (\nabla_i X_j) D^{ij} dV_g = -2 \int_M (\nabla_i D^{ij}) X_j dV_g \end{aligned}$$

for every vector field X , and hence by Lemma 4.7, $\delta D = 0$.

Now let M be a compact symplectic manifold and g an associated metric with corresponding almost complex structure J . A symmetric tensor field D is orthogonal to the orbit of g under the action of \mathcal{S} if and only if there exists a 2-form Ψ such that $(\delta D) \circ J = \delta \Psi$. To see this, first note that the tangent space to the orbit of g under the action of \mathcal{S} at g is the set of tensor fields of the form \mathcal{L}_Xg where X is a symplectic vector field. As we have seen (Theorem 1.7), $X^i = J^{ik}\theta_k$ for some closed 1-form θ . Then as in the argument above,

$$(\mathcal{L}_Xg, D) = 2 \int_M (\delta D)_i J^{ik} \theta_k dV_g,$$

and the result follows from Lemma 4.7.

4.5 Examples of almost contact metric manifolds

4.5.1 \mathbb{R}^{2n+1}

In Example 3.2.1 we considered \mathbb{R}^{2n+1} with its usual contact structure $dz - \sum_{i=1}^n y^i dx^i$ and saw that the contact subbundle \mathcal{D} is spanned by $\frac{\partial}{\partial x^i} + y^i \frac{\partial}{\partial z}$, $\frac{\partial}{\partial y^i}$, $i = 1, \dots, n$. For normalization convenience, we take as the standard contact structure on \mathbb{R}^{2n+1} the 1-form $\eta = \frac{1}{2}(dz - \sum_{i=1}^n y^i dx^i)$.

The characteristic vector field is then $\xi = 2\frac{\partial}{\partial z}$, and the Riemannian metric

$$g = \eta \otimes \eta + \frac{1}{4} \sum_{i=1}^n ((dx^i)^2 + (dy^i)^2)$$

gives a contact metric structure on \mathbb{R}^{2n+1} . For reference purposes, we give the matrix of components of g , namely

$$\frac{1}{4} \begin{pmatrix} \delta_{ij} + y^i y^j & 0 & -y^i \\ 0 & \delta_{ij} & 0 \\ -y^j & 0 & 1 \end{pmatrix}.$$

The tensor field ϕ is given by the matrix

$$\begin{pmatrix} 0 & \delta_{ij} & 0 \\ -\delta_{ij} & 0 & 0 \\ 0 & y^j & 0 \end{pmatrix},$$

and the vector fields $X_i = 2\frac{\partial}{\partial y^i}$, $X_{n+i} = 2(\frac{\partial}{\partial x^i} + y^i \frac{\partial}{\partial z})$, $i = 1, \dots, n$, and ξ form a ϕ -basis for the contact metric structure.

The Riemannian metric given here has the following properties. The vector field ξ is a Killing vector field, i.e., it generates a 1-parameter group of isometries. The sectional curvature of any plane section containing ξ is equal to 1. The sectional curvature of a plane section spanned by a vector X orthogonal to ξ and ϕX is equal to -3 ; for this reason this example is often denoted $\mathbb{R}^{2n+1}(-3)$.

In dimension 3 this example is often identified with the Heisenberg group

$$H_{\mathbb{R}} = \left\{ \begin{pmatrix} 1 & y & z \\ 0 & 1 & x \\ 0 & 0 & 1 \end{pmatrix} \mid x, y, z \in \mathbb{R} \right\};$$

left translation preserves η , and g is a left-invariant metric on $H_{\mathbb{R}}$.

We have already seen that associated metrics are not unique, and in Section 7.2 we shall give another associated metric of the contact form η on \mathbb{R}^{2n+1} that is less standard but has some interesting and basic properties.

4.5.2 $M^{2n+1} \subset \tilde{M}^{2n+2}$ almost complex

We begin with a result of Tashiro [1963] that every C^∞ orientable hypersurface of an almost complex manifold has an almost contact structure.

Let (\tilde{M}^{2n+2}, J) be an almost complex manifold and $\iota : M^{2n+1} \longrightarrow \tilde{M}^{2n+2}$ a C^∞ orientable hypersurface. There exists a transverse vector field ν along M^{2n+1} such that $J\nu$ is tangent. For if $J\iota_*X$ is tangent for every tangent vector X , $J\iota_*X = \iota_*fX$ defines a $(1, 1)$ -tensor field f on M^{2n+1} . Applying J , we have $f^2 = -I$ on M^{2n+1} , making M^{2n+1} an almost complex manifold, a contradiction. Thus there exists a vector field ξ on M^{2n+1} such that $\nu = J\iota_*\xi$ is transverse.

Define a tensor field ϕ of type $(1, 1)$ and a 1-form η on M^{2n+1} by

$$J\iota_*X = \iota_*\phi X + \eta(X)\nu; \quad (*)$$

then applying J , we have

$$-\iota_*X = \iota_*\phi^2X + \eta(\phi X)\nu - \eta(X)\iota_*\xi$$

and hence $\phi^2 = -I + \eta \otimes \xi$ and $\eta \circ \phi = 0$. Taking $X = \xi$ in equation $(*)$ gives $\nu = \iota_*\phi\xi + \eta(\xi)\nu$ and hence $\phi\xi = 0$ and $\eta(\xi) = 1$. Therefore (ϕ, ξ, η) is an almost contact structure on M^{2n+1} .

If \tilde{M}^{2n+2} is almost Hermitian with Hermitian metric \tilde{g} , set $g = \iota^*\tilde{g}$ and take ν to be a unit normal. Then $J\nu$ is tangent and defines ξ by $J\nu = -\xi$. Then again using equation $(*)$,

$$g(X, Y) = \tilde{g}(J\iota_*X, J\iota_*Y) = g(\phi X, \phi Y) + \eta(X)\eta(Y)$$

and we see that (ϕ, ξ, η, g) is an almost contact metric structure.

We can construct the usual contact structure on an odd-dimensional sphere in this way. Let S_r^{2n+1} be a sphere of radius r in \mathbb{C}^{2n+2} with its usual Kähler structure denoted as on \tilde{M}^{2n+2} above with $\tilde{\nabla}$ denoting the connection on \mathbb{C}^{2n+2} . With ν as the unit outer normal, η is the standard contact form (cf. Example 3.2.3). Since S_r^{2n+1} is an umbilical hypersurface, its second fundamental form is $\sigma(X, Y) = -\frac{1}{r}g(X, Y)\nu$. Thus, using the fact that J is parallel and the Gauss–Weingarten equations, we have

$$0 = (\tilde{\nabla}_X J)\xi = \tilde{\nabla}_X\nu - J(\nabla_X\xi + \sigma(X, \xi)) = \frac{1}{r}X - \phi\nabla_X\xi - \frac{1}{r}\eta(X)\xi,$$

where ∇ is the Levi-Civita connection of g . Applying ϕ , we have $\nabla_X\xi = -\frac{1}{r}\phi X$. This in turn yields

$$d\eta(X, Y) = \frac{1}{2}(g(\nabla_X\xi, Y) - g(\nabla_Y\xi, X)) = \frac{1}{r}g(X, \phi Y).$$

Thus for $r \neq 1$, g is not an associated metric, but this situation is easily rectified. The structure $\bar{\eta} = \frac{1}{r}\eta$, $\bar{\xi} = r\xi$, $\bar{\phi} = \phi$ and $\bar{g} = \frac{1}{r^2}g$ is a contact metric structure. Alternatively, the metric $g' = \frac{1}{r}g + (1 - \frac{1}{r})\eta \otimes \eta$ is an associated metric for the induced contact form η on S_r^{2n+1} .

Of course in general, one cannot expect the induced almost contact metric structure to be a contact metric structure. The condition for this when the ambient space is Kähler was obtained by Okumura [1966], and we have the following theorem.

Theorem 4.12 *Let M^{2n+1} be a hypersurface of a Kähler manifold \tilde{M}^{2n+2} , (ϕ, ξ, η, g) its induced almost contact metric structure and A its Weingarten map. Then (ϕ, ξ, η, g) is a contact metric structure if and only if $A\phi + \phi A = -2\phi$.*

Proof. From the Gauss–Weingarten equations we have on the one hand $\tilde{\nabla}_X \xi = \nabla_X \xi + \sigma(X, \xi)$ and on the other

$$\tilde{\nabla}_X \xi = -\tilde{\nabla}_X J\nu = JAX = \phi AX + \eta(AX)\nu.$$

Comparing gives $\nabla_X \xi = \phi AX$. Therefore

$$2d\eta(X, Y) = g(\nabla_X \xi, Y) - g(\nabla_Y \xi, X) = g((\phi A + A\phi)X, Y),$$

from which the result follows. ■

4.5.3 $S^5 \subset S^6$

First consider \mathbb{R}^7 as the imaginary part of the Cayley numbers \mathbb{O} and define a vector product by $\mathbf{u} \times \mathbf{v}$ by the imaginary part of $\mathbf{u}\mathbf{v}$. Then

$$(\mathbf{u} \times \mathbf{v}) \times \mathbf{w} - (\mathbf{u} \cdot \mathbf{w})\mathbf{v} + (\mathbf{v} \cdot \mathbf{w})\mathbf{u} = -\mathbf{u} \times (\mathbf{v} \times \mathbf{w}) + (\mathbf{u} \cdot \mathbf{w})\mathbf{v} - (\mathbf{u} \cdot \mathbf{v})\mathbf{w},$$

both sides not being $\equiv 0$ as in dimension 3. Also

$$(\mathbf{u} \times \mathbf{v}) \cdot (\mathbf{u} \times \mathbf{w}) = (\mathbf{u} \cdot \mathbf{u})(\mathbf{v} \cdot \mathbf{w}) - (\mathbf{u} \cdot \mathbf{w})(\mathbf{v} \cdot \mathbf{u}),$$

though again in dimension 7, this would not hold if the second \mathbf{u} were replaced by a fourth vector.

The unit sphere $(S^6(1), \tilde{g})$ in \mathbb{R}^7 with outer unit normal N inherits an almost complex structure J defined by $JX = N \times X$. From the above vector identities, we have

$$J^2 = N \times (N \times X) = -X,$$

$$\tilde{g}(JX, JY) = (N \times X) \cdot (N \times Y) = X \cdot Y = \tilde{g}(X, Y),$$

giving S^6 an almost Hermitian structure (J, \tilde{g}) . One can also show that $(\tilde{\nabla}_X J)Y + (\tilde{\nabla}_Y J)X = 0$, and hence that the almost Hermitian structure is *nearly Kähler* (see also Example 6.7.3 below).

Now consider the totally geodesic 5-sphere in $S^6(1) \subset \mathbb{R}^7$ defined by $x^7 = 0$ with $\nu = -\frac{\partial}{\partial x^7}$. Let (ϕ, ξ, η, g) be the induced almost contact metric structure; in particular,

$$\begin{aligned} \xi &= -J\nu = N \times \frac{\partial}{\partial x^7} = \sum_{i=1}^6 x^i \frac{\partial}{\partial x^i} \times \frac{\partial}{\partial x^7} \\ &= x^1 \frac{\partial}{\partial x^6} - x^2 \frac{\partial}{\partial x^5} - x^3 \frac{\partial}{\partial x^4} + x^4 \frac{\partial}{\partial x^3} + x^5 \frac{\partial}{\partial x^2} - x^6 \frac{\partial}{\partial x^1}; \end{aligned}$$

η is the restriction of $x^1 dx^6 - x^6 dx^1 + x^5 dx^2 - x^2 dx^5 + x^4 dx^3 - x^3 dx^4$ to S^5 and hence is the usual contact form.

Compare this with the construction of (ϕ', ξ, η, g) on

$$S^5 \subset \mathbb{R}^6(x^7 = 0) \simeq \mathbb{C}^3 = \{x^2 + ix^5, x^3 + ix^4, x^6 + ix^1\};$$

$J' \frac{\partial}{\partial x^2} = \frac{\partial}{\partial x^5}$, etc., so viewing $\mathbb{C}^3 \subset \mathbb{C}^4 \simeq \mathbb{O}$, J' is just left multiplication by $\frac{\partial}{\partial x^7}$ considered as an imaginary unit in \mathbb{O} . Then for $X \perp \xi$,

$$\phi' X = J' X = \frac{\partial}{\partial x^7} X = \frac{\partial}{\partial x^7} \times X,$$

since $\frac{\partial}{\partial x^7} \perp X$. Then $g(\phi X, \phi' X) = (N \times X) \cdot (\frac{\partial}{\partial x^7} \times X) = 0$. Therefore (ϕ, ξ, η, g) is an almost contact metric structure with η contact and ξ its characteristic vector field but is not a contact metric structure, $d\eta(X, Y) = g(X, \phi' Y) \neq g(X, \phi Y)$. The difference between ϕ and ϕ' will be seen again in Example 6.7.3 by comparison of their covariant derivatives.

4.5.4 The Boothby–Wang fibration

Let M^{2n+1} be a compact regular contact manifold and $\pi : M^{2n+1} \longrightarrow M^{2n}$ the Boothby–Wang fibration of M^{2n+1} over a symplectic manifold M^{2n} of integral class with symplectic form Ω . Let G be an associated metric for Ω and J the corresponding almost complex structure; in particular, (M^{2n}, J, G) is almost Kählerian. As we have seen, the contact

form η can be viewed as a connection form on the principal circle bundle M^{2n+1} . Thus denoting the horizontal lift by $\tilde{\pi}$, we define a tensor field ϕ on M^{2n+1} by $\phi X = \tilde{\pi} J \pi_* X$. Then, since the characteristic vector field ξ is vertical, $\phi^2 = -I + \eta \otimes \xi$ and (ϕ, ξ, η) is an almost contact structure. Now define a Riemannian metric on M^{2n+1} by $g = \pi^* G + \eta \otimes \eta$. Since $d\eta = \pi^* \Omega$, we have

$$g(X, \phi Y) = G(\pi_* X, J \pi_* Y) \circ \pi = \Omega(\pi_* X, \pi_* Y) \circ \pi = \pi^* \Omega(X, Y) = d\eta(X, Y).$$

Clearly $\eta(X) = g(X, \xi)$, and hence (ϕ, ξ, η, g) is a contact metric structure on M^{2n+1} . It is also clear that ξ is a Killing vector field, i.e., ξ generates a 1-parameter group of isometries. A contact metric structure for which the characteristic vector field is a Killing vector field is called a *K-contact structure*, a notion that we will discuss further in later chapters.

We can at this point give a topological result on compact regular contact manifolds. Since the characteristic class of the principal circle bundle $\pi : M^{2n+1} \rightarrow M^{2n}$ is $[\Omega] \in H^2(M^{2n}, \mathbb{Z})$, the bundle is nontrivial, and hence the Gysin sequence becomes

$$0 \rightarrow H^1(M^{2n}, \mathbb{R}) \xrightarrow{\pi^*} H^1(M^{2n+1}, \mathbb{R}) \rightarrow H^0(M^{2n}, \mathbb{R}) \xrightarrow{L} H^2(M^{2n}, \mathbb{R}) \rightarrow \dots$$

where L is left exterior multiplication by Ω . Now L is injective and therefore the map π^* is an isomorphism giving the following theorem of Tanno [1967a].

Theorem 4.13 *Let $\pi : M^{2n+1} \rightarrow M^{2n}$ be the Boothby–Wang fibration of a compact regular contact manifold M^{2n+1} . Then the first Betti numbers of M^{2n+1} and M^{2n} are equal.*

In Example 3.2.6 we saw that the 5-dimensional torus carries a contact structure; we note, however, that it is not regular.

Theorem 4.14 *No torus T^{2n+1} can carry a regular contact structure.*

Proof. If T^{2n+1} admitted a regular contact structure, it would be a principal circle bundle over a symplectic manifold M^{2n} by the Boothby–Wang fibration. We have just seen that the first Betti number of the base $b_1(M^{2n})$ is equal to $b_1(T^{2n+1}) = 2n + 1$. On the other hand, we have the homotopy sequence of the bundle

$$0 \rightarrow \pi_2(M^{2n}) \rightarrow \pi_1(S^1) \rightarrow \pi_1(T^{2n+1}) \rightarrow \pi_1(M^{2n}) \rightarrow 0$$

since $\pi_2(T^{2n+1}) = 0$. Now consider the universal covering space \mathbb{R}^{2n+1} of T^{2n+1} and the lift of the fibration; each circle lifts to a line, and hence the fibration of T^{2n+1} by circles has no null-homotopic fibers. Thus the map from $\pi_1(S^1)$ into $\pi_1(T^{2n+1})$ is nontrivial and hence $\pi_2(M^{2n}) = 0$. Then by the exactness of the sequence, $\pi_1(M^{2n}) = \frac{\mathbb{Z} \oplus \dots \oplus \mathbb{Z}}{\mathbb{Z}}$, and hence $b_1(M^{2n}) = 2n$, a contradiction. ■

It is also known that no torus can carry an R-contact structure, Rukimbira [1993], so in particular no torus carries a K-contact structure. For this latter fact see also Itoh [1997].

4.5.5 $M^{2n} \times \mathbb{R}$

Let (M^{2n}, J, G) be an almost Hermitian manifold with local coordinates x^1, \dots, x^{2n} and let t be the coordinate on \mathbb{R} . Then on $M^{2n} \times \mathbb{R}$ set $\eta = f dt$, $\xi = \frac{1}{f} \frac{\partial}{\partial t}$ for some nonvanishing function f . Note that $d\eta = df \wedge dt$ and therefore $\eta \wedge d\eta \equiv 0$. Without stressing notation, for simplicity set $g = G + \eta \otimes \eta$ and define ϕ by $\phi\xi = 0$ and $\phi X = JX$ for $X \perp \xi$. Then (ϕ, ξ, η, g) is an almost contact metric structure that is certainly not a contact metric structure. Generally, in this example f is taken to be identically 1. However, since for a contact metric structure the integral curves of ξ are geodesics, as we have seen, the question of whether for an almost contact metric structure the integral curves of ξ must be geodesics sometimes arises. That the integral curves of ξ need not be geodesics can be seen in this example by choosing f so that it is not independent of x^i . For then using the standard formula for the Levi-Civita connection,

$$\begin{aligned} 2g(\nabla_X Y, Z) &= Xg(Y, Z) + Yg(X, Z) - Zg(X, Y) \\ &\quad + g([X, Y], Z) + g([Z, X], Y) - g([Y, Z], X), \end{aligned}$$

we have

$$\begin{aligned} 2g\left(\nabla_\xi \xi, \frac{\partial}{\partial x^i}\right) &= g\left(\left[\frac{\partial}{\partial x^i}, \xi\right], \xi\right) - g\left(\left[\xi, \frac{\partial}{\partial x^i}\right], \xi\right) \\ &= 2g\left(\left(\frac{\partial}{\partial x^i} \frac{1}{f}\right) \frac{\partial}{\partial t}, \xi\right) = -\frac{2}{f} \frac{\partial f}{\partial x^i}. \end{aligned}$$

4.5.6 Parallelizable manifolds

Let M^{2n+1} be a parallelizable manifold with $\{X_1, \dots, X_{2n+1}\}$ a set of parallelizing vector fields. Define a Riemannian metric by $g(X_A, X_B) = \delta_{AB}$. Let $\xi = X_{2n+1}$ and η its covariant form with respect to g . Similarly let ω^i be the covariant form of X_i , $i = 1 \dots, n$, and ω^{i*} that of $X_{i^*} = X_{n+i}$. Then define ϕ by

$$\phi = \sum_{i=1}^n (\omega^i \otimes X_{i^*} - \omega^{i*} \otimes X_i),$$

and it is easy to check that (ϕ, ξ, η, g) is an almost contact metric structure.

In particular, any odd-dimensional Lie group carries an almost contact structure.

5

Integral Submanifolds and Contact Transformations

In this chapter we first discuss integral submanifolds of a contact manifold, that is, submanifolds whose tangent vectors belong to the contact subbundle. We then study contact transformations, some characterizations, their transitivity, etc., and end the chapter with several examples.

5.1 Integral submanifolds

Let M^{2n+1} be a contact manifold with contact form η . We have seen that $\eta = 0$ defines a $2n$ -dimensional subbundle \mathcal{D} called the *contact distribution* or *subbundle* and that since $\eta \wedge (d\eta)^n \neq 0$, \mathcal{D} is nonintegrable. This nonintegrability was easily visualized on \mathbb{R}^3 in Example 3.2.6.

A submanifold M^r of M^{2n+1} is called an *integral submanifold* if $\eta(X) = 0$ for every tangent vector X . It is clear that for any pair of tangent vector fields we have

$$d\eta(X, Y) = \frac{1}{2}(X\eta(Y) - Y\eta(X) - \eta([X, Y])) = 0.$$

Then in terms of associated metrics, $g(X, \phi Y) = 0$ and for this reason integral submanifolds are often called *C-totally real submanifolds*. In particular, ϕ maps tangent vectors to normal vectors; also, since ξ is a normal vector, the dimension r can be at most n . On the other hand, by the

Darboux theorem we have local coordinates $(x^1, \dots, x^n, y^1, \dots, y^n, z)$ with respect to which $\eta = dz - \sum_{i=1}^n y^i dx^i$. Therefore $x^i = \text{const}$, $z = \text{const}$ define an n -dimensional integral submanifold, and we have the following theorem.

Theorem 5.1 *Let M^{2n+1} be a contact manifold with contact form η . Then there exist integral submanifolds of the contact subbundle \mathcal{D} of dimension n but of no higher dimension.*

Continuing this theme, we have the following result of Sasaki [1964].

Theorem 5.2 *Let $(x^i, y^i, z), i = 1, \dots, n$, be local coordinates about a point $m = (x_0^i, y_0^i, z_0)$ such that $\eta = dz - \sum_{i=1}^n y^i dx^i$ on the coordinate neighborhood. In order that r linearly independent vectors $X_\lambda, \lambda = 1, \dots, r \leq n$, at m with components $(a_\lambda^i, b_\lambda^i, c_\lambda)$ be tangent to an r -dimensional integral submanifold it is necessary and sufficient that $\eta(X_\lambda) = 0$ and $d\eta(X_\lambda, Y_\mu) = 0$, that is, $c_\lambda = \sum_i y_0^i a_\lambda^i$ and $\sum_i a_\lambda^i b_\mu^i = \sum_i a_\mu^i b_\lambda^i$.*

Proof. Again the necessity is clear. To prove the sufficiency, set $c_{\lambda\mu} = \sum_i a_\lambda^i b_\mu^i$ and choose a sufficiently small neighborhood \mathcal{U} of the origin of \mathbb{R}^r with coordinates (u^1, \dots, u^r) such that

$$\begin{aligned} x^i &= x_0^i + \sum_\lambda a_\lambda^i u^\lambda, & y^i &= y_0^i + \sum_\lambda b_\lambda^i u^\lambda, \\ z &= z_0 + \sum_\lambda c_\lambda u^\lambda + \frac{1}{2} \sum_{\lambda, \mu} c_{\lambda\mu} u^\lambda u^\mu \end{aligned}$$

defines a mapping ι of \mathcal{U} into M^{2n+1} . Then $\frac{\partial x^i}{\partial u^\lambda} = a_\lambda^i$, $\frac{\partial y^i}{\partial u^\lambda} = b_\lambda^i$ and

$$\frac{\partial z}{\partial u^\lambda} = c_\lambda + \sum_\mu c_{\lambda\mu} u^\mu = \sum_i y_0^i \frac{\partial x^i}{\partial u^\lambda} + \sum_{i, \mu} \frac{\partial x^i}{\partial u^\lambda} \frac{\partial y^i}{\partial u^\mu} u^\mu = \sum_i y^i \frac{\partial x^i}{\partial u^\lambda},$$

and hence the mapping ι defines an integral submanifold of \mathcal{D} tangent to X_1, \dots, X_r at m . ■

Finally, as in the symplectic case we note the abundance of integral submanifolds of \mathcal{D} ; more precisely, we have the following result (Sasaki [1964]).

Theorem 5.3 *Given a vector $X \in \mathcal{D}$ at $m \in M^{2n+1}$ and any $r, 1 \leq r \leq n$, there exists an r -dimensional integral submanifold M^r of \mathcal{D} through m with X tangent to M^r .*

The proof is again immediate from the Darboux theorem, choosing the Darboux coordinates $(x^i, y^i, z), i = 1, \dots, n$, such that $X = \frac{\partial}{\partial y^1}(m)$.

In Chapter 1 we discussed a theorem of Weinstein (Theorem 1.4) that locally a symplectic manifold is the cotangent bundle of a Lagrangian submanifold. We now state an analogous theorem due to Lychagin [1977] (see also Kriegl–Michor [1997, p. 468]). Recall the Liouville form β on a cotangent bundle (Section 1.1, Examples 3.2.4, 3.2.5).

Theorem 5.4 *If L is an n -dimensional integral submanifold of a contact manifold (M^{2n+1}, η) , then there exist an open neighborhood \mathcal{U} of L in M^{2n+1} , an open neighborhood \mathcal{V} of the zero section in $T^*L \times \mathbb{R}$, and a diffeomorphism $f : \mathcal{U} \rightarrow \mathcal{V}$ such that $f|_L$ is the identity on L and $f^*(\beta - dt) = \eta$.*

5.2 Contact transformations

Recall that a diffeomorphism f of M^{2n+1} is a *contact transformation* if $f^*\eta = \tau\eta$ for some nonvanishing function τ and that f is a *strict contact transformation* if $\tau \equiv 1$. Clearly $f^*d\eta = d\tau \wedge \eta + \tau d\eta$, so if $d\eta$ is invariant, $(\tau - 1)d\eta = -d\tau \wedge \eta$ and hence $(\tau - 1)\eta \wedge d\eta = 0$, giving $\tau \equiv 1$. Thus f is strict if and only if $d\eta$ is invariant.

Theorem 5.5 *f is a contact transformation if and only if $X \in \mathcal{D}$ implies $f_*X \in \mathcal{D}$.*

Proof. $\eta(f_*X) = (f^*\eta)(X) = \tau\eta(X) = 0$, and conversely $0 = \eta(f_*X) = (f^*\eta)(X)$ implies that $f^*\eta$ is proportional to η . ■

Theorem 5.6 *A diffeomorphism f is a contact transformation if and only if f maps r -dimensional integral submanifolds to r -dimensional integral submanifolds.*

Proof. If f is a contact transformation and M^r an integral submanifold, then for X tangent to M^r , $f_*X \in \mathcal{D}$ by the previous result and therefore $f(M^r)$ is an integral submanifold. Conversely, given $X \in \mathcal{D}_m$, we have seen that there exists an integral submanifold M^r through m with X tangent to M^r . Now since $f(M^r)$ is an integral submanifold, $f_*X \in \mathcal{D}$ and hence f is a contact transformation. ■

If $\mathcal{L}_X\eta = \sigma\eta$ for some function σ , X is called an *infinitesimal contact transformation*. If $\sigma \equiv 0$ we say that X is a *strict infinitesimal contact transformation*.

Theorem 5.7 *A vector field X is an infinitesimal contact transformation if and only if there exists a function f on M^{2n+1} such that $X = -\frac{1}{2}\phi\nabla f + f\xi$.*

Proof. For the sufficiency we compute as follows:

$$\begin{aligned} (\mathcal{L}_X\eta)(Y) &= (X \lrcorner d\eta)(Y) + df(Y) = 2d\eta(-\frac{1}{2}\phi\nabla f + f\xi, Y) + Yf \\ &= -d\eta(\phi\nabla f, Y) + Yf = -g(\nabla f, Y) + \eta(\nabla f)\eta(Y) + Yf \\ &= (\xi f)\eta(Y). \end{aligned}$$

Conversely, $\mathcal{L}_X\eta = \sigma\eta$ implies $X\eta(Y) - \eta([X, Y]) = \sigma\eta(Y)$. Then setting $f = \eta(X)$, we have

$$2d\eta(X, Y) + Yf = \sigma\eta(Y),$$

or

$$-2g(\phi X, Y) + g(\nabla f, Y) = \sigma g(\xi, Y),$$

from which $-2\phi X + \nabla f = \sigma\xi$, and applying ϕ we have

$$X = -\frac{1}{2}\phi\nabla f + f\xi$$

as desired; note also that $\sigma = \xi f$, and hence we have the following corollary. ■

Corollary 5.1 *X is strict if and only if $\xi f = 0$.*

As in the symplectic case (Theorem 1.8) we have the following theorem of Hatakeyama [1966] on the transitivity of the group of contact transformations.

Theorem 5.8 *Let M^{2n+1} be a compact contact manifold. Then for any two points p, q , there exists a contact transformation mapping p to q . If M^{2n+1} is regular, there exists a strict contact transformation mapping p to q .*

Proof. We first prove the result for a Darboux neighborhood \mathcal{U} about $p = (0, 0, 0)$. Suppose $q = (A^i, B^i, C)$ in this coordinate system and define a function f on \mathcal{U} by $f = \sum (B^i x^i - A^i y^i) + C - \frac{1}{2} \sum A^i B^i$. Let X be the infinitesimal contact transformation generated by f (strictly speaking, X is determined by $f \in C^\infty(M)$ such that on \mathcal{U} , f is as given and f vanishes outside some larger neighborhood). Writing X as $X^i \frac{\partial}{\partial x^i} + X^{i*} \frac{\partial}{\partial y^i} + X^0 \frac{\partial}{\partial z}$, we have, since $\xi f = 0$ on \mathcal{U} , $d\eta(X, \frac{\partial}{\partial x^i}) = -\frac{1}{2} X^{i*}$. Now by Theorem 5.7,

$$\begin{aligned} d\eta\left(X, \frac{\partial}{\partial x^i}\right) &= d\eta\left(-\frac{1}{2}\phi\nabla f + f\xi, \frac{\partial}{\partial x^i}\right) = -\frac{1}{2}g\left(\phi\nabla f, \phi\frac{\partial}{\partial x^i}\right) \\ &= -\frac{1}{2}\left(g\left(\nabla f, \frac{\partial}{\partial x^i}\right) - \eta(\nabla f)\eta\left(\frac{\partial}{\partial x^i}\right)\right) \\ &= -\frac{1}{2}\frac{\partial f}{\partial x^i} + \frac{1}{2}(\xi f)\eta\left(\frac{\partial}{\partial x^i}\right) = -\frac{1}{2}B^i, \end{aligned}$$

and hence $X^{i*} = B^i$. Similarly $d\eta(X, \frac{\partial}{\partial y^i}) = \frac{1}{2}A^i$, giving $X^i = A^i$. Now $f = \eta(X) = X^0 - y^i X^i = X^0 - y^i A^i$, and so $X^0 - y^i A^i = B^i x^i - y^i A^i + C - \frac{1}{2} \sum A^i B^i$. Therefore

$$X = A^i \frac{\partial}{\partial x^i} + B^i \frac{\partial}{\partial y^i} + \left(B^i x^i + C - \frac{1}{2} \sum A^i B^i\right) \frac{\partial}{\partial z}.$$

Thus the integral curves of X in \mathcal{U} are given by $x^i = A^i t$, $y^i = B^i t$, $z = (C - \frac{1}{2} \sum A^i B^i)t + (\sum A^i B^i) \frac{t^2}{2}$. When $t = 1$ this curve is at q . Thus the corresponding 1-parameter group f_t of X gives a diffeomorphism f_1 mapping p to q . For general p and q in M^{2n+1} , the usual continuation argument gives the result.

Now for M^{2n+1} regular, the result is a consequence of the following lemma.

Lemma 5.1 *If M^{2n+1} is a compact regular contact manifold and f a C^∞ function on a neighborhood \mathcal{U} of p such that $\xi f = 0$, then there exists $\tilde{f} \in C^\infty(M^{2n+1})$ such that $\xi \tilde{f} = 0$ and on some neighborhood \mathcal{V} of p , $\tilde{f} \equiv f$.*

Proof. Let $\pi : M^{2n+1} \rightarrow M^{2n}$ be the principal circle bundle structure of M^{2n+1} . Since $\xi f = 0$, f is constant on fibers and therefore there exist a neighborhood \mathcal{V}' about $\pi(p)$ and a function f' on \mathcal{V}' such that $f = f' \circ \pi$ on $\pi^{-1}(\mathcal{V}')$. Now extend f' to a C^∞ function \tilde{f}' on M^{2n} agreeing with f' on \mathcal{V}' . Then setting $\tilde{f} = \tilde{f}' \circ \pi$, we have the desired function. ■

As in the symplectic case, Boothby [1969] mapped k points to k points; see also Kriegl–Michor [1997, p. 472].

5.3 Examples of integral submanifolds

One can readily cite some simple examples of integral submanifolds, e.g., an n -dimensional integral submanifold in \mathbb{R}^{2n+1} given by $y^i = 0$ as already noted and the fibers of T_1^*M and T_1M with the contact structures given in Example 3.2.4. We give a few more examples here, and we will give further discussion of integral submanifolds and additional examples from time to time.

5.3.1 $S^n \subset S^{2n+1}$

Consider the space \mathbb{C}^{n+1} of $n+1$ complex variables and let J be its usual almost complex structure. Let $S^{2n+1} = \{z \in \mathbb{C}^{n+1} \mid |z| = 1\}$. Then as we have seen, we can give S^{2n+1} its usual contact structure as follows. For every $z \in S^{2n+1}$ and $X \in T_z S^{2n+1}$, $\xi = -Jz$ and ϕX is the tangential part of JX . Let g be the standard metric on S^{2n+1} and η the dual 1-form of ξ . Then (ϕ, ξ, η, g) is a contact metric structure on S^{2n+1} . Now let L be an $(n+1)$ -dimensional linear subspace of \mathbb{C}^{n+1} passing through the origin and such that JL is orthogonal to L . Then, since ξ is simply the negative of the action of J on the position vector, $S^n = S^{2n+1} \cap L$ is orthogonal to ξ and is therefore an n -dimensional integral submanifold of the contact structure on S^{2n+1} . Clearly S^n is a totally geodesic integral submanifold.

5.3.2 $T^2 \subset S^5$

The following embedding of a 3-torus into the unit 5-sphere as a flat minimal submanifold is well known. Given in terms of its position vector $\mathbf{x} : T^3 \rightarrow S^5 \subset E^6 \cong \mathbb{C}^3$ it is

$$\mathbf{x} = \frac{1}{\sqrt{3}}(\cos u, \sin u, \cos v, \sin v, \cos w, \sin w).$$

Now embedding T^2 in T^3 diagonally by $u + v + w = 0$, we have a flat minimal surface in S^5 given by

$$\mathbf{x} = \frac{1}{\sqrt{3}}(\cos u, \sin u, \cos v, \sin v, \cos(u+v), -\sin(u+v)).$$

Since the characteristic vector field is given by $-J$ acting on the position vector,

$$\xi = -J\mathbf{x} = \frac{1}{\sqrt{3}}(-\sin u, \cos u, -\sin v, \cos v, \sin(u+v), \cos(u+v)).$$

Computing the tangent vectors $\mathbf{x}_u = \frac{\partial \mathbf{x}}{\partial u}$ and $\mathbf{x}_v = \frac{\partial \mathbf{x}}{\partial v}$ directly, it follows easily that $\langle \xi, \mathbf{x}_u \rangle = \langle \xi, \mathbf{x}_v \rangle = 0$ and hence that this torus is an integral surface of the contact structure on S^5 .

Examples 5.3.1 and 5.3.2 show that S^5 contains both S^2 and T^2 as integral submanifolds. It is known for topological reasons (see, e.g., Steenrod [1951, p. 144]) that S^5 does not admit a continuous field of 2-planes. Thus S^5 cannot be foliated by integral surfaces of its contact structure.

5.3.3 Legendre curves and Whitney spheres

Recall that a 1-dimensional integral submanifold of a contact manifold is called a *Legendre curve*, and we begin with an elementary property of Legendre curves in the contact manifold $(\mathbb{R}^3, \eta = dz - y dx)$. The projection γ^* of a closed Legendre curve γ in \mathbb{R}^3 to the xy -plane must have self-intersections; moreover, the algebraic (signed) area enclosed by γ^* is zero. Since $dz - y dx = 0$ along γ , this follows from the elementary formula for the area enclosed by a curve given by Green's theorem,

$$0 = - \int_{\gamma} dz = \int_{\gamma^*} -y dx = \text{area},$$

the area being $+$ for γ^* traversed counterclockwise and $-$ for clockwise. Legendre curves and their projections are discussed further in Section 8.3. Here we note that one can think of the pair of γ and its projection γ^* in the following terms. Suppose that γ itself does not have self-intersections and regard γ^* as a Lagrangian submanifold in $\mathbb{R}^2 \cong \mathbb{C}$ with self-intersections; then think of going from γ^* to γ as a way of removing the singularity but preserving the “Lagrangian–Legendre” property.

For example, the map of the circle $u^2 + v^2 = 1$ into \mathbb{R}^2 given by

$$(u, v) \longrightarrow (v, 2uv)$$

has a double point, viz. $(\pm 1, 0) \rightarrow (0, 0)$. On the other hand, the map of the circle $u^2 + v^2 = 1$ into $(\mathbb{R}^3, \eta = dz - y dx)$ given by

$$(u, v) \longrightarrow \left(2uv, v, 2u - \frac{4}{3}u^3 \right)$$

is an embedding and is a Legendre curve.

A generalization of this example and an important Lagrangian submanifold of $\mathbb{R}^{2n} \cong \mathbb{C}^n$ is the Whitney sphere. We give two descriptions of the Whitney sphere. Let $\Omega = \sum_{i=1}^n dx^i \wedge dy^i$ be the standard symplectic form on \mathbb{R}^{2n} and consider the sphere S^n in \mathbb{R}^{n+1} given by $\sum_{i=0}^n (u^i)^2 = 1$ immersed in \mathbb{R}^{2n} by

$$(u^0, \dots, u^n) \longrightarrow (u^1, \dots, u^n, 2u^0u^1, \dots, 2u^0u^n).$$

Again notice the double point $(\pm 1, 0, \dots, 0)$, and it is easy to check that this immersed sphere is a Lagrangian submanifold of \mathbb{R}^{2n} (cf. Weinstein [1977, p. 26], Morvan [1983]). In Section 1.2 we remarked that the sphere S^n can not be embedded in \mathbb{C}^n as a Lagrangian submanifold. In a related vein there are no umbilical, non-totally-geodesic Lagrangian submanifolds isometrically immersed in any complex space-form (Chen–Ogiue [1974b]).

Now embed $\sum_{i=0}^n (u^i)^2 = 1$ in the contact manifold \mathbb{R}^{2n+1} with its standard contact metric structure (Example 4.5.1) by

$$(u^0, \dots, u^n) \longrightarrow \left(2u^0u^1, \dots, 2u^0u^n, u^1, \dots, u^n, 2u^0 - \frac{4}{3}(u^0)^3 \right),$$

giving an embedded sphere as an integral submanifold of the standard contact structure. We refer to this sphere as a *contact Whitney sphere*.

The Whitney sphere is often presented in another form, which, though slightly more complicated, lends itself to natural geometric characterization. For the Whitney sphere M^n as a Lagrangian submanifold of $\mathbb{R}^{2n} \cong \mathbb{C}^n$, the immersion is

$$(u^0, \dots, u^n) \longrightarrow \frac{1}{1 + (u^0)^2} (u^1, \dots, u^n, u^0u^1, \dots, u^0u^n).$$

This submanifold satisfies the relation

$$|\mathbf{H}|^2 = \frac{n+2}{n^2(n-1)}\tau,$$

where \mathbf{H} is the mean curvature vector and τ the scalar curvature of M^n . This equality characterizes the Whitney sphere as a Lagrangian submanifold of \mathbb{C}^n . More precisely, it was proven by Borrelli, Chen and Morvan [1995] and independently by Ros and Urbano [1998] that if M^n is a Lagrangian submanifold of \mathbb{C}^n , then $|\mathbf{H}|^2 \geq \frac{n+2}{n^2(n-1)}\tau$ with equality if and

only if M^n is either totally geodesic or a (piece of a) Whitney sphere. Borrelli, Chen and Morvan [1995] and Ros and Urbano [1998] also gave the characterization that the second fundamental form σ of a Lagrangian submanifold in \mathbb{C}^n is given by

$$\sigma(X, Y) = \frac{n}{n+2}(\tilde{g}(X, Y)\mathbf{H} + \tilde{g}(JX, \mathbf{H})JY + \tilde{g}(JY, \mathbf{H})JX)$$

if and only if the submanifold is either totally geodesic or a (piece of a) Whitney sphere.

In the contact manifold \mathbb{R}^{2n+1} with its standard contact metric structure we also have a second presentation of the contact Whitney sphere as an embedded sphere and an integral submanifold of the contact structure, namely

$$(u^0, \dots, u^n) \longrightarrow \frac{1}{1 + (u^0)^2} \left(u^0 u^1, \dots, u^0 u^n, u^1, \dots, u^n, \frac{u^0}{1 + (u^0)^2} \right).$$

For this contact Whitney sphere the analogues of the above results of Borrelli, Chen and Morvan, and Ros and Urbano were given by A. Carriazo and the author [2000/2001].

5.3.4 Lift of a Lagrangian foliation, Legendre foliations

Let $\pi : M^{2n+1} \longrightarrow M^{2n}$ be the Boothby–Wang fibration of a compact regular contact manifold M^{2n+1} over a symplectic manifold M^{2n} of integral class with symplectic form Ω and recall the details of Example 4.5.4. Let L be a Lagrangian submanifold of M^{2n} and consider the set of fibers over L . Then since $\phi X = \tilde{\pi} J \pi_* X$, this is a submanifold N^{n+1} of M^{2n+1} with the property that ϕ maps the tangent space into the normal space; such a submanifold is sometimes called an *anti-invariant submanifold*. If X and Y are horizontal tangent vector fields to N^{n+1} , then

$$0 = 2g(X, \phi Y) = 2d\eta(X, Y) = X\eta(Y) - Y\eta(X) - \eta([X, Y]) = -\eta([X, Y]).$$

Thus the horizontal distribution in N^{n+1} is integrable, giving n -dimensional integral submanifolds of M^{2n+1} .

If now the base manifold has a foliation by Lagrangian submanifolds, then the bundle space will have a foliation by n -dimensional integral submanifolds. Such a foliation is called a *Legendre foliation*. M.-Y. Pang

[1990] introduced an invariant for Legendre foliations, \mathcal{F} , of contact manifolds, (M, η) , by

$$\Pi_{\mathcal{F}}(X, Y) = -(\mathcal{L}_X \mathcal{L}_Y \eta)(\xi) = 2d\eta(Y, [X, \xi]),$$

where X and Y are vector fields tangent to leaves of the foliation. Since the foliation we just described is contained in a foliation by anti-invariant submanifolds, $[X, \xi]$ is tangent to the leaves of this foliation, and hence $\Pi_{\mathcal{F}} = 0$.

Examples of Legendre foliations for which the the Pang invariant is nonzero are the non-Sasakian (κ, μ) -manifolds, and we will discuss this briefly in Section 7.3. Legendre foliations and the Pang invariant have also been studied by P. Libermann [1991], N. Jayne [1992], [1994], [1998], and B. Cappelletti Montano and L. Di Terlizzi [2008].

6

Sasakian and Cosymplectic Manifolds

In this chapter we define the normality of an almost contact structure and the notion of a Sasakian manifold as a normal contact metric manifold. We also introduce another important structure tensor, h , which will be useful in the study of non-Sasakian contact metric manifolds. As an additional topic, cosymplectic manifolds will be discussed in some detail. We also give several examples and additional commentary.

6.1 Normal almost contact structures

Recall that almost contact manifolds were defined as manifolds with structural group $U(n) \times 1$ and hence can be thought of as odd-dimensional analogues of almost complex manifolds. We now consider almost contact manifolds that are, in a sense to be defined, analogous to complex manifolds.

As is well known, an almost complex structure need not come from a complex structure. The celebrated theorem of Newlander and Nirenberg [1957] states that an almost complex structure J of class $C^{2n+\alpha}$ with vanishing Nijenhuis torsion is integrable, i.e., is the corresponding almost complex structure of a complex structure. The Nijenhuis torsion $[T, T]$

of a tensor field T of type $(1, 1)$ is the tensor field of type $(1, 2)$ given by

$$[T, T](X, Y) = T^2[X, Y] + [TX, TY] - T[TX, Y] - T[X, TY].$$

All manifolds under consideration are of class C^∞ , so the theorem of Newlander and Nirenberg applies. For detailed studies of complex manifolds see for example Goldberg [1962], Kobayashi and Nomizu [1963–69, Chapter IX], Kobayashi and Wu [1983], Morrow and Kodaira [1971], Yano [1965], Zheng [2000].

Let M^{2n+1} be an almost contact manifold with structure tensors (ϕ, ξ, η) and consider the manifold $M^{2n+1} \times \mathbb{R}$. We denote a vector field on $M^{2n+1} \times \mathbb{R}$ by $(X, f \frac{d}{dt})$, where X is tangent to M^{2n+1} , t the coordinate on \mathbb{R} , and f a C^∞ function on $M^{2n+1} \times \mathbb{R}$. Define an almost complex structure J on $M^{2n+1} \times \mathbb{R}$ by

$$J\left(X, f \frac{d}{dt}\right) = \left(\phi X - f\xi, \eta(X) \frac{d}{dt}\right);$$

that $J^2 = -I$ is easy to check. If now J is integrable, we say that the almost contact structure (ϕ, ξ, η) is *normal* (Sasaki and Hatakeyama [1961]).

Since the vanishing of the Nijenhuis torsion of J is a necessary and sufficient condition for integrability, we seek to express the condition of normality in terms of the Nijenhuis torsion of ϕ . Since $[J, J]$ is a tensor field of type $(1, 2)$, it suffices to compute $[J, J]((X, 0), (Y, 0))$ and $[J, J]((X, 0), (0, \frac{d}{dt}))$ for vector fields X and Y on M^{2n+1} :

$$\begin{aligned} [J, J]((X, 0), (Y, 0)) &= -([X, Y], 0) + \left[\left(\phi X, \eta(X) \frac{d}{dt}\right), \left(\phi Y, \eta(Y) \frac{d}{dt}\right)\right] \\ &\quad - J\left[\left(\phi X, \eta(X) \frac{d}{dt}\right), (Y, 0)\right] \\ &\quad - J\left[(X, 0), \left(\phi Y, \eta(Y) \frac{d}{dt}\right)\right] \\ &= (\phi^2[X, Y] - \eta([X, Y])\xi, 0) + \left([\phi X, \phi Y], (\phi X \eta(Y) - \phi Y \eta(X)) \frac{d}{dt}\right) \\ &\quad - \left(\phi[\phi X, Y] + (Y \eta(X))\xi, \eta([\phi X, Y]) \frac{d}{dt}\right) \\ &\quad - \left(\phi[X, \phi Y] - (X \eta(Y))\xi, \eta([X, \phi Y]) \frac{d}{dt}\right) \end{aligned}$$

$$\begin{aligned}
&= \left([\phi, \phi](X, Y) + 2d\eta(X, Y)\xi, ((\mathcal{L}_{\phi X}\eta)(Y) - (\mathcal{L}_{\phi Y}\eta)(X))\frac{d}{dt} \right), \\
[J, J]\left((X, 0), \left(0, \frac{d}{dt}\right)\right) &= \left[\left(\phi X, \eta(X)\frac{d}{dt}\right), (-\xi, 0)\right] \\
&\quad - J\left[\left(\phi X, \eta(X)\frac{d}{dt}\right), \left(0, \frac{d}{dt}\right)\right] - J[(X, 0), (-\xi, 0)] \\
&= \left(-[\phi X, \xi], (\xi\eta(X))\frac{d}{dt}\right) + \left(\phi[X, \xi], \eta([X, \xi])\frac{d}{dt}\right) \\
&= ((\mathcal{L}_{\xi}\phi)X, (\mathcal{L}_{\xi}\eta)(X)).
\end{aligned}$$

We are thus led to define four tensors $N^{(1)}$, $N^{(2)}$, $N^{(3)}$, $N^{(4)}$ by

$$\begin{aligned}
N^{(1)}(X, Y) &= [\phi, \phi](X, Y) + 2d\eta(X, Y)\xi, \\
N^{(2)}(X, Y) &= (\mathcal{L}_{\phi X}\eta)(Y) - (\mathcal{L}_{\phi Y}\eta)(X), \\
N^{(3)} &= (\mathcal{L}_{\xi}\phi)X, \\
N^{(4)} &= (\mathcal{L}_{\xi}\eta)(X).
\end{aligned}$$

Clearly the almost contact structure (ϕ, ξ, η) is normal if and only if these four tensors vanish. However, the vanishing of $N^{(1)}$ implies the vanishing of $N^{(2)}$, $N^{(3)}$ and $N^{(4)}$, so that the normality condition is simply

$$[\phi, \phi](X, Y) + 2d\eta(X, Y)\xi = 0.$$

We now prove this and other properties of these tensors (cf. Sasaki and Hatakeyama [1961], [1962]).

Theorem 6.1 *For an almost contact structure (ϕ, ξ, η) the vanishing of $N^{(1)}$ implies the vanishing of $N^{(2)}$, $N^{(3)}$ and $N^{(4)}$.*

Proof. Setting $Y = \xi$ and applying η we have $d\eta(X, \xi) = 0$, which easily gives $N^{(4)} = 0$. Then

$$0 = [\phi, \phi](X, \xi) = \phi^2[X, \xi] - \phi[\phi X, \xi] = \phi((\mathcal{L}_{\xi}\phi)X).$$

Applying ϕ and noting that $d\eta(\xi, \phi X) = 0$, implies $\eta([\xi, \phi X]) = 0$, we have $N^{(3)} = 0$. Finally, applying η to

$$0 = [\phi, \phi](\phi X, Y) + 2d\eta(\phi X, Y)\xi,$$

we have $\eta([\phi^2 X, \phi Y]) + \phi X\eta(Y) - \eta([\phi X, Y]) = 0$, which simplifies to $N^{(2)} = 0$. ■

Theorem 6.2 *For a contact metric structure (ϕ, ξ, η, g) , $N^{(2)}$ and $N^{(4)}$ vanish. Moreover, $N^{(3)}$ vanishes if and only if ξ is a Killing vector field.*

Proof. We have already seen that $N^{(4)} = 0$. Now $N^{(2)}$ can be written

$$N^{(2)}(X, Y) = 2d\eta(\phi X, Y) - 2d\eta(\phi Y, X) = 2g(\phi X, \phi Y) - 2g(\phi Y, \phi X) = 0.$$

Turning to $N^{(3)}$, we note that since $d\eta$ invariant is under the action of ξ ,

$$\begin{aligned} 0 &= (\mathcal{L}_\xi d\eta)(X, Y) = \xi g(X, \phi Y) - g([\xi, X], \phi Y) - g(X, \phi[\xi, Y]) \\ &= (\mathcal{L}_\xi g)(X, \phi Y) + g(X, (\mathcal{L}_\xi \phi)Y), \end{aligned}$$

from which we see that $N^{(3)} = 0$ if and only if ξ is Killing. ■

Next we establish a formula for the covariant derivative of ϕ for a general almost contact metric structure (ϕ, ξ, η, g) .

Lemma 6.1 *For an almost contact metric structure (ϕ, ξ, η, g) , the covariant derivative of ϕ is given by*

$$\begin{aligned} 2g((\nabla_X \phi)Y, Z) &= 3d\Phi(X, \phi Y, \phi Z) - 3d\Phi(X, Y, Z) + g(N^{(1)}(Y, Z), \phi X) \\ &\quad + N^{(2)}(Y, Z)\eta(X) + 2d\eta(\phi Y, X)\eta(Z) \\ &\quad - 2d\eta(\phi Z, X)\eta(Y). \end{aligned}$$

Proof. Recall that the Riemannian connection ∇ of g is given by

$$\begin{aligned} 2g(\nabla_X Y, Z) &= Xg(Y, Z) + Yg(X, Z) - Zg(X, Y) \\ &\quad + g([X, Y], Z) + g([Z, X], Y) - g([Y, Z], X) \end{aligned}$$

and that the coboundary formula for d on a 2-form Φ is

$$\begin{aligned} d\Phi(X, Y, Z) &= \frac{1}{3} \{X\Phi(Y, Z) + Y\Phi(Z, X) + Z\Phi(X, Y) \\ &\quad - \Phi([X, Y], Z) - \Phi([Z, X], Y) - \Phi([Y, Z], X)\}. \end{aligned}$$

Therefore

$$\begin{aligned} 2g((\nabla_X \phi)Y, Z) &= 2g(\nabla_X \phi Y, Z) + 2g(\nabla_X Y, \phi Z) \\ &= Xg(\phi Y, Z) + \phi Yg(X, Z) - Zg(X, \phi Y) \\ &\quad + g([X, \phi Y], Z) + g([Z, X], \phi Y) - g([\phi Y, Z], X) \end{aligned}$$

$$\begin{aligned}
& + Xg(Y, \phi Z) + Yg(X, \phi Z) - \phi Zg(X, Y) \\
& + g([X, Y], \phi Z) + g([\phi Z, X], Y) - g([Y, \phi Z], X) \\
= & X\Phi(Y, Z) + \phi Y(\Phi(\phi Z, X) + \eta(Z)\eta(X)) - Z\Phi(X, Y) \\
& - \Phi([X, \phi Y], \phi Z) + \eta([X, \phi Y])\eta(Z) \\
& + \Phi([Z, X], Y) - g(\phi[\phi Y, Z], \phi X) + \eta(X)\eta([Z, \phi Y]) \\
& + X\Phi(\phi Y, \phi Z) - Y\Phi(Z, X) - \phi Z(\Phi(\phi Y, X) \\
& + \eta(Y)\eta(X)) + \Phi([X, Y], Z) - \Phi([\phi Z, X], \phi Y) \\
& + \eta([\phi Z, X])\eta(Y) - g(\phi[Y, \phi Z], \phi X) + \eta(X)\eta([\phi Z, Y]) \\
& + \{\Phi([Y, Z], X) - g([Y, Z], \phi X)\} \\
& - \{\Phi([\phi Y, \phi Z], X) - g([\phi Y, \phi Z], \phi X)\} \\
& + \{g(2d\eta(Y, Z)\xi, \phi X)\} \\
= & 3d\Phi(X, \phi Y, \phi Z) - 3d\Phi(X, Y, Z) + g(N^{(1)}(Y, Z), \phi X) \\
& + N^{(2)}(Y, Z)\eta(X) + 2d\eta(\phi Y, X)\eta(Z) - 2d\eta(\phi Z, X)\eta(Y).
\end{aligned}$$

■

Corollary 6.1 *For a contact metric structure the formula of Lemma 6.1 becomes*

$$2g((\nabla_X \phi)Y, Z) = g(N^{(1)}(Y, Z), \phi X) + 2d\eta(\phi Y, X)\eta(Z) - 2d\eta(\phi Z, X)\eta(Y).$$

Taking $X = \xi$ in Corollary 6.1, we see that $\nabla_\xi \phi = 0$ for any contact metric structure. By choosing a ϕ -basis, Corollary 6.1 also yields

$$\nabla_i \phi^j{}_j = -2n\eta_j$$

on a contact metric manifold.

While our main interest is in contact manifolds, we mention, in regard to the normality of almost contact structures, papers of Sato [1977] and Geiges [1997b]. Sato proved that if a compact 3-dimensional normal almost contact manifold M is not homotopic to $S^1 \times S^2$, then $\pi_2(M) = 0$, and Geiges gives a complete classification.

6.2 The tensor field h

We have seen that on a contact metric manifold, $N^{(3)}$ vanishes if and only if ξ is Killing (Theorem 6.2) and a contact metric structure for which ξ is Killing is called a *K-contact structure*. For a general contact metric

structure the tensor field $N^{(3)}$ enjoys many important properties, and for simplicity we define a tensor field h on a contact metric manifold by

$$h = \frac{1}{2}\mathcal{L}_\xi\phi = \frac{1}{2}N^{(3)}.$$

The first property to note is immediate, namely $h\xi = 0$. We now give a number of important properties of h .

Lemma 6.2 *On a contact metric manifold h is a symmetric operator,*

$$\nabla_X\xi = -\phi X - \phi hX,$$

h anticommutes with ϕ , and $\text{tr}h = 0$.

Proof. We have already seen that on a contact metric manifold $\nabla_\xi\phi = 0$ and $\nabla_\xi\xi = 0$. Thus

$$\begin{aligned} g((\mathcal{L}_\xi\phi)X, Y) &= g(\nabla_\xi\phi X - \nabla_{\phi X}\xi - \phi\nabla_\xi X + \phi\nabla_X\xi, Y) \\ &= g(-\nabla_{\phi X}\xi + \phi\nabla_X\xi, Y), \end{aligned}$$

which vanishes if either X or Y is ξ . For X and Y orthogonal to ξ , $N^{(2)} = 0$ becomes $\eta([\phi X, Y]) + \eta([X, \phi Y]) = 0$; continuing the computation, we then have

$$\begin{aligned} g((\mathcal{L}_\xi\phi)X, Y) &= \eta(\nabla_{\phi X}Y) + \eta(\nabla_X\phi Y) \\ &= \eta(\nabla_Y\phi X) + \eta(\nabla_{\phi Y}X) \\ &= g((\mathcal{L}_\xi\phi)Y, X). \end{aligned}$$

For the second statement, using Lemma 6.1 we have

$$\begin{aligned} 2g((\nabla_X\phi)\xi, Z) &= g(\phi^2[\xi, Z] - \phi[\xi, \phi Z], \phi X) - 2d\eta(\phi Z, X) \\ &= -g(\phi(\mathcal{L}_\xi\phi)Z, \phi X) - 2g(\phi Z, \phi X) \\ &= -g((\mathcal{L}_\xi\phi)Z, X) + \eta((\mathcal{L}_\xi\phi)Z)\eta(X) - 2g(Z, X) \\ &\quad + 2\eta(Z)\eta(X) \\ &= -g((\mathcal{L}_\xi\phi)X, Z) - 2g(X, Z) + 2g(\eta(X)\xi, Z) \end{aligned}$$

and hence $-\phi\nabla_X\xi = -\frac{1}{2}(\mathcal{L}_\xi\phi)X - X + \eta(X)\xi$. Applying ϕ we then have

$$\nabla_X\xi = -\phi X - \phi hX.$$

To see the anticommutativity, note that

$$\begin{aligned} 2g(X, \phi Y) &= 2d\eta(X, Y) = g(\nabla_X \xi, Y) - g(\nabla_Y \xi, X) \\ &= g(-\phi X - \phi hX, Y) - g(-\phi Y - \phi hY, X). \end{aligned}$$

Therefore $0 = g(-\phi hX, Y) - g(Y, h\phi X)$, giving $h\phi + \phi h = 0$.

An immediate consequence of this anticommutativity is that if $hX = \lambda X$, then $h\phi X = -\lambda\phi X$. Thus if λ is an eigenvalue of h , so is $-\lambda$ and hence $\text{tr} h = 0$. \blacksquare

As a result we get the following easy corollary.

Corollary 6.2 *On a contact metric manifold $\delta\eta = 0$.*

Proof. Choosing an eigenvector basis $\{e_i\}$ of h we have

$$g(\nabla_{e_i} \xi, e_i) = g(-\phi e_i - \phi h e_i, e_i) = 0.$$


\blacksquare

Example 3.2.6 provides a nice illustration of the tensor field h . Let $\eta = \frac{1}{2}(\cos x^3 dx^1 + \sin x^3 dx^2)$ be the contact form on \mathbb{R}^3 . Then $\xi = 2(\cos x^3 \frac{\partial}{\partial x^1} + \sin x^3 \frac{\partial}{\partial x^2})$ is the characteristic vector field. The flat metric $g_{ij} = \frac{1}{4}\delta_{ij}$ is an associated metric and ϕ is given by

$$\phi = \begin{pmatrix} 0 & 0 & \sin x^3 \\ 0 & 0 & -\cos x^3 \\ -\sin x^3 & \cos x^3 & 0 \end{pmatrix}.$$

The contact subbundle \mathcal{D} is spanned by $X = -\sin x^3 \frac{\partial}{\partial x^1} + \cos x^3 \frac{\partial}{\partial x^2}$ and $\frac{\partial}{\partial x^3}$. Since the metric is Euclidean on \mathbb{R}^3 , $\nabla_X \xi = 0$. Therefore by Lemma 6.2, $h\phi X = \phi X$, but $\phi X = \frac{\partial}{\partial x^3}$, and so $h\frac{\partial}{\partial x^3} = \frac{\partial}{\partial x^3}$, i.e., $\frac{\partial}{\partial x^3}$ spans the $+1$ eigenspace of h and in turn X spans the -1 eigenspace. See figure on page 86.

Conditions on $\nabla_\xi h$ arise frequently, as we will see in later chapters, but we mention a couple at this point. Calvaruso and Perrone [2000] prove that a 3-dimensional contact metric manifold is locally homogeneous if and only if it is ball-homogeneous and $\nabla_\xi h = ah\phi$ where a is a constant. Moreover, recalling the contact circles (Example 3.2.9) of Geiges and Gonzalo [1995], Calvaruso and Perrone prove that a compact orientable 3-manifold admits a taut contact circle if and only if it admits a locally homogeneous contact metric structure satisfying $\nabla_\xi h = 0$.



6.3 Definition of a Sasakian manifold

In this short section we give an important definition, namely that of a Sasakian manifold. A *Sasakian manifold* is a normal contact metric manifold. In some respects Sasakian manifolds may be viewed as odd-dimensional analogues of Kähler manifolds. This point of view is reflected in many of the examples and results on Sasakian manifolds that will be discussed. To begin with, the following theorem is analogous to an almost Hermitian manifold being Kähler if and only if the almost complex structure is parallel with respect to the Levi-Civita connection.

Theorem 6.3 *An almost contact metric structure (ϕ, ξ, η, g) is Sasakian if and only if*

$$(\nabla_X \phi)Y = g(X, Y)\xi - \eta(Y)X.$$

Proof. The necessity follows easily from Lemma 6.1, for if (ϕ, ξ, η, g) is a normal contact metric structure, then $\Phi = d\eta$, $N^{(1)} = 0$ and $N^{(2)} = 0$,

and hence

$$\begin{aligned}
 2g((\nabla_X \phi)Y, Z) &= 2d\eta(\phi Y, X)\eta(Z) - 2d\eta(\phi Z, X)\eta(Y) \\
 &= 2(g(Y, X) - \eta(Y)\eta(X))\eta(Z) - 2g(Z, X) \\
 &\quad - \eta(Z)\eta(X))\eta(Y) \\
 &= 2g(g(X, Y)\xi - \eta(Y)X, Z).
 \end{aligned}$$

Conversely, assuming $(\nabla_X \phi)Y = g(X, Y)\xi - \eta(Y)X$, setting $Y = \xi$ gives $-\phi\nabla_X \xi = \eta(X)\xi - X$ and hence $\nabla_X \xi = -\phi X$. Therefore

$$d\eta(X, Y) = \frac{1}{2}(g(\nabla_X \xi, Y) - g(\nabla_Y \xi, X)) = g(X, \phi Y),$$

showing that (ϕ, ξ, η, g) is a contact metric structure. Now

$$[\phi, \phi](X, Y) = (\phi\nabla_Y \phi - \nabla_{\phi Y} \phi)X - (\phi\nabla_X \phi - \nabla_{\phi X} \phi)Y,$$

and straightforward substitution of the hypothesis simplifies this to

$$[\phi, \phi](X, Y) = -2d\eta(X, Y)\xi.$$

■

Recall that a contact metric structure (ϕ, ξ, η, g) is said to be *K-contact* if ξ is a Killing vector field, in particular if $h = 0$. Since $\nabla_X \xi = -\phi X$ in the above proof, $h = 0$ and we have the following corollary.

Corollary 6.3 *A Sasakian manifold is K-contact.*


In dimension 3 the converse is true, Corollary 6.5. For K-contact structures that are not Sasakian, see Example 6.7.2.

While we will see other examples of Sasakian manifolds in due course, let us show here that the standard contact metric structure on an odd-dimensional unit sphere is Sasakian. Recall that the standard contact metric on an odd-dimensional sphere was exhibited in Example 4.5.2, and in particular, if the radius is 1, the constant curvature metric is an associated metric. Using the notation there and the Kähler property of \mathbb{C}^{2n+2} , we have

$$\begin{aligned}
 0 &= (\tilde{\nabla}_X J)Y = \tilde{\nabla}_X(\phi Y + \eta(Y)\nu) - J(\nabla_X Y - g(X, Y)\nu) \\
 &= \nabla_X \phi Y - g(X, \phi Y)\nu + (X\eta(Y))\nu + \eta(Y)X \\
 &\quad - \phi\nabla_X Y - \eta(\nabla_X Y)\nu - g(X, Y)\xi \\
 &= (\nabla_X \phi)Y - g(X, Y)\xi + \eta(Y)X + ((\nabla_X \eta)(Y) - g(X, \phi Y))\nu.
 \end{aligned}$$

Taking the tangential part, we see that $(\nabla_X \phi)Y = g(X, Y)\xi - \eta(Y)X$ and hence that the structure is Sasakian.


We close this section with a couple of diagrams indicating some analogies between almost Hermitian manifolds and almost contact metric manifolds. Let $(M^{2n}J, g)$ be an almost Hermitian manifold and let Ω denote the fundamental 2-form. Then we have the following schematic array of structures.



Recall that S^6 carries an almost Hermitian structure that is neither Hermitian nor almost Kähler (cf. Example 4.5.3 and Example 6.7.3 below). The well known Calabi–Eckmann manifolds $S^{2p+1} \times S^{2q+1}$, $p, q \geq 1$, are Hermitian manifolds (see also Section 6.6 below) that are not Kähler for the topological reason that the second Betti number of a compact Kähler manifold is nonzero. As noted in Section 1.1, there are many compact symplectic (and hence almost Kähler) manifolds with no Kähler structure. Also the tangent bundle of a nonflat Riemannian manifold carries an almost Kähler structure that is not Kählerian (Dombrowski [1962]; Tachibana and Okumura [1962]; see also Section 9.1 below). Finally, there are many well-known Kähler manifolds.

The corresponding diagram for almost contact metric manifolds is the following, the notion of a K-contact manifold being intermediate between a contact metric manifold and a Sasakian manifold.

We have already seen in Example 4.5.3 that S^5 carries an almost contact metric structure that is not a contact metric structure, and we will



see in Example 6.7.3 that the structure is not normal. Cosymplectic manifolds as discussed in Section 6.5 are examples of normal almost contact metric manifolds that are not Sasakian. Since the first Betti number of a compact Sasakian manifold is even (cf. Section 6.8), odd-dimensional tori have no Sasakian structures though they have the contact structures discussed in Example 3.2.6. Also we will see in Section 9.2 that the tangent sphere bundles are in general not Sasakian. Finally, we just noted that the odd-dimensional spheres are Sasakian, and other examples are given in Section 6.7. Also in his classification of compact 3-dimensional normal almost contact manifolds, Geiges [1997b] identifies those which are normal contact (Sasakian). In particular he proves the following.

Theorem 6.4 *A compact 3-dimensional manifold admits a Sasakian structure if and only if it is diffeomorphic to a left-invariant quotient of $SU(2)$, the Heisenberg group, or $\widetilde{SL}(2, \mathbb{R})$ by a discrete group.*

A similar treatment can be found in Belgun [2000].

6.4 CR-manifolds

In this section we discuss some aspects of the relation of almost contact structures and contact metric structures to CR-structures. Let N be an n -dimensional C^∞ manifold and $T^{\mathbb{C}}N$ its complexified tangent bundle, i.e., $T_p^{\mathbb{C}}N = T_pN \otimes_{\mathbb{R}} \mathbb{C} \simeq T_pN \oplus iT_pN$. Let \mathcal{H} be a C^∞ complex subbundle

of complex dimension l . A *CR-manifold*, as introduced by Greenfield [1968], of real dimension n and *CR-dimension* l is a pair (N, \mathcal{H}) such that $\mathcal{H}_p \cap \bar{\mathcal{H}}_p = 0$ and \mathcal{H} is involutive, i.e., for vector fields $X, Y \in \mathcal{H}$, $[X, Y] \in \mathcal{H}$. Then there exist a unique subbundle \mathcal{D} of TN such that $\mathcal{D}^\mathbb{C} = \mathcal{H} \oplus \bar{\mathcal{H}}$ and a unique bundle map $\mathcal{J} : \mathcal{D} \longrightarrow \mathcal{D}$ such that $\mathcal{J}^2 = -I$ and $\mathcal{H} = \{X - i\mathcal{J}X | X \in \mathcal{D}\}$.

Now let (M, \mathcal{H}) be a CR-manifold with M of real dimension $2n + 1$ and \mathcal{H} of complex dimension n . Consider the space F_x of all covectors $f \in T_x^*M$ such that $\mathcal{D} \subseteq \ker f$. This defines a real line bundle $F \subset T^*M$. If M is orientable, then $F \longrightarrow M$ admits a global nowhere vanishing section η which is called a *pseudo-Hermitian structure* and (M, \mathcal{H}, η) is called a *pseudo-Hermitian manifold*. The *Levi form* of (M, \mathcal{H}, η) is defined by

$$L_\eta(X, Y) = -d\eta(X, \mathcal{J}Y), \quad X, Y \in \mathcal{D};$$

(M, \mathcal{H}, η) is *nondegenerate* if L_η is nondegenerate. In this case M has a natural volume form $\eta \wedge (d\eta)^n$; thus η is a contact form and its characteristic vector field ξ is transversal to \mathcal{D} . If L_η is positive definite (M, \mathcal{H}, η) is said to be *strongly pseudoconvex*. Using the direct sum decomposition $TM = \mathcal{D} \oplus \{\xi\}$ we may extend L_η to a Riemannian metric g_η on M , called the *Webster metric* (see, e.g., S. Dragomir [1995]), by $g_\eta(\xi, \xi) = 1$, $g_\eta(\xi, X) = 0$ for $X \in \mathcal{D}$, and $g_\eta(X, Y) = L_\eta(X, Y)$ for $X, Y \in \mathcal{D}$. Moreover, we may extend \mathcal{J} to a tensor field ϕ on M by $\phi\xi = 0$ and $\phi X = \mathcal{J}X$ for $X \in \mathcal{D}$. Therefore a strongly pseudoconvex CR manifold (M, \mathcal{H}, η) carries a contact metric structure $(\phi, \xi, \eta, g_\eta)$.

In [1978] Bejancu defined the notion of a CR-submanifold of an almost Hermitian manifold. A submanifold N of an almost Hermitian manifold (M, J, g) is a *CR-submanifold* if there exists a holomorphic (J -invariant) subbundle \mathcal{D} of TN , with $\mathcal{D}_p \neq \{0\}$ or $T_p N$ for any $p \in N$, and such that its orthogonal complement $\mathcal{D}^\perp \subset TN$ is totally real, i.e., $J\mathcal{D}_p^\perp \subset T_p^\perp N$. Clearly real hypersurfaces are CR-submanifolds.

Let P denote the projection from TN to \mathcal{D} and Q the projection from TN to \mathcal{D}^\perp . Now since \mathcal{D} is J -invariant, set $\mathcal{J} = JP$ and define a complex subbundle \mathcal{H} of $T^\mathbb{C}N$ by $\mathcal{H} = \{X - i\mathcal{J}X | X \in \mathcal{D}\}$. The following lemma is immediate.

Lemma 6.3 $\mathcal{J}(X - i\mathcal{J}X) = JPX - i\mathcal{J}JPX$.

Now suppose that the ambient space (M, J, g) is Hermitian, i.e., the almost complex structure J is integrable. We then have the following lemma and the theorem of B.-Y. Chen and the author [1979].

Lemma 6.4 *Let N be a CR-submanifold of a Hermitian manifold (M, J, g) . Then for $X, Y \in \mathcal{D}$, $Q([JX, Y] + [X, JY]) = 0$.*

Proof. Since M is Hermitian, $[J, J] = 0$ and hence

$$0 = [J, J](JX, Y) = -[JX, Y] - [X, JY] + J([X, Y] - [JX, JY]),$$

but $[X, Y]$ and $[JX, JY]$ are tangent to N , so $J([X, Y] - [JX, JY])$ has no \mathcal{D}^\perp -component. Therefore $[JX, Y] + [X, JY]$ has no \mathcal{D}^\perp -component. ■

Theorem 6.5 *Let M be a Hermitian manifold and N a CR-submanifold. Then N is a CR-manifold.*

Proof. Let $X, Y \in \mathcal{D}$. Then

$$\begin{aligned} [X - i\mathcal{J}X, Y - i\mathcal{J}Y] &= [X, Y] - [JX, JY] - i[JX, Y] - i[X, JY] \\ &= -J[JX, Y] - J[X, JY] - iP[JX, Y] - iP[X, JY] \end{aligned}$$

by virtue of $[J, J] = 0$ and Lemma 6.4. Continuing the computation using Lemma 6.4 again and then Lemma 6.3,

$$\begin{aligned} [X - i\mathcal{J}X, Y - i\mathcal{J}Y] &= -\mathcal{J}[JX, Y] - \mathcal{J}[X, JY] + i\mathcal{J}^2[JX, Y] \\ &\quad + i\mathcal{J}^2[X, JY] \\ &= -\mathcal{J}([JX, Y] - i\mathcal{J}[JX, Y]) - \mathcal{J}([X, JY] \\ &\quad - i\mathcal{J}[X, JY]) \in \mathcal{H}. \end{aligned}$$

■

Turning to the case of almost contact structures, consider an almost contact manifold M^{2n+1} with structure tensors (ϕ, ξ, η) . Since $\phi^2 = -I + \eta \otimes \xi$ and $\phi\xi = 0$, the eigenvalues of ϕ are 0 and $\pm i$ each with multiplicity n ; in particular, ϕ is an almost complex structure on the subbundle \mathcal{D} defined by $\eta = 0$. Thus the complexification of \mathcal{D}_p is decomposable as $\mathcal{D}'_p \oplus \mathcal{D}''_p$, where $\mathcal{D}'_p = \{X - i\phi X | X \in \mathcal{D}_p\}$ and $\mathcal{D}''_p = \{X + i\phi X | X \in \mathcal{D}_p\}$, the eigenspaces of $\pm i$ respectively.

Lemma 6.5 $\phi(X - i\phi X) \in \mathcal{D}'_p$.

Proof. $\phi(\phi(X - i\phi X)) = \phi(i(X - i\phi X)) = i\phi(X - i\phi X)$. ■

We now prove a theorem of Ianus [1972] that a normal almost contact manifold is a CR-manifold. The converse is not true, and in Theorem 6.7 we will obtain a necessary and sufficient condition for a contact metric manifold to be a CR-manifold.

Theorem 6.6 *If $(M^{2n+1}, \phi, \xi, \eta, g)$ is a normal almost contact manifold, then (M^{2n+1}, \mathcal{D}') is a CR-manifold.*

Proof. First of all, $\bar{\mathcal{D}}'_p = \mathcal{D}''_p$ and $\mathcal{D}'_p \cap \mathcal{D}''_p = 0$; thus it remains to show that $[X - i\phi X, Y - i\phi Y] \in \mathcal{D}'$ for $X, Y \in \mathcal{D}$. By the normality, $[\phi, \phi] + 2d\eta \otimes \xi = 0$, so that

$$0 = -[X, Y] + [\phi X, \phi Y] - \phi[\phi X, Y] - \phi[X, \phi Y].$$

Also since $N^{(2)} = 0$, $(\mathcal{L}_{\phi X}\eta)(Y) - (\mathcal{L}_{\phi Y}\eta)(X) = 0$, from which

$$\eta([\phi X, Y] + [X, \phi Y]) = 0.$$

Now

$$\begin{aligned} [X - i\phi X, Y - i\phi Y] &= [X, Y] - [\phi X, \phi Y] - i[\phi X, Y] - i[X, \phi Y] \\ &= -\phi[\phi X, Y] - \phi[X, \phi Y] + i\phi^2[\phi X, Y] \\ &\quad - i\eta([\phi X, Y])\xi + i\phi^2[X, \phi Y] - i\eta([X, \phi Y])\xi \\ &= -\phi([\phi X, Y] - i\phi[\phi X, Y]) - \phi([X, \phi Y] \\ &\quad - i\phi[X, \phi Y]) \in \mathcal{D}'. \quad \blacksquare \end{aligned}$$

On a contact metric manifold M^{2n+1} , (M^{2n+1}, \mathcal{D}') might be CR without the structure being normal. We present an important result of Tanno [1989] giving a necessary and sufficient condition for a contact metric manifold to be a CR-manifold.

Theorem 6.7 *Let (M^{2n+1}, η, g) be a contact metric manifold. Then the pair (M^{2n+1}, \mathcal{D}') is a (strongly pseudoconvex) CR-manifold if and only if*

$$(\nabla_X \phi)Y = g(X + hX, Y)\xi - \eta(Y)(X + hX). \quad (*)$$

Proof. The strong pseudoconvexity refers to the positive definiteness of the Levi form

$$L(X, Y) = -d\eta(X, \phi|_{\mathcal{D}}Y), \quad X, Y \in \mathcal{D}.$$

We must show the equivalence of $(*)$ and

$$[X - i\phi X, Y - i\phi Y] \in \mathcal{D}' \text{ for } X, Y \in \mathcal{D}. \quad (\dagger)$$

Since $N^{(2)} = 0$, $\eta([X, Y] - [\phi X, \phi Y]) = 0$ for $X, Y \in \mathcal{D}$, from which we can see that (\dagger) is equivalent to

$$\phi[X, Y] - \phi[\phi X, \phi Y] - [\phi X, Y] - [X, \phi Y] = 0 \quad X, Y \in \mathcal{D}. \quad (\ddagger)$$

Now replacing X, Y by $\phi X, \phi Y$, we have an expression for our condition in terms of general vector fields X, Y , viz.

$$\begin{aligned} & \phi[\phi X, \phi Y] - \phi[-X + \eta(X)\xi, -Y + \eta(Y)\xi] \\ & - [-X + \eta(X)\xi, \phi Y] - [\phi X, -Y + \eta(Y)\xi] = 0. \end{aligned}$$

Applying ϕ and changing the sign, we get

$$\begin{aligned} & [\phi, \phi](X, Y) - \eta([\phi X, \phi Y])\xi - \phi^2[\eta(X)\xi, Y] - \phi^2[X, \eta(Y)\xi] \\ & + \phi[\eta(X)\xi, \phi Y] + \phi[\phi X, \eta(Y)\xi] = 0. \end{aligned}$$

Expressing the Nijenhuis torsion and Lie brackets in terms of covariant derivatives and noting that $-\eta([\phi X, \phi Y]) = 2d\eta(\phi X, \phi Y) = 2d\eta(X, Y)$, we have

$$\begin{aligned} & (\phi \nabla_Y \phi - \nabla_{\phi Y} \phi)X - (\phi \nabla_X \phi - \nabla_{\phi X} \phi)Y + 2d\eta(X, Y)\xi \\ & + \eta(X)(\phi^2 \nabla_Y \xi - \phi \nabla_{\phi Y} \xi) - \eta(Y)(\phi^2 \nabla_X \xi - \phi \nabla_{\phi X} \xi) = 0. \end{aligned}$$

Now recall that since $g(X, \phi Y) = d\eta(X, Y)$, the cyclic sum on $\{X, Y, Z\}$ in $g((\nabla_X \phi)Y, Z)$ must vanish. Thus taking the inner product of our condition with a vector field Z and using this cyclic sum property twice, we obtain

$$\begin{aligned} & g(\phi(\nabla_Y \phi)X, Z) - g(\phi(\nabla_X \phi)Y, Z) + g((\nabla_X \phi)Z, \phi Y) + g((\nabla_Z \phi)\phi Y, X) \\ & - g((\nabla_Y \phi)Z, \phi X) - g((\nabla_Z \phi)\phi X, Y) + 2d\eta(X, Y)\eta(Z) \\ & + \eta(X)(-g((\nabla_Y \xi, Z) + g((\nabla_{\phi Y} \xi, \phi Z)) \\ & - \eta(Y)(-g((\nabla_X \xi, Z) + g((\nabla_{\phi X} \xi, \phi Z))) = 0. \end{aligned}$$

Also, since $\phi^2 = -I + \eta \otimes \xi$, we have $(\nabla_X \phi)\phi Y + \phi(\nabla_X \phi)Y = g(\nabla_X \xi, Y)\xi + \eta(Y)\nabla_X \xi$. Using this we obtain

$$\begin{aligned} & 2g((\nabla_Z \phi)\phi Y, X) - g(\nabla_Z \xi, Y)\eta(X) - \eta(Y)g(\nabla_Z \xi, X) \\ & + \eta(X)g(\nabla_{\phi Y} \xi, \phi Z) - \eta(Y)g(\nabla_{\phi X} \xi, \phi Z) = 0. \end{aligned}$$

Lemma 6.5 then yields $-(\nabla_Z \phi)\phi Y = (g(\phi Z, Y) + g(\phi hZ, Y))\xi$. Finally, replacing Y by ϕY , $(\nabla_Z \phi)Y = g(Z + hZ, Y)\xi - \eta(Y)(Z + hZ)$ as desired. \blacksquare

In contrast to the fact that not every 3-dimensional contact metric manifold is Sasakian, every 3-dimensional contact metric manifold is a strongly pseudoconvex CR-manifold, as we see in the following corollary.

Corollary 6.4 *A three-dimensional contact metric manifold is a strongly pseudoconvex CR-manifold; in particular, on a 3-dimensional contact metric manifold,*

$$(\nabla_X \phi)Y = g(X + hX, Y)\xi - \eta(Y)(X + hX).$$

Proof. The left-hand side of (‡) when restricted to \mathcal{D} is just $-\phi[\phi, \phi]$, and hence it is enough to verify (‡) on the basis $\{X, Y = \phi X\}$ of \mathcal{D} , which is straightforward. ■

Corollary 6.5 *A 3-dimensional K-contact manifold is Sasakian.*

Proof. Since being K-contact is equivalent to a contact metric manifold satisfying $h = 0$, the result follows from Corollary 6.4 and Theorem 6.3. ■

Remark. In the literature there is also the following definition of a CR-structure which does not include the integrability. A *CR-structure* on a manifold M is a contact form η together with a complex structure \mathcal{J} on the contact subbundle \mathcal{D} (see, e.g., Chern–Hamilton [1985]). Let (M, η, \mathcal{J}) be a CR-structure in this sense and set

$$\mathcal{H} = \{X - i\mathcal{J}X | X \in \mathcal{D}\} = \{Z \in \mathcal{D}^{\mathbb{C}} | \mathcal{J}Z = iZ\}.$$

If the Levi form L_η is Hermitian, then (M, η, \mathcal{J}) is called a *nondegenerate pseudo-Hermitian manifold* (the condition is equivalent to the partial integrability condition: $[X, Y] \in \mathcal{D}^{\mathbb{C}}$ for $X, Y \in \mathcal{H}$). Then (M, η, \mathcal{J}) is said to be *integrable* if \mathcal{H} is involutive, i.e., $[X, Y] \in \mathcal{D}$ for $X, Y \in \mathcal{D}$.

If L_η is positive definite, (M, η, \mathcal{J}) is called a *strongly pseudoconvex CR-manifold*. So the notion, in this sense, of a strongly pseudoconvex CR-structure on M is equivalent to our notion of a contact metric structure (ϕ, ξ, η, g) by the relations $g = L_\eta + \eta \otimes \eta$, $\mathcal{J} = \phi|_{\mathcal{D}}$, where L_η also denotes its natural extension to a $(0, 2)$ tensor field on M . Using this definition of CR-structure, Theorem 6.7 can be given in the following form: Let (M, η, g) be a contact metric manifold and (η, \mathcal{J}) the corresponding strongly pseudoconvex CR-structure. Then (η, \mathcal{J}) is integrable if and only if the condition $(*)$ of Theorem 6.7 holds. In particular, (M, η, g) is Sasakian if and only if $h = 0$ and (η, \mathcal{J}) is integrable.

We shall continue to include the integrability in our definition of a (strongly pseudoconvex) CR-manifold.

6.5 Cosymplectic manifolds and remarks on the Sasakian definition

There are two notions of a cosymplectic structure in the literature (in fact, counting “Hermitian cosymplectic” ($\delta\Omega = 0$) there are at least three). P. Libermann [1959] defines a cosymplectic manifold to be a $(2n + 1)$ -dimensional manifold admitting a closed 1-form η and a closed 2-form Φ such that $\eta \wedge \Phi^n$ is a volume element. As before, on the subbundle defined by $\eta = 0$, Φ may be polarized to give an almost contact metric structure (ϕ, ξ, η, g) for which both η and the fundamental 2-form $\Phi(X, Y) = g(X, \phi Y)$ are closed. In [1967] the author defined a *cosymplectic structure* to be a normal almost contact metric structure (ϕ, ξ, η, g) with both η and Φ closed, and we adopt this point of view here.

Corresponding to Theorem 6.3 for Sasakian structures we have the following result.

Theorem 6.8 *An almost contact metric structure (ϕ, ξ, η, g) is cosymplectic if and only if ϕ is parallel.*

Proof. Since $N^{(1)} = 0$ implies $N^{(2)} = 0$, that a cosymplectic manifold satisfies $\nabla_X \phi = 0$ follows immediately from Lemma 6.1. Conversely, $\nabla_X \phi = 0$ implies that $d\Phi = 0$ and $N^{(1)} = 2d\eta \otimes \xi$. Now

$$\begin{aligned} N^{(2)}(Y, \xi) &= (\mathcal{L}_{\phi Y} \eta)(\xi) = -\eta([\phi Y, \xi]) \\ &= -g(\xi, \nabla_{\phi Y} \xi - \nabla_{\xi} \phi Y) = g(\xi, \phi \nabla_{\xi} Y) = 0; \end{aligned}$$

so setting $Z = \xi$ in Lemma 6.1, $d\eta(\phi Y, X) = 0$ for all X . Therefore $d\eta = 0$ and in turn $N^{(1)} = 0$. ■

Since $d\eta = 0$ on a cosymplectic manifold, it is clear that the subbundle defined by $\eta = 0$ is integrable; moreover, one can show that η is parallel. Also, the projection map to the tangent spaces of the integral submanifolds, $-\phi^2 = I - \eta \otimes \xi$, is parallel. Thus from Theorem 6.8 we see that locally a cosymplectic manifold is the product of a Kähler manifold and an interval, the complex structure being the restriction of ϕ to integral submanifolds. There are, however, cosymplectic manifolds that are not globally the product of a Kähler manifold and a 1-dimensional manifold; a 3-dimensional example was given by Chinea, de Leon and Marrero [1993], and higher-dimensional examples were given by Marrero and Padron [1998].

We have remarked that a Sasakian manifold is sometimes viewed as an odd-dimensional analogue of a Kähler manifold. In view of the present theorem, the same can be said of cosymplectic manifolds. Moreover, recalling the almost complex structure $J(X, f \frac{d}{dt}) = (\phi X - f\xi, \eta(X) \frac{d}{dt})$ on $M^{2n+1} \times \mathbb{R}$ defined at the beginning of this chapter, consider the product metric $G = g + dt^2$. An easy calculation shows that

$$G\left(J\left(X, f_1 \frac{d}{dt}\right), J\left(Y, f_2 \frac{d}{dt}\right)\right) = G\left(\left(X, f_1 \frac{d}{dt}\right), \left(Y, f_2 \frac{d}{dt}\right)\right),$$

so that $(M^{2n+1} \times \mathbb{R}, J, G)$ is an almost Hermitian manifold. Denote by $\bar{\nabla}$ the Levi-Civita connection of G . If now one assumes that this structure is Kähler, then another straightforward computation gives

$$0 = (\bar{\nabla}_{(X,0)} J)(Y, 0) = \left((\nabla_X \phi)Y, (\nabla_X \eta)(Y) \frac{d}{dt} \right),$$

from which we see that M^{2n+1} is cosymplectic.

Thus we have the question of the relation between the metric structure on $M^{2n+1} \times \mathbb{R}$ and the Sasakian condition. This was discussed by Tashiro [1963] and Oubina [1985]. Let $G = e^{2\rho} G'$ be a conformal change of a Kähler metric. Then, as is well known, the respective connections are related by

$$D_X Y = D'_X Y + (X\rho)Y + (Y\rho)X - G'(X, Y)P,$$

where $P = \mathbf{grad}'\rho$, and we have

$$(D_X J)Y = (JY\rho)X - G'(X, JY)P - (Y\rho)JX + G'(X, Y)JP.$$

Now suppose that the product metric G on $M^{2n+1} \times \mathbb{R}$ is conformally equivalent to a Kähler metric G' with $\rho = -t$. Then $\mathbf{grad}'\rho = -e^{-2t} \frac{d}{dt}$. Thus for X, Y vector fields on M^{2n+1} ,

$$(D_{(X,0)} J)(Y, 0) = -\eta(Y)(X, 0) + g(X, \phi Y) \left(0, \frac{d}{dt}\right) + g(X, Y)(\xi, 0)$$

on the one hand, and on the other,

$$\begin{aligned} (D_{(X,0)} J)(Y, 0) &= D_{(X,0)} \left(\phi Y, \eta(Y) \frac{d}{dt} \right) - J(\nabla_X Y, 0) \\ &= \left(\nabla_X \phi Y, (X\eta(Y)) \frac{d}{dt} \right) - \left(\phi \nabla_X Y, \eta(\nabla_X Y) \frac{d}{dt} \right). \end{aligned}$$

Comparing the components, we see that $(\nabla_X \phi)Y = g(X, Y)\xi - \eta(Y)X$ and hence that M^{2n+1} is Sasakian. Also $(\nabla_X \eta)(Y) = g(X, \phi Y)$, since ξ is now Killing.

Conversely, it is clear that if M^{2n+1} is Sasakian, then $(D_{(X,0)}J)(Y, 0)$ is given as above. Therefore, making the inverse conformal change,

$$(D'_{(X,0)}J)(Y, 0) = 0.$$

Moreover,

$$(D_{(X,0)}J)\left(0, \frac{d}{dt}\right) = D_{(X,0)}(-\xi, 0) = (-\nabla_X \xi, 0) = (\phi X, 0),$$

and this is equal to

$$(D'_{(X,0)}J)\left(0, \frac{d}{dt}\right) - g(X, \xi)\frac{d}{dt} + \left(\phi X, \eta(X)\frac{d}{dt}\right).$$

Therefore $(D'_{(X,0)}J)(0, \frac{d}{dt}) = 0$, and similarly

$$\left(D'_{(0, \frac{d}{dt})}J\right)(Y, 0) = 0, \quad \left(D'_{(0, \frac{d}{dt})}J\right)\left(0, \frac{d}{dt}\right) = 0.$$

Combining these cases, we see that G' is a Kähler metric.

The idea of using a deformed metric on $M \times \mathbb{R}$ to characterize Sasakian manifolds can be done in another way which is sometimes taken as the definition of a Sasakian manifold (see, e.g., Boyer and Galicki [2008, Definition 6.5.15]). Let (M^m, g) be a Riemannian manifold, \mathbb{R}_+ the positive reals and

$$C(M^m) = (\mathbb{R}_+ \times M^m, dr^2 + r^2g)$$

the cone over M^m . Then (M^m, g) is Sasakian if and only if the holonomy group of $C(M^m)$ reduces to a subgroup of $U(\frac{m+1}{2})$. Thus $(\mathbb{R}_+ \times M^m, dr^2 + r^2g)$ is Kähler and $m = 2n + 1$, $n \geq 1$. Boyer and Galicki are particularly interested in defining 3-Sasakian manifolds in an analogous way; see Chapter 14 of this book, Boyer, Galicki, and Mann [1994], or Boyer and Galicki [2008, p. 477].

6.6 Products of almost contact manifolds

We continue our discussion in the last section with a look at products of almost contact manifolds. However, we stress only the statement of results rather than the details of their proofs; the interested reader may

consult the references. Let M_1 and M_2 be almost contact manifolds with almost contact structures $(\phi_i, \xi_i, \eta_i), i = 1, 2$. A. Morimoto [1963] defined an almost complex structure J on the product $M_1 \times M_2$ by

$$J(X_1, X_2) = (\phi_1 X_1 - \eta_2(X_2)\xi_1, \phi_2 X_2 + \eta_1(X_1)\xi_2);$$

that $J^2(X_1, X_2) = -(X_1, X_2)$ is an easy calculation. Morimoto then proved the following theorem.

Theorem 6.9 *J is integrable if and only if both (ϕ_1, ξ_1, η_1) and (ϕ_2, ξ_2, η_2) are normal.*

An interesting corollary of this is a result of Calabi and Eckmann [1953] that the product of two odd-dimensional spheres is a complex manifold.

Corollary 6.6 *$S^{2p+1} \times S^{2q+1}$ is a complex manifold.*

Turning again to metric considerations, let M_1 and M_2 be almost contact metric manifolds with almost contact metric structures $(\phi_i, \xi_i, \eta_i, g_i), i = 1, 2$. Capursi [1984] studied the product metric $G = g_1 + g_2$, and it is again an easy calculation that

$$G(J(X_1, X_2), J(Y_1, Y_2)) = G((X_1, X_2), (Y_1, Y_2)).$$

The result of Capursi is the following.

Theorem 6.10 *$(M_1 \times M_2, J, G)$ is Kähler if and only if both $(M_1, \phi_1, \xi_1, \eta_1, g_1)$ and $(M_2, \phi_2, \xi_2, \eta_2, g_2)$ are cosymplectic.*

In view of this and the observation of Tashiro and Oubina in the last section on how to obtain the Sasakian condition from the structure on $M \times \mathbb{R}$, one might ask how to get both M_1 and M_2 Sasakian out of the almost Hermitian structure (J, G) on $M_1 \times M_2$. To the author's knowledge this is an open question. On the other hand, a special almost contact metric structure introduced by Kenmotsu [1972] seems to play a role here.

An almost contact metric manifold (M, ϕ, ξ, η, g) is called a *Kenmotsu manifold* if it satisfies

$$(\nabla_X \phi)Y = g(\phi X, Y)\xi - \eta(Y)\phi X.$$

Kenmotsu [1972] gave a local characterization of this structure.

Theorem 6.11 *Every point of a Kenmotsu manifold has a neighborhood that is a warped product $(-\epsilon, \epsilon) \times_f V$, where $f(t) = ce^t$ and V is Kähler.*

Returning to the product space $M_1 \times M_2$, define an almost complex structure by

$$J(X_1, X_2) = (\phi_1 X_1 - e^{-2\mu} \eta_2(X_2) \xi_1, \phi_2 X_2 + e^{-2\mu} \eta_1(X_1) \xi_2).$$

Again $J^2(X_1, X_2) = -(X_1, X_2)$ is an easy calculation. Let $G = e^{2\rho} g_1 + e^{2\tau} g_2$. Then G is Hermitian if and only if $\mu = \frac{1}{2}(\rho - \tau)$. Oubina and the author [1990] then noted the following.

Theorem 6.12 *Let M_1 and M_2 be almost contact metric manifolds and \mathcal{U} a coordinate neighborhood on M_2 such that $\xi_2 = \frac{\partial}{\partial t}$. Consider the change of metric $G = e^{2\rho} g_1 + e^{2\tau} g_2$ on $M_1 \times \mathcal{U}$, where $\rho = \log(k - e^{-t})$ and $\tau = -t$ for some constant k . Then $(M_1 \times \mathcal{U}, J, G)$ is Kähler if and only if the structure on M_1 is Sasakian and the structure on \mathcal{U} is Kenmotsu.*

A variation of the above result is that if $(M_1 \times \mathcal{U}, J, G)$ is Kähler for the conformal change $\rho = \tau = -t$, then M_1 is Sasakian and the structure $(\phi_2, -\xi_2, -\eta_2, g_2)$ is Kenmotsu.

The fact that this theorem is local in regard to the second manifold M_2 is not unnatural. Even for $M_1 \times \mathbb{R}$, the 1-dimensional case for M_2 , note that the Hopf manifold $S^{2n+1} \times S^1$ is locally conformally Kähler but not globally conformally Kähler.

We close this section with a remark on a generalization of these structures. In the classification of Gray and Hervella [1980] of almost Hermitian manifolds there appears a class, \mathcal{W}_4 , of Hermitian manifolds which are closely related to locally conformally Kähler manifolds. Again consider $M_1 \times \mathbb{R}$ with the almost complex structure $J(X, f \frac{d}{dt}) = (\phi X - f\xi, \eta(X) \frac{d}{dt})$ and product metric G . Oubina [1985] introduced the notion of a *trans-Sasakian structure* as an almost contact metric structure (ϕ, ξ, η, g) for which the almost Hermitian manifold $(M_1 \times \mathbb{R}, J, G)$ belongs to the class \mathcal{W}_4 . This may be expressed by the condition

$$(\nabla_X \phi)Y = \alpha(g(X, Y)\xi - \eta(Y)X) + \beta(g(\phi X, Y)\xi - \eta(Y)\phi X)$$

for functions α and β on M , and the trans-Sasakian structure is said to be of type (α, β) . If β but not α (respectively α but not β) vanishes, the structure is α -Sasakian (resp. β -Kenmotsu) (see also Janssens and Vanhecke [1981]). Marrero [1992] showed that a trans-Sasakian manifold of dimension ≥ 5 is either α -Sasakian, β -Kenmotsu or cosymplectic.

He also showed that if M is a 3-dimensional Sasakian manifold with structure tensors (ϕ, ξ, η, g) , f a positive nonconstant function and $\bar{g} = fg + (1-f)\eta \otimes \eta$, then $(\phi, \xi, \eta, \bar{g})$ is trans-Sasakian of type $(\frac{1}{f}, \frac{1}{2}\xi(\log f))$.

6.7 Examples

6.7.1 \mathbb{R}^{2n+1}

In Example 4.5.1 we gave explicitly an associated almost contact metric structure (ϕ, ξ, η, g) to the Darboux contact form $\eta = \frac{1}{2}(dz - \sum_{i=1}^n y^i dx^i)$ on \mathbb{R}^{2n+1} . From the matrix expression for ϕ given in Example 4.5.1 it is easy to check that $[\phi, \phi] + 2d\eta \otimes \xi = 0$ and hence that this contact metric structure is Sasakian.

6.7.2 Principal circle bundles

In Example 4.5.4 we saw that a compact regular contact manifold M^{2n+1} carries a K-contact structure (ϕ, ξ, η, g) , defined in terms of the almost Kähler structure (J, G) of the base manifold M^{2n} . Since $\mathcal{L}_\xi \phi = N^{(3)} = 0$,

$$[\phi, \phi](\xi, X) + 2d\eta(\xi, X)\xi = \phi^2[\xi, X] - \phi[\xi, \phi X] = 0.$$

Now $\phi X = \tilde{\pi} J \pi_* X$, so for projectable horizontal vector fields X and Y ,

$$\begin{aligned} [\phi, \phi](X, Y) + 2d\eta(X, Y)\xi &= \tilde{\pi} J^2 \pi_* [X, Y] + [\tilde{\pi} J \pi_* X, \tilde{\pi} J \pi_* Y] \\ &\quad - \tilde{\pi} J \pi_* [\tilde{\pi} J \pi_* X, Y] - \tilde{\pi} J \pi_* [X, \tilde{\pi} J \pi_* Y] \\ &\quad + d\eta(X, Y)\xi \\ &= \tilde{\pi} J^2 [\pi_* X, \pi_* Y] + \tilde{\pi} [J \pi_* X, J \pi_* Y] \\ &\quad + \eta([\tilde{\pi} J \pi_* X, \tilde{\pi} J \pi_* Y])\xi - \tilde{\pi} J [J \pi_* X, \pi_* Y] \\ &\quad - \tilde{\pi} J [\pi_* X, J \pi_* Y] + d\eta(X, Y)\xi \\ &= \tilde{\pi} [J, J](\pi_* X, \pi_* Y) - 2(\Omega(J \pi_* X, J \pi_* Y) \circ \pi \\ &\quad - \Omega(\pi_* X, \pi_* Y) \circ \pi)\xi \\ &= \tilde{\pi} [J, J](\pi_* X, \pi_* Y). \end{aligned}$$

Thus we see that the K-contact structure (ϕ, ξ, η, g) is Sasakian if and only if the base manifold $(M^{2n} J, G)$ is Kählerian (Hatakeyama [1963]). If $(M^{2n} J, G)$ is only almost Kähler, then (ϕ, ξ, η, g) is only K-contact.

Similar to the Boothby–Wang fibration of compact regular contact manifolds, A. Morimoto [1964] obtained a fibration of compact normal almost contact manifolds with ξ regular. First, however, let $\pi : M^{2n+1} \longrightarrow M^{2n}$ be a principal circle bundle over a complex manifold M^{2n} and suppose there exists a connection form η such that $d\eta = \pi^*\Psi$, where Ψ is a form of bidegree $(1, 1)$ on M^{2n} . Then we again define ϕ by $\phi X = \tilde{\pi} J \pi_* X$, where J is the almost complex structure on M^{2n} and $\tilde{\pi}$ the horizontal lift with respect to η . Let ξ be a vertical vector field with $\eta(\xi) = 1$. Then (ϕ, ξ, η) is an almost contact structure. Noting that $\mathcal{L}_\xi \eta = 0$, since η is a connection form, and $\mathcal{L}_\xi \phi = 0$ by the definition of ϕ , a computation of $N^{(1)}$ similar to the one just given shows that M^{2n+1} is a normal almost contact manifold (Morimoto [1963]).

Conversely, we give the following theorem of Morimoto [1964] and just sketch its proof, since the major ideas have already been given.

Theorem 6.13 *Let M^{2n+1} be a compact normal almost contact manifold with structure tensors (ϕ, ξ, η) and suppose that ξ is a regular vector field. Then M^{2n+1} is the bundle space of a principal circle bundle $\pi : M^{2n+1} \longrightarrow M^{2n}$ over a complex manifold M^{2n} . Moreover, η is a connection form and the 2-form Ψ on M^{2n} such that $d\eta = \pi^*\Psi$ is of bidegree $(1, 1)$.*

Proof. In the proof of the Boothby–Wang theorem (Section 3.3) we defined the period function λ of the vector field ξ and showed that λ was constant on M^{2n+1} , which we then took to be 1. The argument (cf. Tanno [1965]) required only that $\eta(\xi) = 1$ and $\mathcal{L}_\xi \eta = N^{(4)} = 0$. Thus we again have a circle bundle structure as in the Boothby–Wang fibration with η a connection form. Now since $\mathcal{L}_\xi \phi = N^{(3)} = 0$, ϕ is projectable and we can define an almost complex structure J on M^{2n} by $JX = \pi_* \phi \tilde{\pi} X$, where $\tilde{\pi}$ denotes the horizontal lift with respect to η . That $J^2 = -I$ is immediate and

$$\begin{aligned} \tilde{\pi}[J, J](X, Y) &= -[\tilde{\pi}X, \tilde{\pi}Y] + \eta([\tilde{\pi}X, \tilde{\pi}Y])\xi + [\phi\tilde{\pi}X, \phi\tilde{\pi}Y] \\ &\quad - \eta([\phi\tilde{\pi}X, \phi\tilde{\pi}Y])\xi - \phi[\phi\tilde{\pi}X, \tilde{\pi}Y] - \phi[\tilde{\pi}X, \phi\tilde{\pi}Y] \\ &= [\phi, \phi](\tilde{\pi}X, \tilde{\pi}Y) + 2d\eta(\phi\tilde{\pi}X, \phi\tilde{\pi}Y)\xi \\ &= [\phi, \phi](\tilde{\pi}X, \tilde{\pi}Y) + 2d\eta(\tilde{\pi}X, \tilde{\pi}Y)\xi = 0, \end{aligned}$$

the last equality following from $N^{(1)} = 0$ and the next to last from $N^{(2)} = 0$. Finally, $\Psi(JX, JY) \circ \pi = d\eta(\phi\tilde{\pi}X, \phi\tilde{\pi}Y) = d\eta(\tilde{\pi}X, \tilde{\pi}Y) = \Psi(X, Y) \circ \pi$, showing that Ψ is of bidegree $(1, 1)$. ■

6.7.3 A nonnormal almost contact structure on S^5

In Example 4.5.3 we saw that S^5 inherits from the almost Hermitian structure on S^6 an almost contact metric structure different from the standard one. Recall that the almost Hermitian structure (J, \tilde{g}) on S^6 given in Example 4.5.3 is a nearly Kähler structure, i.e., $(\tilde{\nabla}_X J)X = 0$ for all vector fields X . The geometric meaning of this condition is that geodesics are holomorphically planar curves. A curve γ on an almost Hermitian manifold is *holomorphically planar* if the holomorphic section determined by its tangent field is parallel along the curve.

We will now show that the induced almost contact metric structure (ϕ, ξ, η, g) on S^5 satisfies a similar condition, namely $(\nabla_X \phi)X = 0$. This is an immediate consequence of the following theorem of the author [1971]; for notation see Example 4.5.2.

Theorem 6.14 *Let M^{2n+1} be a hypersurface of a nearly Kähler manifold \tilde{M}^{2n+2} . Then the induced almost contact metric structure (ϕ, ξ, η, g) satisfies $(\nabla_X \phi)X = 0$ if and only if the second fundamental form σ is proportional to $(\eta \otimes \eta)\nu$.*

Proof. From equation (*) in Example 4.5.2 and $\tilde{\nabla}_X Y = \nabla_X Y + \sigma(X, Y)$ we have

$$(\nabla_X \Phi)(Y, Z) = (\tilde{\nabla}_X \Omega)(Y, Z) + \tilde{g}(\sigma(X, Y), \nu)\eta(Z) - \tilde{g}(\sigma(X, Z), \nu)\eta(Y),$$

where Ω is the fundamental 2-form of the nearly Kähler structure. Interchanging X and Z and adding, we obtain

$$\begin{aligned} (\nabla_X \Phi)(Y, Z) + (\nabla_Z \Phi)(Y, X) \\ = -2\tilde{g}(\sigma(X, Z), \nu)\eta(Y) + \tilde{g}(\sigma(X, Y), \nu)\eta(Z) \\ + \tilde{g}(\sigma(Z, Y), \nu)\eta(X). \end{aligned}$$

Now if σ is proportional to $(\eta \otimes \eta)\nu$, $(\nabla_X \phi)X = 0$. Conversely, if $(\nabla_X \phi)X = 0$,

$$0 = -2\tilde{g}(\sigma(X, Z), \nu)\eta(Y) + \tilde{g}(\sigma(X, Y), \nu)\eta(Z) + \tilde{g}(\sigma(Z, Y), \nu)\eta(X).$$

Setting $Y = \xi$ gives $2\tilde{g}(\sigma(X, Z), \nu) = \tilde{g}(\sigma(X, \xi), \nu)\eta(Z) + \tilde{g}(\sigma(Z, \xi), \nu)\eta(X)$, but now setting $X = \xi$, we have $\tilde{g}(\sigma(\xi, Z), \nu) = \tilde{g}(\sigma(\xi, \xi), \nu)\eta(Z)$ and consequently $\tilde{g}(\sigma(X, Z), \nu) = \tilde{g}(\sigma(\xi, \xi), \nu)\eta(X)\eta(Z)$. ■

A cosymplectic version of this theorem was given by Goldberg in [1968a] (see also Okumura [1965]). An almost contact metric structure

(ϕ, ξ, η, g) satisfying $(\nabla_X \phi)X = 0$ is called a *nearly cosymplectic structure*. As a further justification of this name we remark, without proof, that a normal nearly cosymplectic manifold is cosymplectic (see the author's paper [1971] for details), but we do give the following two propositions.

Proposition 6.1 *On a nearly cosymplectic manifold, ξ is a Killing vector field.*

Proof. Clearly $(\nabla_\xi \phi)\xi = 0$, from which one easily obtains $\nabla_\xi \xi = 0$. Now differentiating the compatibility condition $g(\phi X, \phi Y) = g(X, Y) - \eta(X)\eta(Y)$ with respect to ξ , we obtain

$$g((\nabla_\xi \phi)X, \phi Y) + g(\phi X, (\nabla_\xi \phi)Y) = 0.$$

The nearly cosymplectic condition then gives

$$g((\nabla_X \phi)\xi, \phi Y) + g(\phi X, (\nabla_Y \phi)\xi) = 0,$$

which easily simplifies to

$$g(\nabla_X \xi, Y) + g(\nabla_Y \xi, X) = 0.$$

■

Proposition 6.2 *On a normal nearly cosymplectic manifold, $d\eta = 0$.*

Proof. Since the structure is normal, $N^{(1)}$ and $N^{(2)}$ vanish; thus setting $Y = X$ and $Z = \xi$ in Lemma 6.1 we have $d\eta(X, \phi X) = 0$ for all vector fields X . Linearizing this, we obtain $d\eta(X, \phi Y) + d\eta(Y, \phi X) = 0$, but from the vanishing of $N^{(2)}$, $d\eta(X, \phi Y) = -d\eta(\phi X, Y)$ and hence $d\eta(X, \phi Y) = 0$. Also $d\eta(X, \xi) = \frac{1}{2}(g(\nabla_X, \xi, \xi) - g(\nabla_\xi \xi, X)) = 0$, and so we obtain $d\eta = 0$. ■

Turning now to S^5 as a totally geodesic hypersurface of S^6 (Example 4.5.3), its induced structure is nearly cosymplectic by Theorem 6.14. This structure is not normal, for if it were, then by our two propositions η would be respectively coclosed and closed and hence harmonic, contradicting the vanishing of the first Betti number of S^5 .

6.7.4 $M^{2n+1} \subset \tilde{M}^{2n+2}$

We have already seen that the odd-dimensional spheres are Sasakian manifolds and that this structure may be obtained both by viewing the sphere as a hypersurface in \mathbb{C}^{n+1} and by considering the Hopf fibration $\pi : S^{2n+1} \longrightarrow \mathbb{C}P^n$ as a special case of the Boothby–Wang fibration. In Theorem 4.12 we saw the condition for the induced almost contact metric structure on a hypersurface of a Kähler manifold to be contact metric. Now similar to Theorem 6.14 we give the condition for the hypersurface to be Sasakian; this is a result of Tashiro [1963] and we will omit the proof since it is similar to the proof of Theorem 6.14.

Theorem 6.15 *Let M^{2n+1} be a hypersurface of a Kähler manifold \tilde{M}^{2n+2} . Then the induced almost contact metric structure (ϕ, ξ, η, g) is Sasakian if and only if the second fundamental form $\sigma = (-g + \beta(\eta \otimes \eta))\nu$, for some function β .*

An almost contact metric structure (ϕ, ξ, η, g) is said to be *nearly Sasakian* if $(\nabla_X \phi)Y + (\nabla_Y \phi)X = 2g(X, Y)\xi - \eta(X)Y - \eta(Y)X$. The above theorem also holds for this structure on a hypersurface of a nearly Kähler manifold (Showers, Yano and the author [1976]). Similar to the structure on S^5 obtained in Examples 4.5.3 and 6.7.3, consider S^5 as an umbilical hypersurface of the unit sphere S^6 at a “latitude” of 45° , so that $\sigma(X, Y) = -g(X, Y)\nu$. Then the induced almost contact metric structure is nearly Sasakian but not Sasakian.

For other results on hypersurfaces of Kähler manifolds see Okumura [1964a, 1964b, 1966], Vernon [1987].

6.7.5 Brieskorn manifolds

In this section we will show that the Brieskorn manifolds admit Sasakian structures which are often nonregular.

Consider \mathbb{C}^{n+1} with coordinates $z = (z_0, \dots, z_n)$ and let (a_0, \dots, a_n) be an $(n+1)$ -tuple of positive integers. A polynomial of the form $P(z) = z_0^{a_0} + \dots + z_n^{a_n}$ is called a *Brieskorn polynomial*, and we let $V^{2n}(a_0, \dots, a_n)$, or just V^{2n} , denote the zero set of P . Then $\Sigma^{2n-1}(a_0, \dots, a_n) = V^{2n} \cap S^{2n+1}(1)$, or just Σ^{2n-1} , is called a *Brieskorn manifold*.

For $w \in \mathbb{C}$ define $f_w : \mathbb{C}^{n+1} \longrightarrow \mathbb{C}^{n+1}$ by

$$f_w(z) = (e^{\frac{mw}{a_0}} z_0, \dots, e^{\frac{mw}{a_n}} z_n),$$

where m is the least common multiple of (a_0, \dots, a_n) . $G_{\mathbb{C}} = \{f_w | w \in \mathbb{C}\}$ is an abelian group of diffeomorphisms of \mathbb{C}^{n+1} . Let $G_{\mathbb{R}}$ be the 1-parameter subgroup given by $\{f_s | s \in \mathbb{R}\}$. The \mathbb{R} -action leaves V^{2n} invariant, and differentiating with respect to s at $s = 0$, we see that the vector field that generates $G_{\mathbb{R}}$ is

$$\mathbf{a} = \left(\frac{m}{a_{\alpha}} z_{\alpha} \right).$$

Now let $G_S = \{f_{it} | t \in \mathbb{R}\}$; f_{it} equals $f_{i(t+2\pi)}$ and is an S^1 -action on \mathbb{C}^{n+1} leaving Σ^{2n-1} invariant. The vector field that generates G_S is

$$\mathbf{b} = i\mathbf{a} = \left(\frac{m}{a_{\alpha}} iz_{\alpha} \right).$$

The almost complex structure J on \mathbb{C}^{n+1} restricts to V^{2n} , and $\xi = -\mathbf{b} = -J\mathbf{a}$ is a vector field tangent to Σ^{2n-1} . Moreover, for an arbitrary vector field tangent to Σ^{2n-1} , decomposing JX as

$$JX = \phi X + \eta(X)\mathbf{a}$$

gives Σ^{2n-1} an almost contact structure (ϕ, ξ, η) as in Example 4.5.2; see Sasaki and Takahashi [1976] and Sasaki [1985]. (In Sasaki [1985], the structure tensors have the opposite sign from our construction here.) Sasaki and Takahashi prove the following.

Theorem 6.16 *The almost contact structure (ϕ, ξ, η) on the Brieskorn manifold Σ^{2n-1} is normal.*

Recalling the theorem of Morimoto, Theorem 6.9, we have following result of Brieskorn and Van de Ven [1968].

Corollary 6.7 *Let Σ^{2n-1} and Σ^{2m-1} be Brieskorn manifolds. Then $\Sigma^{2n-1} \times \mathbb{R}$ and $\Sigma^{2n-1} \times \Sigma^{2m-1}$ are complex manifolds.*

In regard to nonregularity, Sasaki and Takahashi prove the following.

Theorem 6.17 *The almost contact structure (ϕ, ξ, η) on the Brieskorn manifold $\Sigma^{2n-1}(a_0, \dots, a_n)$, is nonregular if and only if there exist three positive integers a_{λ} , a_{μ} , a_{ν} among the positive integers a_0, \dots, a_n such that the least common multiple of a_{λ} and a_{μ} is not equal to the least common multiple of a_{λ} , a_{μ} , a_{ν} .*

Now let H be the inner product of \mathbf{a} with the position vector $z \in \mathbb{C}^{n+1}$, i.e., denoting the inner product on \mathbb{C}^{n+1} by $\langle \cdot, \cdot \rangle$, $H = \langle \mathbf{a}, z \rangle = m \sum_{\alpha=0}^n \frac{z_{\alpha} \bar{z}_{\alpha}}{a_{\alpha}}$. Taking the inner product of $JX = \phi X + \eta(X)\mathbf{a}$ with the position vector z , we have $H\eta(X) = \langle JX, z \rangle = -\langle X, Jz \rangle$. Thus if $\iota : \Sigma^{2n-1} \rightarrow \mathbb{C}^{n+1}$ denotes the embedding and $\omega = \frac{i}{2} \sum_{\alpha=0}^n (\bar{z}_{\alpha} dz_{\alpha} - z_{\alpha} d\bar{z}_{\alpha})$, then $\eta = \iota^* \omega / H$ and $\iota^* \omega$ are contact forms on Σ^{2n-1} . This was shown by Sasaki and Hsu [1976] and simpler proofs were given by Abe [1977] and Vaisman [1978].

Turning to the question of an associated metric for η , set

$$g(X, Y) = \frac{1}{H} (\langle X, Y \rangle - \eta(X)\langle Y, \xi \rangle - \eta(Y)\langle X, \xi \rangle + \eta(X)\eta(Y)\langle \xi, \xi \rangle) + \eta(X)\eta(Y).$$

Note that g is not the induced metric from the embedding $\iota : \Sigma^{2n-1} \rightarrow \mathbb{C}^{n+1}$. Sasaki [1985] then showed that (ϕ, ξ, η, g) is a Sasakian structure on Σ^{2n-1} .

Thus we see that there are in fact many nonregular Sasakian manifolds. Much of the above goes over to more general spaces. For discussion of these generalizations, see the references mentioned as well as Abe [1976], Abe and Erbacher [1975], Lutz and Meckert [1976] and Sato [1977]. An earlier example of a nonregular Sasakian manifold was given by Tanno [1969].

In recent years, interest in the homotopy spheres as Sasakian manifolds has increased. For example, Boyer, Galicki and Nakamaye [2003a] prove the following result.

Theorem 6.18 *Every homotopy sphere of dimension ≥ 5 that can be realized as the boundary of a parallelizable manifold admits a Sasakian metric of positive Ricci curvature.*

For more information on Sasakian structures on homotopy spheres see the book of Boyer and Galicki [2008, Chapter 9].

6.8 Some early topology

In the 1960s, a great deal of work was done on the topology of compact Sasakian and to a lesser extent cosymplectic manifolds. The idea was to see how much a compact Sasakian manifold must be like a sphere. In the

case of a compact Kähler manifold, the even-dimensional Betti numbers are different from zero and the odd-dimensional Betti numbers are even, properties which are certainly enjoyed by complex projective space (see, e.g., Goldberg [1962, Chapter V]). Furthermore, the Betti numbers b_p of a compact Kähler manifold of positive constant holomorphic curvature are equal to 1 for p even and vanish for p odd (see, e.g., Goldberg [1962, Chapter VI]).

In [1965] Tachibana proved that the first Betti number of a compact Sasakian manifold M^{2n+1} is zero or even. This is proved by first showing that on a compact K-contact manifold, a harmonic 1-form ω is orthogonal to the contact form η . Then letting $\tilde{\omega} = \omega \circ \phi$ and computing the Laplacian of $\tilde{\omega}$, one obtains the harmonicity of $\tilde{\omega}$ as well. Thus the number of independent harmonic 1-forms is even. The computation uses the fact that on a Sasakian manifold the Ricci operator commutes with ϕ , a fact that we will prove in our subsequent discussions of curvature. More generally, the p th Betti number is even for p odd and $1 \leq p \leq n$ and by duality for p even and $n+1 \leq p \leq 2n$ (Fujitani [1966]; see also S. I. Goldberg and the author [1967]).

For a regular Sasakian manifold M^{2n+1} fibering over a Kähler manifold M^{2n} , the Betti numbers are related by $b_p(M^{2n+1}) = b_p(M^{2n}) - b_{p-2}(M^{2n})$, $0 \leq p \leq n$ (see Boyer and Galick: [2008 chapter 7]).

In Section 3.4 we noted the result of Rukimbira [1995a] that on a compact K-contact manifold M^{2n+1} , ξ has at least $n+1$ closed orbits. In the same paper he proves that a compact Sasakian manifold for which ξ has only a finite number of closed orbits has vanishing first Betti number.

Considerable attention has been given to the vanishing of the second Betti number of a Sasakian manifold under some curvature restrictions as well as such a manifold being isometric to the unit sphere under stronger conditions. A compact Sasakian manifold of strictly positive curvature has vanishing second Betti number (Moskal [1966], Tachibana and Ogawa [1966], Tanno [1968], Goldberg [1967] in the regular case and Goldberg [1968b] in the regular case with nonnegative curvature). A compact, simply connected Sasakian Einstein space of strictly positive curvature is isometric to the unit sphere (Moskal [1966]). Pinching theorems have been obtained by Tanno [1968] including an analogue of holomorphic pinching.

In [1986] Goldberg showed that a compact simply connected regular Sasakian manifold M of strictly positive curvature is homeomorphic

to a sphere. If, in addition, M has constant scalar curvature, then Goldberg had shown earlier [1967] that M is isometric to a sphere, but not necessarily with a constant curvature metric (cf. the metrics on the sphere in Example 7.8.1).

Allowing some negative curvature, Tanno [1968] showed that if M^{2n+1} is a compact K-contact manifold with sectional curvature greater than $\frac{-3}{2n-1}$, then $b_1 = 0$. Similarly, if the Ricci tensor ρ is such that $\rho + 2g$ is positive definite, then $b_1 = 0$. By duality in dimension 3, one also has $b_2 = 0$.

In dimension 5, Perrone [1989] showed that if M^5 is a compact simply connected regular Sasakian manifold with $b_2 = 0$ and with scalar curvature $\tau > -4$, then M^5 is homeomorphic to a sphere. If, in addition, M^5 has constant scalar curvature, M is isometric to a sphere (but not necessarily with a constant curvature metric).

A classical result on the topology of a compact Kähler manifold M^{2n} is the monotonicity of the Betti numbers, namely $b_p \leq b_{p+2}$, $p \leq n-1$; this is proved using the idea of effective harmonic forms (see Goldberg [1962, Chapter V]). We briefly describe the idea of effective forms in the almost contact context. Let $(M^{2n+1}, \phi, \xi, \eta, g)$ be a compact almost contact manifold, and as before, we denote the fundamental 2-form by Φ . Define two operators \mathbf{L} and $\mathbf{\Lambda}$ acting on differential forms α by

$$\mathbf{L}\alpha = \alpha \wedge \Phi, \quad \mathbf{\Lambda}\alpha = *\mathbf{L}*\alpha,$$

where $*$ denotes the Hodge star isomorphism ($\mathbf{L}\alpha = \alpha \wedge \Omega$ in the symplectic case). A p -form α is said to be *effective* if $\mathbf{\Lambda}\alpha = 0$. We now have the following proposition (see Goldberg and the author [1967], Chinea, de Leon and Marrero [1993]).

Proposition 6.3 *On an almost contact metric manifold of dimension $2n+1$, every p -form α with $p \leq n+1$ may be written uniquely as a sum*

$$\alpha = \sum_{k=0}^r \mathbf{L}^k \beta_{p-2k},$$

where the β_{p-2k} 's, $0 \leq k \leq r$, are effective forms of degree $p-2k$ and $r = \lfloor \frac{p}{2} \rfloor$.

In the cosymplectic case we cite the work of Chinea, de Leon and Marrero [1993]. They prove that on a compact cosymplectic manifold

M^{2n+1} , $b_0 \leq b_1 \leq \cdots \leq b_n = b_{n+1}$ and $b_{n+1} \geq b_{n+2} \geq \cdots \geq b_{2n+1}$. Moreover, the differences $b_{2p+1} - b_{2p}$ with $0 \leq p \leq n$ are even, and so in particular, the first Betti number of M^{2n+1} is odd. They also prove a strong Lefschetz property for a compact cosymplectic manifold and show that such a manifold is formal (i.e., the homotopy type of the differential graded algebra of differential forms is the same as the homotopy type of the cohomology ring).

This topological work involves generalizing from Kähler geometry notions of the bidegree of differential forms, effective harmonic forms, etc. For studies of the *tridegree* of differential forms and applications, see Chineza, de Leon and Marrero [1997], Moskal [1977], and Fujitani [1966]. A p -form α is said to be *coeffective* if $\mathbf{L}\alpha = 0$. Coeffective cohomology was studied in the symplectic case by Bouche [1990] and in the almost cosymplectic case by Chineza, de Leon and Marrero [1995]. In both papers the relation between the coeffective and the de Rham cohomologies of the manifolds is discussed (for the almost contact case see also M. Fernández, R. Ibáñez and M. de Leon [1997], [1998]).

Curvature of Contact Metric Manifolds

In this chapter we discuss many aspects of the curvature of contact metric manifolds. As such, it is to be regarded as one of the most important chapters in this book.

7.1 Basic curvature properties

We begin with some preliminaries concerning the tensor field h . Let M^{2n+1} be a contact metric manifold with structure tensors (ϕ, ξ, η, g) and $h = \frac{1}{2}\mathcal{L}_\xi\phi$ as before. Recall that in Lemma 6.2 we saw that $\nabla_X\xi = -\phi X - \phi hX$.

Proposition 7.1 *On a contact metric manifold M^{2n+1} we have the following formulas:*

$$\begin{aligned}(\nabla_\xi h)X &= \phi X - h^2\phi X - \phi R_{X\xi}\xi, \\ \frac{1}{2}(R_{\xi X}\xi - \phi R_{\xi\phi X}\xi) &= h^2X + \phi^2X.\end{aligned}$$

Proof. We compute $R_{\xi X}\xi = \nabla_\xi\nabla_X\xi - \nabla_X\nabla_\xi\xi - \nabla_{[\xi,X]}\xi$ using $\nabla_X\xi = -\phi X - \phi hX$; thus

$$R_{\xi X}\xi = \nabla_\xi(-\phi X - \phi hX) + \phi[\xi, X] + \phi h[\xi, X].$$

Applying ϕ and recalling that $\nabla_\xi \phi = 0$, we have

$$\begin{aligned}\phi R_{\xi X} \xi &= \nabla_\xi (X + hX) - \eta(\nabla_\xi (X + hX))\xi - [\xi, X] + \eta([\xi, X])\xi - h[\xi, X] \\ &= (\nabla_\xi h)X + \nabla_X \xi + h\nabla_X \xi.\end{aligned}$$

Using $\nabla_X \xi = -\phi X - \phi hX$ and $\phi h + h\phi = 0$ (Lemma 6.2), this becomes

$$\phi R_{\xi X} \xi = (\nabla_\xi h)X - \phi X + h^2 \phi X,$$

which is the first formula.

Now from the first formula we have

$$R_{\xi X} \xi = h^2 X + \phi^2 X - \phi(\nabla_\xi h)X$$

and

$$\phi R_{\xi \phi X} \xi = -h^2 X - \phi^2 X - \phi(\nabla_\xi h)X;$$

subtracting then yields the second formula. ■

Corollary 7.1 *On a contact metric manifold M^{2n+1} the Ricci curvature in the direction ξ is given by*

$$Ric(\xi) = 2n - \text{tr} h^2.$$

Proof. Choosing X to be a unit vector orthogonal to ξ , the inner product of X with the second formula yields the following formula for sectional curvatures:

$$K(\xi, X) + K(\xi, \phi X) = 2(1 - g(h^2 X, X)).$$

Therefore if $\{X_1, \dots, X_n, \phi X_1, \dots, \phi X_n, \xi\}$ is a ϕ -basis, then summing over $\{X_1, \dots, X_n\}$ yields the result. ■

Recall that a K-contact structure is a contact metric structure for which the vector field ξ is Killing and that this is the case if and only if the symmetric operator h vanishes. Thus from the above corollary we have the following immediate result (the author [1977]).

Theorem 7.1 *A contact metric manifold M^{2n+1} is K-contact if and only if $Ric(\xi) = 2n$.*

With regard to sectional curvature we have an earlier result obtained by Hatakeyama, Ogawa, and Tanno [1963].

Theorem 7.2 *A contact metric manifold is K-contact if and only if the sectional curvature of all plane sections containing ξ are equal to 1. Moreover, on a K-contact manifold,*

$$R_X \xi \xi = X - \eta(X)\xi.$$

Proof. In view of the above, the sufficiency is clear. Conversely, if the structure is K-contact, then since $\nabla_X \xi = -\phi X$, we have for X orthogonal to ξ ,

$$R_{\xi X} \xi = \nabla_{\xi}(-\phi X) + \phi[\xi, X] = -\phi \nabla_X \xi = \phi^2 X = -X.$$

The second statement is easily obtained. ■

Furthermore, for the Ricci operator Q acting on ξ we have the following.

Proposition 7.2 *On a K-contact metric manifold M^{2n+1} , $Q\xi = 2n\xi$.*

Proof. Since ξ is Killing, it is affine and therefore

$$\nabla_X \nabla_Y \xi - \nabla_{\nabla_X Y} \xi = R_{X\xi} Y.$$

From this we have $-(\nabla_X \phi)Y = R_{X\xi} Y$, but in Section 6.1 we saw that on a contact metric manifold $\nabla_i \phi^i_j = -2n\eta_j$. Thus letting $\{X_A\}$ be a local orthonormal basis of the contact subbundle we have

$$Q\xi = \sum R_{\xi X_A} X_A = \sum (\nabla_{X_A} \phi) X_A = 2n\xi.$$

■

We noted in Theorem 7.2 that on a K-contact manifold, $R_{X\xi} \xi = X - \eta(X)\xi$. On a Sasakian manifold we have the following stronger result.

Proposition 7.3 *On a Sasakian manifold,*

$$R_{XY} \xi = \eta(Y)X - \eta(X)Y.$$

Proof.

$$\begin{aligned} R_{XY} \xi &= -\nabla_X \phi Y + \nabla_Y \phi X + \phi[X, Y] \\ &= -(\nabla_X \phi)Y + (\nabla_Y \phi)X \\ &= \eta(Y)X - \eta(X)Y. \end{aligned}$$

■

As converses of these results one often sees propositions of the following type (Hatakeyama, Ogawa, and Tanno [1963]).

Proposition 7.4 *Let (M^{2n+1}, g) be a Riemannian manifold admitting a unit Killing field ξ such that $R_X \xi = X$ for X orthogonal to ξ . Then M^{2n+1} is a K-contact manifold.*

Proof. Let $\eta(X) = g(X, \xi)$ and $\phi X = -\nabla_X \xi$. Since ξ is unit Killing, we have $\nabla_\xi \xi = 0$ and

$$\nabla_X \nabla_Y \xi - \nabla_{\nabla_X Y} \xi = R_{X\xi} Y. \quad (*)$$

Thus for X orthogonal to ξ ,

$$\phi^2 X = \nabla_{\nabla_X \xi} \xi = R_{\xi X} \xi = -X$$

and $\phi \xi = 0$. Therefore $\phi^2 = -I + \eta \otimes \xi$. Moreover,

$$d\eta(X, Y) = \frac{1}{2}(g(\nabla_X \xi, Y) - g(\nabla_Y \xi, X)) = -g(\nabla_Y \xi, X) = g(X, \phi Y).$$

Therefore (ϕ, ξ, η, g) is a contact metric structure on M^{2n+1} . ■

An interesting variation is a result of Rukimbira [1995b] that if a Riemannian manifold admits a unit Killing field ξ such that the sectional curvatures of plane sections containing ξ is positive, then it also admits a K-contact structure but with a possibly different metric.

Proposition 7.5 *If in Proposition 7.4, $R_{X\xi} = g(\xi, Y)X - g(X, \xi)Y$, then M^{2n+1} is Sasakian.*

Proof. From equation $(*)$ in the proof of Proposition 7.4, $(\nabla_X \phi)Y = R_{\xi X} Y$ and hence

$$g((\nabla_X \phi)Y, Z) = g(R_{\xi X} Y, Z) = g(R_{Y\xi} X, Z) = g(\eta(Z)Y - \eta(Y)Z, X).$$

■

These results start with a Riemannian structure and construct the desired structure. However, given a contact metric structure we have the following proposition.

Proposition 7.6 *A contact metric structure is Sasakian if and only if*

$$R_{X\xi} = \eta(Y)X - \eta(X)Y.$$

Proof. The necessity is Proposition 7.3 above. To prove the sufficiency first note that for X orthogonal to ξ , $R_\xi X\xi = -X$, and so from the second equation of Proposition 7.1 we have

$$\frac{1}{2}(-X - \phi(-\phi X)) = h^2 X + \phi^2 X = h^2 X - X.$$

Therefore $h^2 = 0$, but h is a symmetric operator and so $h = 0$. Thus ξ is Killing and the result follows as above. ■

Finally, for future use we establish some additional lemmas on Sasakian manifolds. The reader will recognize these curvature properties as being analogous to well-known curvature properties of Kähler manifolds. Let M^{2n+1} be a Sasakian manifold with structure tensors (ϕ, ξ, η, g) and define a tensor field P of type (0,4) by

$$\begin{aligned} P(X, Y, Z, W) &= d\eta(X, Z)g(Y, W) - d\eta(X, W)g(Y, Z) \\ &\quad - d\eta(Y, Z)g(X, W) + d\eta(Y, W)g(X, Z). \end{aligned}$$

Lemma 7.1 *On a Sasakian manifold we have*

$$(a) \quad g(R_{X \ Y} Z, \phi W) + g(R_{X \ Y} \phi Z, W) = -P(X, Y, Z, W).$$

For X, Y, Z, W orthogonal to ξ we have

$$(b) \quad g(R_{\phi X \ \phi Y} \phi Z, \phi W) = g(R_{X \ Y} Z, W)$$

and

$$(c) \quad g(R_{X \ \phi X} Y, \phi Y) = g(R_{X \ Y} X, Y) + g(R_{X \ \phi Y} X, \phi Y) - 2P(X, Y, X, \phi Y).$$

Proof. A direct computation or the Ricci identity shows that

$$\begin{aligned} (\nabla_X \nabla_Y \Phi - \nabla_Y \nabla_X \Phi - \nabla_{[X, Y]} \Phi)(Z, W) \\ = -g(R_{X \ Y} Z, \phi W) - g(R_{X \ Y} \phi Z, W). \end{aligned}$$

Computing the left-hand side using $(\nabla_X \phi)Y = g(X, Y)\xi - \eta(Y)X$ yields (a). Using (a) and the definition of P we obtain (b). Finally, applying the first Bianchi identity to $g(R_{X \ \phi X} Y, \phi Y)$ and using (a), we obtain (c). ■

Lemma 7.2 *On a Sasakian manifold, $Q\phi = \phi Q$.*

Proof. Choosing a ϕ -basis $\{X_i, X_{n+i} = \phi X_i, \xi\}$, we have for X and Y orthogonal to ξ ,

$$\begin{aligned} g(Q\phi X, \phi Y) &= \sum_{A=1}^{2n} g(R_{\phi X \phi X_A} X_A, \phi Y) + g(R_{\phi X \xi} \xi, \phi Y) \\ &= \sum_{A=1}^{2n} g(R_{\phi X \phi X_A} \phi X_A, \phi Y) + g(X, Y) \\ &= g(QX, Y), \end{aligned}$$

where we have used (b) from the previous lemma. We already know that $Q\xi = 2n\xi$, and hence the Ricci operator Q commutes with ϕ on a Sasakian manifold. ■

7.2 Curvature of contact metric manifolds

Before giving our main curvature results, we present some rather complicated lemmas from the paper [1979] of Olszak.

Lemma 7.3 *On a contact metric manifold,*

$$(\nabla_X \phi)Y + (\nabla_{\phi X} \phi)\phi Y = 2g(X, Y)\xi - \eta(Y)(X + hX + \eta(X)\xi).$$

Proof. Using Corollary 6.1 or by direct differentiation of $\nabla_Y \xi = -\phi Y - \phi hY$ we obtain

$$\begin{aligned} (\nabla_X \Phi)(\phi Y, Z) - (\nabla_X \Phi)(Y, \phi Z) &= -\eta(Y)g(X + hX, \phi Z) \\ &\quad - \eta(Z)g(X + hX, \phi Y), \end{aligned} \quad (*)$$

and replacing Z by ϕZ and using Corollary 6.1 again we obtain

$$(\nabla_X \Phi)(\phi Y, \phi Z) + (\nabla_X \Phi)(Y, Z) = \eta(Y)g(X + hX, Z) - \eta(Z)g(X + hX, Y). \quad (**)$$

Now since $d\Phi = 0$, we have

$$\begin{aligned} &(\nabla_X \Phi)(Y, Z) + (\nabla_Y \Phi)(Z, X) + (\nabla_Z \Phi)(X, Y) \\ &+ (\nabla_{\phi X} \Phi)(\phi Y, Z) + (\nabla_{\phi Y} \Phi)(Z, \phi X) + (\nabla_{\phi Z} \Phi)(\phi X, \phi Y) \\ &+ (\nabla_{\phi X} \Phi)(Y, \phi Z) + (\nabla_Y \Phi)(\phi Z, \phi X) + (\nabla_{\phi Z} \Phi)(\phi X, Y) \\ &- (\nabla_X \Phi)(\phi Y, \phi Z) - (\nabla_{\phi Y} \Phi)(\phi Z, X) - (\nabla_{\phi Z} \Phi)(X, \phi Y) = 0. \end{aligned}$$

Now (*) and (**) give

$$\begin{aligned} & (\nabla_{\phi X}\Phi)(Z, \phi Y) + (\nabla_X\Phi)(Z, Y) \\ &= 2\eta(Z)g(X, Y) - \eta(Y)g(X + hX, Z) - \eta(X)\eta(Y)\eta(Z), \end{aligned}$$

from which the result follows. ■

Lemma 7.4 *The curvature tensor of a contact metric manifold satisfies*

$$\begin{aligned} g(R_{\xi X}Y, Z) &= -(\nabla_X\Phi)(Y, Z) - g(X, (\nabla_Y\phi h)Z) + g(X, (\nabla_Z\phi h)Y), \\ g(R_{\xi X}Y, Z) - g(R_{\xi X}\phi Y, \phi Z) &+ g(R_{\xi\phi X}Y, \phi Z) + g(R_{\xi\phi X}\phi Y, Z) \\ &= 2(\nabla_{hX}\Phi)(Y, Z) - 2\eta(Y)g(X + hX, Z) + 2\eta(Z)g(X + hX, Y). \end{aligned}$$

Proof. Differentiating $\nabla_Z\xi = -\phi Z - \phi hZ$, we have

$$R_{YZ}\xi = -(\nabla_Y\phi)Z + (\nabla_Z\phi)Y - (\nabla_Y\phi h)Z + (\nabla_Z\phi h)Y,$$

which, since $d\Phi = 0$, yields the first formula. Now set

$$\begin{aligned} A(X, Y, Z) &= -(\nabla_X\Phi)(Y, Z) + (\nabla_X\Phi)(\phi Y, \phi Z) \\ &\quad - (\nabla_{\phi X}\Phi)(Y, \phi Z) - (\nabla_{\phi X}\Phi)(\phi Y, Z) \end{aligned}$$

and

$$\begin{aligned} B(X, Y, Z) &= -g(X, (\nabla_Y\phi h)Z) + g(X, (\nabla_{\phi Y}\phi h)\phi Z) \\ &\quad - g(\phi X, (\nabla_Y\phi h)\phi Z) - g(\phi X, (\nabla_{\phi Y}\phi h)Z). \end{aligned}$$

Then by the first formula, the left side of the second formula is $A(X, Y, Z) + B(X, Y, Z) - B(X, Z, Y)$. Now by Lemma 7.3 and equation (**) in its proof we have

$$A(X, Y, Z) = 2g(X, Y)\eta(Z) - 2g(X, Z)\eta(Y).$$

Also it is straightforward to show that $\eta((\nabla_{\phi Y}h)Z) = g(-Y + hY, hZ)$. Now rewrite B as

$$\begin{aligned} B(X, Y, Z) &= -g(X, (\nabla_Y\phi)hZ) + g(X, h(\nabla_Y\phi)Z) \\ &\quad + g(X, h\phi(\nabla_{\phi Y}\phi)Z) + g(X, \phi(\nabla_{\phi Y}\phi)hZ) \\ &\quad + \eta(X)\eta((\nabla_{\phi Y}h)Z). \end{aligned}$$

Then using Lemma 7.3 again,

$$B(X, Y, Z) = 2g(hX, (\nabla_Y \phi)Z) + 2\eta(Z)g(hX, Y) - 2\eta(X)g(hY, hZ).$$

Finally, computing $A(X, Y, Z) + B(X, Y, Z) - B(X, Z, Y)$ and using $d\Phi = 0$, we have the result. ■

Let ρ denote the Ricci tensor and τ the scalar curvature. In addition, we define the **-Ricci tensor* ρ^* and **-scalar curvature* τ^* by contracting the curvature tensor by ϕ instead of the metric. Precisely,

$$\rho_{ij}^* = R_{iklm}\phi^{kl}\phi_j^m, \quad \tau^* = \rho^*{}^i{}_i.$$

These notions have their origin in almost Hermitian geometry, and we shall see their almost Hermitian analogues in Section 10.2. We now have an important proposition due to Olszak [1979].

Proposition 7.7 *On a contact metric manifold M^{2n+1} ,*

$$\tau^* - \tau + 4n^2 = \text{tr}h^2 + \frac{1}{2}(|\nabla\phi|^2 - 4n) \geq 0,$$

with equality if and only if M^{2n+1} is Sasakian.

Proof. We have seen that $\nabla_i\phi^i{}_j = -2n\eta_j$ and $\nabla_k\xi^j = -\phi^j{}_k - \phi^j{}_m h^m{}_k$. Therefore using $\phi^2 = -I + \eta \otimes \xi$ and basic properties of h (Section 6.4), we have

$$\phi^{kj}\nabla_k\nabla_i\phi^i{}_j = -4n^2. \quad (*)$$

Differentiation of $\phi^2 = -I + \eta \otimes \xi$ yields $\phi^{kj}\nabla_t\phi_{kj} = 0$. Since $d\Phi = 0$, we have $\phi^{kj}\nabla_k\phi_{tj} = \phi^{kj}(-\nabla_t\phi_{jk} - \nabla_j\phi_{kt})$ from which we obtain $\phi^{kj}\nabla_k\phi^t{}_j = 0$. In turn, $\phi^{kj}\nabla_t\nabla_k\phi^t{}_j = -(\nabla_t\phi^{kj})(\nabla_k\phi^t{}_j)$. Using $d\Phi = 0$ on the second factor on the right and simplifying, we have

$$\phi^{kj}\nabla_t\nabla_k\phi^t{}_j = -\frac{1}{2}|\nabla\phi|^2. \quad (**)$$

From (*) and (**),

$$\phi^{kj}(\nabla_k\nabla_t\phi^t{}_j - \nabla_t\nabla_k\phi^t{}_j) = \frac{1}{2}|\nabla\phi|^2 - 4n^2.$$

Therefore by Corollary 7.1,

$$\begin{aligned} \frac{1}{2}|\nabla\phi|^2 - 4n^2 &= \phi^{kj}(R_{kta}{}^t\phi^a{}_j - R_{ktj}{}^a\phi^t{}_a) = (g^{ka} - \xi^k\xi^a)(-\rho_{ka}) + \tau^* \\ &= \tau^* - \tau + (2n - \text{tr}h^2), \end{aligned}$$

which is the desired formula.

Now $(\nabla_i \phi_{jk} - g_{ik} \eta_j + g_{ij} \eta_k)(\nabla^i \phi^{jk} - g^{ik} \xi^j + g^{ij} \xi^k) \geq 0$ is equivalent to $|\nabla \phi|^2 - 4n \geq 0$, giving the inequality, and by Theorem 6.3, equality holds if and only if the structure is Sasakian. ■

We now prove the following important result of Olszak [1979].

Theorem 7.3 *If a contact metric manifold M^{2n+1} is of constant curvature c and dimension ≥ 5 , then $c = 1$ and the structure is Sasakian.*

Proof. Recall from Proposition 7.1 that $\frac{1}{2}(R_{\xi X} \xi - \phi R_{\xi \phi X} \xi) = h^2 X + \phi^2 X$; thus if $R_{XY}Z = c(g(Y, Z)X - g(X, Z)Y)$, then $\frac{c}{2}(\eta(X)\xi - X + \phi^2 X) = h^2 X + \phi^2 X$. Therefore $h^2 X + (c - 1)\phi^2 X$ and hence $\text{tr} h^2 = 2n(1 - c)$. Now from the second equation in Lemma 7.4,

$$\begin{aligned} c(g(X, Y)\eta(Z) - \eta(Y)g(X, Z)) - 0 - c\eta(Y)g(\phi X, \phi Z) + c\eta(Z)g(\phi X, \phi Y) \\ = 2(\nabla_{hX}\Phi)(Y, Z) - 2\eta(Y)g(X + hX, Z) + 2\eta(Z)g(X + hX, Y). \end{aligned}$$

Therefore

$$\begin{aligned} (\nabla_{hX}\Phi)(Y, Z) &= (1 - c)(\eta(Y)g(X, Z) - \eta(Z)g(X, Y)) \\ &\quad + \eta(Y)g(hX, Z) - \eta(Z)g(hX, Y). \end{aligned}$$

Replacing X by hX , we have

$$\begin{aligned} -g((\nabla_{(c-1)(-X+\eta(X)\xi)}\phi)Y, Z) &= (1 - c)(\eta(Y)g(hX, Z) - \eta(Z)g(hX, Y)) \\ &\quad + \eta(Y)g((c - 1)(-X + \eta(X)\xi), Z) \\ &\quad - \eta(Z)g((c - 1)(-X + \eta(X)\xi), Y), \end{aligned}$$

and hence

$$(\nabla_X \phi)Y = g(X + hX, Y)\xi - \eta(Y)(X + hX).$$

Using this to compute $|\nabla \phi|^2$ and using $\text{tr} h^2 = 2n(1 - c)$, we obtain

$$|\nabla \phi|^2 = 4n(2 - c).$$

On the other hand, $\tau = 2n(2n + 1)c$, and $\tau^* = 2nc$, as is easily checked. Now from Olszak's formula in Proposition 7.7,

$$\tau - \tau^* - 4n^2 = -\text{tr} h^2 - \frac{1}{2}|\nabla \phi|^2 + 2n \leq 0,$$

with equality if and only if the structure is Sasakian. Computing both sides of this formula, we find that $4n^2(c-1) = 4n(c-1)$; thus if $n > 1$, then $c = 1$ and hence $\tau - \tau^* - 4n^2 = 0$, giving M^{2n+1} Sasakian. ■

Earlier, the present author [1976] showed that in dimension ≥ 5 , there are no flat associated metrics. Thus while the 5-torus carries a contact structure (Example 3.2.6), the flat metric is not an associated metric. In dimension 3, the only constant curvature, cases are constant curvature 0 and 1, as we will note below. The nonexistence of flat associated metrics does raise the question whether there are contact metric manifolds of everywhere non-positive curvature, except for the flat 3-dimensional case. If the manifold is compact and we ask for strictly negative curvature, we can answer this question in the negative using the following deep result of A. Zeghib [1995] on geodesic plane fields. Recall that a plane field on a Riemannian manifold is said to be *geodesic* if any geodesic tangent to the plane field at one point is tangent to it at every point.

Theorem 7.4 *A compact negatively curved Riemannian manifold has no C^1 geodesic plane field (of nontrivial dimension).*

Since for any contact metric structure the integral curves of ξ are geodesics (Theorem 4.5), ξ determines a geodesic line field to which we can apply the theorem of Zeghib, as was pointed out by Rukimbira [1998]. Thus we have the following corollary.

Corollary 7.2 *On a compact contact manifold, there is no associated metric of strictly negative curvature.*

The author conjectures that this and a bit more is true locally, viz. that except for the flat 3-dimensional case, any contact metric manifold has some positive sectional curvature. The fact that hyperbolic space has many 1-dimensional totally geodesic foliations does not violate such a conjecture, since the hyperbolic metric cannot be an associated metric of any contact structure by the above theorem of Olszak.

Along the line of the influence of a contact structure on the curvature of its associated metrics we also make the following remark. In [1941] Myers proved that a complete Riemannian manifold for which $Ric \geq \delta > 0$ is compact and has finite fundamental group. In [1981] Hasegawa and Seino generalized Myers' theorem for a K-contact manifold by proving that a complete K-contact manifold for which $Ric \geq -\delta > -2$ is compact (they state their result in the Sasakian case, but their proof uses only the

K-contact property). As we have seen in the K-contact case, all sectional curvatures of plane sections containing ξ are equal to 1, and hence there is some of positive curvature from the outset. In an attempt to weaken the K-contact requirement in this result, R. Sharma and the author [1990a] considered a contact metric manifold M^{2n+1} for which ξ is an eigenvector field of the Ricci operator, equivalently, the divergence of $h\phi$ is proportional to η . In this case, if $Ric \geq -\delta > -2$ and the sectional curvatures of plane sections containing ξ are $\geq \epsilon > \delta' \geq 0$, where

$$\delta' = 2\sqrt{n(\delta - 2\sqrt{2\delta} + n + 2) - (\delta - 2\sqrt{2\delta} + 1 + 2n)},$$

then M^{2n+1} is compact.

The flat case was investigated further by Rukimbira [1998], who showed that a compact flat contact metric manifold is isometric to the quotient of a flat 3-torus by a finite cyclic group of isometries of order 1, 2, 3, 4 or 6.

In Example 3.2.6 and Section 6.2 we saw explicitly a flat contact metric structure on \mathbb{R}^3 and in turn on the 3-dimensional torus T^3 . As in Section 6.2, let $\eta = \frac{1}{2}(\cos x^3 dx^1 + \sin x^3 dx^2)$ be the contact form; $g_{ij} = \frac{1}{4}\delta_{ij}$ is an associated metric, and the nonzero eigenvalues of h are ± 1 . It is interesting to study this example in Darboux coordinates and to generalize it to higher dimensions. By the Darboux theorem there exist coordinates (x, y, z) such that the contact form η is given by $\frac{1}{2}(dz - y dx)$. Consider the map $f: \mathbb{R}^3 \rightarrow \mathbb{R}^3$ given by

$$x^1 = z \cos x - y \sin x, \quad x^2 = -z \sin x - y \cos x, \quad x^3 = -x.$$

Then $\frac{1}{2}(dz - y dx) = f^*\eta$ and $g_0 = f^*g$ is a flat associated metric for the Darboux form $\eta_0 = \frac{1}{2}(dz - y dx)$. The metric g_0 is given by

$$g_0 = \frac{1}{4} \begin{pmatrix} 1 + y^2 + z^2 & z & -y \\ z & 1 & 0 \\ -y & 0 & 1 \end{pmatrix}.$$

Now consider the Darboux form $\eta = \frac{1}{2}(dz - \sum y^i dx^i)$ on \mathbb{R}^{2n+1} . The metric

$$g = \frac{1}{4} \begin{pmatrix} \delta_{ij} + y^i y^j + \delta_{ij} z^2 & \delta_{ij} z & -y^i \\ \delta_{ij} z & \delta_{ij} & 0 \\ -y^j & 0 & 1 \end{pmatrix}$$

is an associated metric quite different from the Sasakian metric given in Examples 4.5.1 and 6.7.1. Direct computation shows that $h \frac{\partial}{\partial y^i} = -\frac{\partial}{\partial y^i}$. Therefore -1 is an eigenvalue of h of multiplicity n , and hence in turn $+1$ is also an eigenvalue of multiplicity n . This metric enjoys the following curvature properties: $R_{\xi X}\xi = 0$ for every X and $R_{XY}\xi = 0$ for $X, Y \in \{-1\}$. However, $R_{XY}\xi \neq 0$ in general; for example in dimension 5,

$$R_{\frac{\partial}{\partial x^1} \frac{\partial}{\partial x^2}}\xi = \frac{1}{2} \left(-y^2 \frac{\partial}{\partial x^1} + y^1 \frac{\partial}{\partial x^2} + y^2 z \frac{\partial}{\partial y^1} - y^1 z \frac{\partial}{\partial y^2} \right).$$

We now show that the condition $R_{XY}\xi = 0$ for all X, Y has a strong and interesting implication for a contact metric manifold, namely that it is locally the product of Euclidean space E^{n+1} and a sphere of constant curvature $+4$ (the author [1977]).

Theorem 7.5 *A contact metric manifold M^{2n+1} satisfying $R_{XY}\xi = 0$ is locally isometric to $E^{n+1} \times S^n(4)$ for $n > 1$ and flat for $n = 1$.*

Proof. $R_{XY}\xi = 0$ implies, by Proposition 7.1, that $h^2 + \phi^2 = 0$ and hence that the nonzero eigenvalues of h are ± 1 , each with multiplicity n . For $X, Y \in [-1]$,

$$0 = -\nabla_{[X,Y]}\xi = \phi[X, Y] + \phi h[X, Y]$$

and $\eta([X, Y]) = -2d\eta(X, Y) = -2g(X, \phi Y) = 0$. Thus $[-1]$ is integrable. Also $0 = -\nabla_{[X,\xi]}\xi$, so that $[X, \xi] \in [-1]$. Therefore $[-1] \oplus \{\xi\}$ is integrable.

Choose coordinates (u^0, \dots, u^{2n}) such that $\frac{\partial}{\partial u^0}, \dots, \frac{\partial}{\partial u^n} \in [-1] \oplus \{\xi\}$. Let $X_i = \frac{\partial}{\partial u^{n+i}} + \sum_{j=0}^n f_i^j \frac{\partial}{\partial u^j}$, where the f_i^j 's are functions chosen so that $X_i \in [+1]$. Now $[\frac{\partial}{\partial u^k}, X_i] \in [-1] \oplus \{\xi\}$, $k = 0, \dots, n$, and hence

$$0 = \nabla_{[\frac{\partial}{\partial u^k}, X_i]}\xi = \nabla_{\frac{\partial}{\partial u^k}}\nabla_{X_i}\xi - \nabla_{X_i}\nabla_{\frac{\partial}{\partial u^k}}\xi = -2\nabla_{\frac{\partial}{\partial u^k}}\phi X_i,$$

from which

$$\nabla_{\phi X_j}\phi X_i = 0.$$

We therefore see that the integral submanifolds of $[-1] \oplus \{\xi\}$ are totally geodesic and flat.

From the second equation of Lemma 7.4, we have for X, Y, Z orthogonal to ξ that $g((\nabla_X \phi)Y, Z) = 0$. Also for $X, Y \in [+1]$,

$$\begin{aligned} 0 &= R_{XY}\xi = -2\nabla_X \phi Y + 2\nabla_Y \phi X - \nabla_{[X,Y]}\xi \\ &= -2(\nabla_X \phi)Y + 2(\nabla_Y \phi)X - 2\phi[X, Y] + \phi[X, Y] + \phi h[X, Y]. \end{aligned}$$

Taking the inner product with $Z \in [+1]$ gives $g(-\phi[X, Y] - h\phi[X, Y], Z) = 0$, and hence $g(\phi[X, Y], Z) = 0$. Thus $[X, Y]$ is orthogonal to $[-1]$. Also $\eta([X, Y]) = 0$, and therefore $[+1]$ is integrable.

Now take $X \in [-1]$ and $Y \in [+1]$. Then

$$\begin{aligned} 0 &= R_{XY}\xi = -2\nabla_X\phi Y + \phi[X, Y] + \phi h[X, Y] \\ &= -2(\nabla_X\phi)Y - \phi\nabla_XY - \phi\nabla_YX - h\phi\nabla_XY + h\phi\nabla_YX. \end{aligned}$$

Taking the inner product with $Z \in [-1]$, we have $g(\phi\nabla_YX, Z) = 0$ and hence that ∇_YX is orthogonal to $[-1]$. Also $\nabla_Y\xi = -2\phi Y$ is orthogonal to $[-1]$. Therefore the integral submanifolds of $[+1]$ are totally geodesic. Moreover, we are now at the point that M^{2n+1} has a local Riemannian product structure.

For $X \in [+1]$,

$$\begin{aligned} g((\nabla_X\phi)Y, \xi) &= -g((\nabla_X\phi)\xi, Y) = g(\phi\nabla_X\xi, Y) \\ &= g(\phi(-2\phi X), Y) = 2g(X, Y), \end{aligned}$$

and hence by the second formula of Lemma 7.4, $(\nabla_X\phi)Y = 2g(X, Y)\xi$ for $X, Y \in [+1]$. Now for $X, Y, Z, W \in [+1]$,

$$\begin{aligned} &g(\nabla_X\nabla_Y\phi Z, \phi W) - g(\nabla_X\nabla_YZ, W) \\ &= g(\nabla_X(2g(Y, Z)\xi + \phi\nabla_YZ), \phi W) - g(\nabla_X\nabla_YZ, W) \\ &= -4g(Y, Z)g(X, W), \end{aligned}$$

using the property we noted above that $g((\nabla_X\phi)Y, Z) = 0$ for X, Y, Z orthogonal to ξ . Using this property again to treat the terms involving $\nabla_{[X, Y]}$, we have

$$g(R_{XY}\phi Z, \phi W) - g(R_{XY}Z, W) = -4(g(Y, Z)g(X, W) - g(X, Z)g(Y, W))$$

but $g(R_{XY}\phi Z, \phi W) = 0$ by virtue of the Riemannian product structure. Therefore the integral submanifolds of $[+1]$ are locally isometric to $S^n(4)$.

In dimension 3, the integrability of $[-1]$ and $[+1]$ is immediate and the rest of the proof goes through, giving that M^3 is flat. ■

7.3 The (κ, μ) -manifolds

As a generalization of both $R_{XY}\xi = 0$ (Theorem 7.5 above) and the Sasakian case, $R_{XY}\xi = \eta(Y)X - \eta(X)Y$, consider

$$R_{XY}\xi = \kappa(\eta(Y)X - \eta(X)Y) + \mu(\eta(Y)hX - \eta(X)hY)$$

for constants κ and μ . A contact metric manifold satisfying this condition is called a (κ, μ) -manifold. Despite the technical appearance of this condition there are good reasons for considering it; we first mention them here, referring to Koufogiorgos and Papantoniou and the author [1995] for details and then give a classification theorem due to Boeckx [2000] and other results.

Theorem 7.6 *A (κ, μ) -manifold M is a strongly pseudoconvex CR-manifold.*

Theorem 7.7 *Let M^{2n+1} be a (κ, μ) -manifold. Then $\kappa \leq 1$. If $\kappa = 1$, the structure is Sasakian. If $\kappa < 1$, the (κ, μ) condition determines the curvature of M^{2n+1} completely.*

We remark that when $\kappa = 1$, the proof shows that $h = 0$, and hence μ is indeterminate in this case. When $\kappa < 1$, the nonzero eigenvalues of h are $\pm\sqrt{1-\kappa}$, each with multiplicity n . Let λ be the positive eigenvalue. Then M^{2n+1} admits three mutually orthogonal subbundles, $\mathcal{D}(0)$, $\mathcal{D}(\lambda)$ and $\mathcal{D}(-\lambda)$, which are integrable.

The curvature tensor of a non-Sasakian (κ, μ) -manifold may be found in Boeckx [1999]. We mention only that for a unit vector $X \in [\lambda]$, $\phi X \in [-\lambda]$, we have the following sectional curvatures:

$$K(X, \xi) = \kappa + \lambda\mu, \quad K(\phi X, \xi) = \kappa - \lambda\mu, \quad K(X, \phi X) = -(\kappa + \mu).$$

Thus, turning to the question of the sign of the curvature, if a (κ, μ) -manifold were of negative curvature, then $\lambda > 1$ by Corollary 7.1 and now with both $\kappa \pm \lambda\mu < 0$, we have $\lambda^2 - 1 > \lambda|\mu|$. Then $K(X, \phi X) = -(\kappa + \mu) < 0$ which gives $\lambda|\mu| < \lambda^2 - 1 < \mu$, a contradiction.

The Ricci operator and the scalar curvature of a $(2n+1)$ -dimensional (κ, μ) -manifold are given by

$$\begin{aligned} QX &= (2(n-1) - n\mu)X + (2(n-1) + \mu)hX \\ &\quad + (n(2\kappa + \mu) - 2(n-1))\eta(X)\xi, \\ \tau &= 2n(2(n-1) + \kappa - n\mu). \end{aligned}$$

Theorem 7.8 *Let M be a 3-dimensional (κ, μ) -manifold. Then M is either Sasakian or locally isometric to one of the unimodular Lie groups $SU(2)$, $SL(2, \mathbb{R})$, $E(2)$, $E(1, 1)$ with a left-invariant metric.*

Theorem 7.9 *The standard contact metric structure on the tangent sphere bundle T_1M (see Section 9.2) satisfies the (κ, μ) condition if and only if the base manifold M is of constant curvature. In particular, if M has constant curvature c , then $\kappa = c(2 - c)$ and $\mu = -2c$.*

Given a contact metric structure (ϕ, ξ, η, g) , consider the deformed structure

$$\bar{\eta} = a\eta, \quad \bar{\xi} = \frac{1}{a}\xi, \quad \bar{\phi} = \phi, \quad \bar{g} = ag + a(a-1)\eta \otimes \eta,$$

where a is a positive constant. Such a deformation is called a \mathcal{D} -homothetic deformation, since the metrics restricted to the contact subbundle \mathcal{D} are homothetic. This deformation was introduced by Tanno in [1968] and has many applications. While such a change preserves the state of being contact metric, K-contact, Sasakian, or strongly pseudoconvex CR, it destroys a condition like $R_{XY}\xi = 0$ or $R_{XY}\xi = \kappa(\eta(Y)X - \eta(X)Y)$. However, the form of the (κ, μ) condition is preserved under a \mathcal{D} -homothetic deformation with

$$\bar{\kappa} = \frac{\kappa + a^2 - 1}{a^2}, \quad \bar{\mu} = \frac{\mu + 2a - 2}{a}.$$

For a non-Sasakian (κ, μ) -manifold M , Boeckx [2000] introduced an invariant

$$I_M = \frac{1 - \frac{\mu}{2}}{\sqrt{1 - \kappa}}$$

and showed that for two non-Sasakian (κ, μ) -manifolds $(M_i, \phi_i, \xi_i, \eta_i, g_i)$, $i = 1, 2$, we have $I_{M_1} = I_{M_2}$ if and only if up to a \mathcal{D} -homothetic deformation, the two spaces are locally isometric as contact metric manifolds. Thus we know all non-Sasakian (κ, μ) -manifolds locally as soon as we have for every odd dimension $2n + 1$ and for every possible value of the invariant I , one (κ, μ) -manifold (M, ϕ, ξ, η, g) with $I_M = I$. From Theorem 7.9 we see that for the standard contact metric structure on the tangent sphere bundle of a manifold of constant curvature c , $I = \frac{1+c}{|1-c|}$. Therefore as c varies over the reals, I takes on every value > -1 . Boeckx now gives an example for any odd dimension and value of $I \leq -1$; his construction is as follows.

Let \mathfrak{g} be a $(2n + 1)$ -dimensional Lie algebra, $n \geq 2$. Introduce a basis for \mathfrak{g} , $\{\xi, X_1, \dots, X_n, Y_1, \dots, Y_n\}$, and for real numbers α and β define

the Lie bracket by

$$\begin{aligned}
[\xi, X_1] &= -\frac{\alpha\beta}{2}X_2 - \frac{\alpha^2}{2}Y_1, \quad [\xi, X_2] = \frac{\alpha\beta}{2}X_1 - \frac{\alpha^2}{2}Y_2, \\
[\xi, X_i] &= -\frac{\alpha^2}{2}Y_i, \quad i \geq 3, \\
[\xi, Y_1] &= \frac{\beta^2}{2}X_1 - \frac{\alpha\beta}{2}Y_2, \quad [\xi, Y_2] = \frac{\beta^2}{2}X_2 + \frac{\alpha\beta}{2}Y_1, \\
[\xi, Y_i] &= \frac{\beta^2}{2}X_i, \quad i \geq 3, \\
[X_1, X_i] &= \alpha X_i, \quad i \neq 1, \quad [X_i, X_j] = 0, \quad i, j \neq 1, \\
[Y_2, Y_i] &= \beta Y_i, \quad i \neq 2, \quad [Y_i, Y_j] = 0, \quad i, j \neq 2, \\
[X_1, Y_1] &= -\beta X_2 + 2\xi, \quad [X_1, Y_i] = 0, \quad i \geq 2, \\
[X_2, Y_1] &= \beta X_1 - \alpha Y_2, \quad [X_2, Y_2] = \alpha Y_1 + 2\xi, \quad [X_2, Y_i] = \beta X_i, \quad i \geq 3, \\
[X_i, Y_1] &= -\alpha Y_i, \quad i \geq 3, \quad [X_i, Y_2] = 0, \quad i \geq 3, \\
[X_i, Y_j] &= \delta_{ij}(-\beta X_2 + \alpha Y_1 + 2\xi), \quad i, j \geq 3.
\end{aligned}$$

The associated Lie group G is not unimodular for $\dim \mathfrak{g} \geq 5$, and not both α and β are equal to zero, since $\text{tr ad}_{X_1} = (n-1)\alpha$ and $\text{tr ad}_{Y_2} = (n-1)\beta$. Now define a metric on G by left translation of the basis $\{\xi, X_1, \dots, X_n, Y_1, \dots, Y_n\}$, taken as orthonormal at the identity. Then taking η as the metric dual of ξ and defining ϕ by $\phi\xi = 0$, $\phi X_i = Y_i$ and $\phi Y_i = -X_i$, we have a contact metric structure on G . Now for the present purpose suppose that $\beta^2 > \alpha^2$. Then G is a non-Sasakian (κ, μ) -manifold and

$$I_G = -\frac{\beta^2 + \alpha^2}{\beta^2 - \alpha^2} \leq -1;$$

thus for appropriate choices of $\beta > \alpha \geq 0$, I_G attains any value ≤ -1 .

For the 3-dimensional case, keeping Theorem 7.8 in mind, consider the Lie algebra

$$[\xi, X] = -\frac{\alpha^2}{2}Y, \quad [\xi, Y] = \frac{\beta^2}{2}X, \quad [X, Y] = 2\xi,$$

which corresponds to a unimodular Lie group. Boeckx [2000] now points out that for $\beta > \alpha > 0$ we obtain a left-invariant contact metric structure

on $SL(2, \mathbb{R})$ with $I_{SL(2, \mathbb{R})} < -1$. For $\beta > 0$ and $\alpha = 0$ we have a left-invariant contact metric structure on $E(1, 1)$ with $I_{E(1, 1)} = -1$. It is worth mentioning that there are also (κ, μ) -structures on $SL(2, \mathbb{R})$ with $-1 < I_{SL(2, \mathbb{R})} < +1$, on $E(2)$ with $I_{E(2)} = +1$ and on $SU(2)$ with $I_{SU(2)} > 1$. (see also Koufogiorgos, Papantoniou and the author [1995]).

In [2000] Koufogiorgos and Tsihlias considered the question of contact metric manifolds for which ξ satisfies the (κ, μ) condition but where κ and μ are functions rather than constants and called these spaces *generalized (κ, μ) -manifolds*. They showed that in dimensions ≥ 5 , κ and μ must be constant and in dimension 3 gave an example for which κ and μ are not constants; this case is studied further in Koufogiorgos and Tsihlias [2003]. Moreover, this idea is closely related to the question of the characteristic vector field as a map into the tangent sphere bundle being a harmonic map. We discuss this topic in Subsection 10.3.1.

We have already noted that for a non-Sasakian (κ, μ) -manifold, h admits two nonzero eigenvalues $\pm\lambda$ determining n -dimensional integrable subbundles $\mathcal{D}(\lambda)$ and $\mathcal{D}(-\lambda)$ and hence two complementary Legendre foliations. Cappelletti Montano and Di Terlizzi [2008] studied these foliations in detail. The Pang invariants (see Example 5.3.4) are

$$\begin{aligned}\Pi_{\mathcal{D}(\lambda)} &= \frac{(\lambda + 1)^2 - \kappa - \mu\lambda}{\lambda} g|_{\mathcal{D}(\lambda) \times \mathcal{D}(\lambda)}, \\ \Pi_{\mathcal{D}(-\lambda)} &= \frac{-(\lambda - 1)^2 + \kappa - \mu\lambda}{\lambda} g|_{\mathcal{D}(-\lambda) \times \mathcal{D}(-\lambda)}.\end{aligned}$$

Cappelletti Montano and Di Terlizzi then prove the following theorem.

Theorem 7.10 *Let (M, ϕ, ξ, η, g) be a non-Sasakian contact metric manifold. Then the manifold is a (κ, μ) -manifold if and only if it admits two mutually orthogonal Legendre subbundles L and Q and a unique linear connection $\bar{\nabla}$ such that*

$$\bar{\nabla}L \subset L, \quad \bar{\nabla}Q \subset Q,$$

$$\bar{\nabla}\eta = 0, \quad \bar{\nabla}d\eta = 0, \quad \bar{\nabla}g = 0, \quad \bar{\nabla}\phi = 0, \quad \bar{\nabla}h = 0,$$

$$\bar{T}(X, Y) = 2d\eta(X, Y)\xi \text{ for all } X, Y \in \mathcal{D},$$

$$\bar{T}(X, \xi) = [\xi, X_L]_Q + [\xi, X_Q]_L \text{ for all tangent vector fields } X,$$

where \bar{T} denotes the torsion of $\bar{\nabla}$ and the subscripts indicate the projections to L and Q . Furthermore, L and Q are integrable and coincide with the eigenspaces $\mathcal{D}(\lambda)$ and $\mathcal{D}(-\lambda)$.

For a Legendre foliation with $\Pi_{\mathcal{F}}$ nondegenerate, Libermann [1991] introduced a linear map $\Lambda : TM \longrightarrow T\mathcal{F}$ by

$$\Pi_{\mathcal{F}}(\Lambda X, Y) = d\eta(X, Y).$$

While the kernel of this map is $T\mathcal{F} \oplus \mathbb{R}\xi$ and $\Lambda^2 = 0$, this map and $\Pi_{\mathcal{F}}$ can be used in particular cases to define a field of endomorphisms ϕ and a metric g to give a contact metric structure to a contact manifold as in the works of Jayne [1998] and Cappelletti Montano [to appear]. The latter author especially gives conditions under which a contact manifold admits a (κ, μ) -structure.

If on a (κ, μ) -manifold, $\mu = 0$, the contact metric manifold is said to be one for which ξ belongs to the κ -nullity distribution. In general, the κ -nullity distribution of a Riemannian manifold (M, g) is the subbundle $N(k)$ defined by

$$N_p(\kappa) = \{Z \in T_p M \mid R_{XY}Z = \kappa((g(Y, Z)X - g(X, Z)Y) \forall X, Y \in T_p M\}.$$

In dimension 3, Koufogiorgos, Sharma and the author [1990] showed that ξ belonging to $N(\kappa)$ is equivalent to the Ricci operator Q commuting with ϕ and equivalent to the contact metric manifold being η -Einstein, i.e.,

$$Q = aI + b\eta \otimes \xi$$

for some functions a and b . In dimensions ≥ 5 it is known that for any η -Einstein K-contact manifold, a and b are constants. The main result of the author's papers with Koufogiorgos and Sharma [1990] and H. Chen [1992] is that a 3-dimensional contact metric manifold for which ξ belongs to the κ -nullity distribution is either Sasakian, flat, or locally isometric to a left-invariant metric on the Lie group $SU(2)$ or $SL(2, \mathbb{R})$.

We now give another example of a manifold with ξ belonging to the κ -nullity distribution. Using the Boeckx invariant we construct an example with $\kappa = 1 - \frac{1}{n}$, $n > 1$. Since the Boeckx invariant for a $(1 - \frac{1}{n}, 0)$ -manifold is $\sqrt{n} > -1$, referring to Theorem 7.9 above, we consider the tangent sphere bundle of an $(n+1)$ -dimensional manifold of constant curvature c . Choose a \mathcal{D} -homothetic deformation so that the resulting manifold will be a $(1 - \frac{1}{n}, 0)$ -manifold. That is, for $\kappa = c(2 - c)$ and $\mu = -2c$, we solve

$$1 - \frac{1}{n} = \frac{\kappa + a^2 - 1}{a^2}, \quad 0 = \frac{\mu + 2a - 2}{a}$$

for a and c . The result is

$$c = \frac{(\sqrt{n} \pm 1)^2}{n-1}, \quad a = 1 + c,$$

and taking c and a to be these values, we have a contact metric manifold with ξ belonging to the $(1 - \frac{1}{n})$ -nullity distribution.

Now consider the concircular curvature tensor

$$Z_{XY}V = R_{XY}V - \frac{\tau}{2n(2n+1)}(g(Y, V)X - g(X, V)Y)$$

and let $Z_{\xi X} \cdot Z$ denote the action of $Z_{\xi X}$ on Z as a derivation. J.-S. Kim, M. M. Tripathi and the author [2005] proved the following theorem.

Theorem 7.11 *A $(2n+1)$ -dimensional contact metric manifold M with ξ belonging to the κ -nullity distribution satisfies*

$$Z_{\xi X} \cdot Z = 0$$

if and only if M is 3-dimensional and flat, or locally isometric to the sphere $S^{2n+1}(1)$, or locally isometric to the above example $(\kappa = 1 - \frac{1}{n}, n > 1)$.

An interesting side question associated with ξ belonging to the κ -nullity distribution is the dimension of the κ -nullity distribution itself. Clearly if a vector Z belongs to $N(\kappa)$, then the sectional curvatures of all plane sections containing Z are equal to κ . In particular, on any Riemannian manifold, $N(\kappa)$ is nontrivial for at most one value of κ . It is known that $N(\kappa)$ is an integrable subbundle with totally geodesic leaves of constant curvature κ ; see, e.g., Tanno [1978a]. In unpublished work, Baikoussis, Koufogiorgos and the author showed that if the characteristic vector field ξ on a contact metric manifold M^{2n+1} with $n > 1$ belongs to $N(\kappa)$, then $(\kappa \leq 1)$ and if $\kappa < 1$ and $\kappa \neq 0$, then $\dim N(\kappa) = 1$. The corresponding result for $n = 1$ is due to R. Sharma [1995]. If $\kappa = 0$, M^{2n+1} is locally $E^{n+1} \times S^n(4)$, as we have seen, and ξ is tangent to the Euclidean factor, giving $\dim N(0) = n + 1$. If $\kappa = 1$, the structure is Sasakian. This leaves the question of the dimension of $N(1)$ on a Sasakian manifold. P. Rukimbira [2009] showed that the dimension of $N(1)$ is either $\leq n$ or $2n + 1$, i.e., $N(1)$ is the whole tangent bundle. F. Gouli-Andreou and the author conjecture that the dimension of $N(1)$ must be either $2n + 1$ (and M^{2n+1} is of constant curvature) or 3 (and M^{2n+1} has a Sasakian 3-structure; see Chapter 14) or 1.

7.4 Sasakian Einstein manifolds

Sasakian Einstein manifolds have recently been shown to be very numerous, and so a brief discussion is in order. For a more complete treatment see the book of Boyer and Galicki [2008, Chapter 11] and for very recent results see the papers of J. Kollár [2009], J. Sparks [2009], and C. van Coevering [2009].

First note that on a Sasakian Einstein manifold M , $Q\xi = 2n\xi$ and $Q = \frac{\tau}{2n+1}I$, and hence the scalar curvature is $\tau = 2n(2n+1) > 0$. The Einstein constant being $2n$ implies that in the compact case, $\pi_1(M)$ is finite by Myers' theorem [1941].

An important recent result is the following theorem of Boyer and Galicki [2001]; see also Boyer and Galicki [2008, pp. 372–374]. This remarkable result should be compared with the Goldberg conjecture that a compact almost Kähler Einstein space is Kähler (see Section 10.2). The Goldberg conjecture is true for nonnegative scalar curvature (Sekigawa [1987]). Since Sasakian Einstein manifolds have positive scalar curvature, one might hope for a result similar to the Goldberg conjecture in contact geometry, and Boyer and Galicki achieved the following.

Theorem 7.12 *A compact K-contact Einstein manifold M^{2n+1} is Sasakian.*

For an almost regular (quasiregular) K-contact manifold the proof of this theorem by Boyer and Galicki is to consider the quotient by the flow of ξ , which is an almost Kähler orbifold and is Einstein with positive scalar curvature. Since Sekigawa's proof of the Goldberg conjecture uses only local computations and certain integral formulas, one has that the orbifold is Kähler Einstein. Thus in turn, one has that the given manifold is Sasakian. If ξ is not almost regular, Boyer and Galicki show that it may be approximated by a sequence of almost regular vector fields that define K-contact structures for a sequence of metrics.

An alternative proof of this theorem was given by Apostolov, Draghici and Moroianu [2006]. Their approach is to use the cone

$$C(M^{2n+1}) = (\mathbb{R}_+ \times M^{2n+1}, \bar{g} = dr^2 + r^2g)$$

over the K-contact manifold M^{2n+1} . The curvature \bar{R} of the cone is given by $\bar{R}_{XY}Z = R_{XY}Z + g(X, Z)Y - g(Y, Z)X$ for vectors tangent to M^{2n+1} and vanishes if any of the vectors is $\frac{\partial}{\partial r}$. Then $\bar{Q}X = QX - 2nX$, and

we see that g is Einstein (with Einstein constant $2n$) if and only if the cone is Ricci flat. Thus the bulk of the work in this proof is to prove the integrability.

It is also known that a contact metric Einstein manifold of dimension ≥ 5 with ξ belonging to the κ -nullity distribution is Sasakian (Tanno [1988]). In fact, aside from the flat 3-dimensional case there are no non-Sasakian Einstein (κ, μ) -manifolds, as can easily be seen from the formula for the Ricci operator of a (κ, μ) -manifold given in the last section.

In [2006] Apostolov, Draghici and Moroianu also prove the following theorem.

Theorem 7.13 *Let (M, ϕ, ξ, η, g) be a Sasakian Einstein manifold.. Then any contact metric structure on M with characteristic vector field ξ' and the same metric g is Sasakian. Moreover, if $\xi' \neq \pm\xi$, then either (M, g) admits a 3-Sasakian structure (see Chapter 14) or (M, g) is covered by a round sphere.*

There are many Sasakian Einstein manifolds, as shown by Boyer and Galicki in [2000] and in their book [2008, Chapter 11], including the well-known Sasakian Einstein structure on $S^3 \times S^2$ (see Tanno [1978b] and Section 9.2). Generalizing this last example, Boyer, Galicki and Nakamaye [2003b] showed that $S^3 \times S^2$ carries 14 inequivalent Sasakian Einstein structures and that there are infinite families of such structures on the connected sums $S^3 \times S^2 \# \cdots \# S^3 \times S^2$ with $2, \dots, 7$ summands. More recently, J. Kollár [2007] proved the existence of families of Sasakian Einstein metrics for any number of summands ≥ 6 . We also note that very explicit expressions for Sasakian Einstein metrics on $S^3 \times S^2$ were given in Gauntlett, Martelli, Sparks and Waldram [2004].

It is also known that there exist Sasakian Einstein metrics on many homotopy spheres; see Boyer, Galicki and Kollár [2005] and Boyer and Galicki [2008, Section 11.5].

Let us now turn briefly to the η -Einstein case. Recall that this means that the Ricci tensor $\rho = ag + b\eta \otimes \eta$, where initially a and b are functions, but if the dimension is ≥ 5 , a and b must be constants. S. Morimoto [1992] and independently Boyer and Galicki [2001] prove the following theorem.

Theorem 7.14 *If M^{2n+1} is a compact η -Einstein K -contact manifold with Ricci tensor, then $\rho = ag + b\eta \otimes \eta$, and if $a \geq -2$, then M^{2n+1} is Sasakian.*

Boyer and Galicki in their treatment of this question prove more, namely, that if $a > -2$, then the \mathcal{D} -homothetically deformed metric $g = \alpha g + \alpha(\alpha - 1)\eta \otimes \eta$ with $\alpha = \frac{a+2}{2n+2}$ is Sasakian Einstein and hence $\pi_1(M^{2n+1})$ is finite. For further results on Sasakian η -Einstein structures we refer to the paper of Boyer, Galicki and Matzeu [2006] and to the book of Boyer and Galicki [2008, Sections 11.1 and 11.8].

7.5 Locally symmetric contact metric manifolds

The question of locally symmetric contact metric manifolds has a long history, but the list of such manifolds is very short, and we highlight just a few of the results in this direction.

Already in [1962a] Okumura proved that a locally symmetric Sasakian manifold is of constant curvature $+1$. This was generalized to the K-contact case by Tanno [1967b].

In [1989] the author showed that the standard contact metric structure of the tangent sphere bundle (or unit tangent bundle but with a homothetic change of metric) T_1M of a Riemannian manifold M is locally symmetric if and only if either M is flat, in which case T_1M is locally isometric to $E^{n+1} \times S^n(4)$, or M is 2-dimensional and of constant curvature $+1$, in which case T_1M is locally isometric to $T_1S^2 \sim \mathbb{R}P^3 \sim SO(3)$. We remark that even though T_1S^3 is topologically $S^3 \times S^2$, the product metric is not an associated metric to the natural contact structure (see Section 9.2).

The above results raise the question of the classification of all locally symmetric contact metric manifolds, and one might conjecture that the only two possibilities are contact metric manifolds that are locally isometric to $S^{2n+1}(1)$ or $E^{n+1} \times S^n(4)$. In dimension 3, R. Sharma and the author [1990b] showed that a locally symmetric contact metric manifold is of constant curvature 0 or 1. In dimension 5, A. M. Pastore [1998] showed that indeed local isometry with $S^5(1)$ or $E^3 \times S^2(4)$ are the only two possibilities.

In [1994] K. Bang showed that a locally symmetric contact metric manifold with $R_X \xi \xi = 0$ is locally isometric to $E^{n+1} \times S^n(4)$; this answered positively a question posed by D. Perrone [1992a].

When the contact metric manifold is a CR-manifold (i.e., $(\nabla_X \phi)Y = g(X + hX, Y)\xi - \eta(Y)(X + hX)$) of dimension $2n + 1 > 3$ but $\neq 7$,

Ghosh and Sharma showed in [1999] that the condition of being locally symmetric implies that the manifold is locally isometric to $S^{2n+1}(1)$ or $E^{n+1} \times S^n(4)$.

Finally, in [2006], without any further restrictions, Boeckx and Cho proved the following classification theorem.

Theorem 7.15 *A locally symmetric contact metric manifold is locally isometric to $S^{2n+1}(1)$ or $E^{n+1} \times S^n(4)$.*

7.6 Conformally flat contact metric manifolds

Instead of asking when a contact metric manifold is locally symmetric, we now consider the question of a contact metric manifold being conformally flat. In [1962a] Okumura showed that a conformally flat Sasakian manifold of dimension ≥ 5 is of constant curvature $+1$ and in [1963,1967b] Tanno extended this result to the K-contact case and for dimensions ≥ 3 .

Theorem 7.16 *A conformally flat K-contact manifold is of constant curvature $+1$ and Sasakian.*

In a similar vein, if a contact metric manifold of dimension ≥ 5 is a CR-manifold, Ghosh, Koufogiorgos and Sharma [2001] showed that conformal flatness implies constant curvature $+1$.

In dimension ≥ 5 , as we have seen, a contact metric structure of constant curvature must be of constant curvature $+1$ and in dimension 3 a contact metric structure of constant curvature must be of constant curvature 0 or $+1$. For simplicity and future use we introduce another symmetric operator l by

$$lX = R_{X\xi}\xi.$$

K. Bang [1994] showed that in dimension ≥ 5 there are no conformally flat contact metric structures with $l = 0$, even though there are many contact metric manifolds satisfying $l = 0$; see Theorem 9.16. Bang's result was extended to dimension 3 and in fact generalized by Gouli-Andreou and Xenos [1999] who showed that in dimension 3 the only conformally flat contact metric structures satisfying $\nabla_\xi l = 0$ (equivalent to $\nabla_\xi h = 0$, Perrone [1992a]) are those of constant curvature 0 or 1.

In the case of the standard contact metric structure on the tangent sphere bundle (Section 9.2), the condition of conformal flatness is quite extreme; in fact, the metric is conformally flat if and only if the base

manifold is a surface of constant Gaussian curvature 0 or 1 (Theorem 9.6), as was shown by Th. Koufogiorgos and the author [1994].

In [1999] Calvaruso, Perrone and Vanhecke showed that in dimension 3 the only conformally flat contact metric structures for which ξ is an eigenvector of the Ricci tensor are those of constant curvature 0 or 1. This seems to be a somewhat key condition and has attracted the attention of several authors. Finally, in [2004] the corresponding result in higher dimensions was achieved by F. Gouli-Andreou and N. Tsolakidou. An independent and basis-free proof was given by K. Bang and the author in [2008].

Theorem 7.17 *A conformally flat contact metric manifold whose characteristic vector field is everywhere an eigenvector of the Ricci operator is of constant curvature.*

A contrasting condition to ξ being an eigenvector of the Ricci tensor, namely that $Q\xi$ be orthogonal to ξ , was considered by F. Gouli-Andreou and R. Sharma [2003]. They showed that a compact 3-dimensional conformally flat contact metric manifold satisfying this condition is flat.

In view of these strong curvature results one may ask whether there are any conformally flat contact metric structures that are not of constant curvature. We devote the rest of this section to showing the local existence and giving some additional remarks.

We will work in \mathbb{R}^3 , primarily with cylindrical coordinates (r, θ, z) . Let $\eta = \frac{1}{2}(\alpha dr + \beta r d\theta + \gamma dz)$ be a contact form on \mathbb{R}^3 . Then

$$d\eta = \frac{1}{2}((\beta + r\beta_r - \alpha_\theta)dr \wedge d\theta + (\gamma_r - \alpha_z)dr \wedge dz + (\gamma_\theta - r\beta_z)d\theta \wedge dz).$$

If g is a conformally flat metric, we may write it as $ds^2 = \frac{1}{4}e^{2\sigma}(dr^2 + r^2d\theta^2 + dz^2)$. If g is also an associated metric, the characteristic vector field is given by

$$\xi = 2e^{-2\sigma}\left(\alpha\frac{\partial}{\partial r} + \beta\frac{1}{r}\frac{\partial}{\partial\theta} + \gamma\frac{\partial}{\partial z}\right),$$

and $\eta(\xi) = 1$ gives $e^{2\sigma} = \alpha^2 + \beta^2 + \gamma^2$. Computing g^{-1} , we can obtain ϕ from $d\eta$ but we must have $d\eta(\xi, X) = 0$ and $\phi^2 = -I + \eta \otimes \xi$.

Much of our analysis will actually be done with respect to the Euclidean metric on \mathbb{R}^3 , and we denote the Euclidean length of a vector field \mathbf{B} by $|\mathbf{B}|$. In particular, if $\mathbf{B} = \alpha\frac{\partial}{\partial r} + \beta\frac{1}{r}\frac{\partial}{\partial\theta} + \gamma\frac{\partial}{\partial z}$, then $|\mathbf{B}| = e^\sigma$.

Now $d\eta(\xi, X) = 0$ for all X gives

$$\begin{aligned}\frac{1}{r}\beta(\beta + r\beta_r - \alpha_\theta) + \gamma(\gamma_r - \alpha_z) &= 0, \\ \alpha(\alpha_\theta - \beta - r\beta_r) + \gamma(\gamma_\theta - r\beta_z) &= 0, \\ \alpha(\alpha_z - \gamma_r) + \frac{1}{r}\beta(r\beta_z - \gamma_\theta) &= 0,\end{aligned}$$

and therefore $\mathbf{curl} \mathbf{B}$ is proportional to \mathbf{B} , say $\mathbf{curl} \mathbf{B} = f\mathbf{B}$. Then $\phi^2 = -I + \eta \otimes \xi$ yields $f^2 = e^{2\sigma}$. Therefore $\mathbf{curl} \mathbf{B} = \pm e^\sigma \mathbf{B}$. Since we may change the sign of η without changing the problem, our study reduces to the study of the differential equation

$$\mathbf{curl} \mathbf{B} = |\mathbf{B}|\mathbf{B}.$$

Thus we are at the point that the existence of 3-dimensional conformally flat contact metric manifolds corresponds to finding solutions of $\mathbf{curl} \mathbf{B} = |\mathbf{B}|\mathbf{B}$. Had we carried out our analysis in Cartesian coordinates (x, y, z) , we would have been led to the same differential equation for a vector field $\mathbf{B} = \alpha \frac{\partial}{\partial x} + \beta \frac{\partial}{\partial y} + \gamma \frac{\partial}{\partial z}$. The familiar unit vector field $\mathbf{B} = \sin z \frac{\partial}{\partial x} + \cos z \frac{\partial}{\partial y}$ that is equal to its own \mathbf{curl} corresponds to the flat case. Stereographic projection from S^3 to \mathbb{R}^3 gives the vector field

$$\xi = (xz - y) \frac{\partial}{\partial x} + (x + yz) \frac{\partial}{\partial y} + \frac{1}{2}(1 + z^2 - x^2 - y^2) \frac{\partial}{\partial z}$$

corresponding to the contact form

$$\eta = \frac{4(xz - y)dx + 4(x + yz)dy + 2(1 + z^2 - x^2 - y^2)dz}{(1 + x^2 + y^2 + z^2)^2}$$

for which $g = \frac{4(dx^2 + dy^2 + dz^2)}{(1 + x^2 + y^2 + z^2)^2}$ is an associated metric of constant curvature +1. The vector field $\mathbf{B} = \frac{8}{(1 + x^2 + y^2 + z^2)^2} \xi$ satisfies $\mathbf{curl} \mathbf{B} = |\mathbf{B}|\mathbf{B}$, though we remark that it is not of vanishing divergence.

In both of the two constant curvature examples above, the component functions are not functions of the radial coordinate alone. If $\mathbf{curl} \mathbf{B} = |\mathbf{B}|\mathbf{B}$ and the component functions are functions of r alone, then $\alpha = 0$ and the other components, $\beta(r)$ and $\gamma(r)$, satisfy

$$\frac{1}{r}\beta + \beta' = \sqrt{\beta^2 + \gamma^2} \gamma, \quad -\gamma' = \sqrt{\beta^2 + \gamma^2} \beta. \quad (*)$$

Local existence away from $r = 0$ follows easily from the standard existence theorems for systems of ordinary differential equations.

Finally, one can show that the metric $ds^2 = \frac{1}{4}e^{2\sigma}(dr^2 + r^2d\theta^2 + dz^2)$ with $e^{2\sigma} = \beta^2 + \gamma^2$ is not of constant curvature. Differentiating $e^{2\sigma} = \beta^2 + \gamma^2$ using (*), we have $\sigma' = -\beta^2/re^{2\sigma}$. If g were of constant curvature, the value of the curvature would be 0 or +1. Consider the unit vector field $X = 2e^{-2\sigma}(\frac{\gamma}{r}\frac{\partial}{\partial\theta} - \beta\frac{\partial}{\partial z})$ belonging to the contact subbundle \mathcal{D} . Then computing the sectional curvature of the plane section spanned by ξ and X , we have

$$g(R_{X\xi\xi}, X) = -4e^{-2\sigma}\left(\sigma'^2 + \frac{\sigma'}{r}\right) = \frac{4e^{-6\sigma}}{r^2}\beta^2\gamma^2.$$

If the constancy of the curvature were 0, then β or γ would vanish, and from (*) both would vanish, making $\eta \equiv 0$, a contradiction. If the curvature is +1, then $re^{3\sigma} = 2\beta\gamma$; differentiating this using (*) and $\sigma' = -\frac{\beta^2}{re^{2\sigma}}$, we again have that $\beta = 0$.

This class of examples was studied further by G. Calvaruso [2000]; he showed that

$$\nabla_\xi h = ah\phi, \quad a = \frac{2\beta\gamma}{re^{3\sigma}} \neq \text{constant}.$$

He also showed that if a is a constant $\neq 2$, then a 3-dimensional conformally flat contact metric manifold satisfying $\nabla_\xi h = ah\phi$ has constant curvature. It is not known whether there exist conformally flat contact metric manifolds of dimension ≥ 5 that are not locally isometric to the standard Sasakian structure on the unit sphere.

The equation $\mathbf{curl} \mathbf{B} = f\mathbf{B}$ for some function f arises in solar physics, and $\mathbf{curl} \mathbf{B} = |\mathbf{B}|\mathbf{B}$ can be viewed as a special case; $\mathbf{B} = \mu\mathbf{H}$, where \mathbf{H} is the magnetic field and μ the magnetic permeability. An “active region” on the sun is an area of extreme magnetic flux. These regions contain sunspots as well as the highly volatile phenomena of solar flares. For some solar phenomena, such as flares and prominences, the Lorentz force, $\mathbf{j} \times \mathbf{B}$, \mathbf{j} being the current density, dominates the pressure gradient and gravitational forces and thus for a relatively slow moving plasma we have from the equations of motion the approximation $\mathbf{j} \times \mathbf{B} = 0$, the so-called “force-free” field. From Maxwell’s equations we have for an electrically neutral field $\mathbf{curl} \mathbf{B} = \mu\mathbf{j}$. Thus one is led to $\mathbf{curl} \mathbf{B} = f\mathbf{B}$. There are many solutions in the literature (see, e.g., the book by E. Priest [1982]) which by Maxwell’s equation must also satisfy $\mathbf{div} \mathbf{B} = 0$.

Except for the simple case $\mathbf{B} = \sin z \frac{\partial}{\partial x} + \cos z \frac{\partial}{\partial y}$, the solutions listed in Priest [1982] do not satisfy $\mathbf{curl} \mathbf{B} = |\mathbf{B}|\mathbf{B}$. Our vector field is divergence-free and hence another solution; also its series expansion converges rapidly, so that its calculation for physical purposes is not prohibitive.

7.7 ϕ -sectional curvature

In this section we introduce the notion of ϕ -sectional curvature. This idea plays the role in Sasakian geometry that holomorphic sectional curvature plays in Kähler geometry. A plane section in $T_m M^{2n+1}$ is called a ϕ -section if there exists a vector $X \in T_m M^{2n+1}$ orthogonal to ξ such that $\{X, \phi X\}$ span the section. The sectional curvature $K(X, \phi X)$, denoted $H(X)$, is called ϕ -sectional curvature.

Recall that the sectional curvatures of a Riemannian manifold determine the curvature transformation $R_{X,Y}Z$. It is also well known that the holomorphic sectional curvatures of a Kähler manifold determine the curvature completely. We shall show that on a Sasakian manifold the ϕ -sectional curvatures determine the curvature completely (Moskal [1966]). Let $B(X, Y) = g(R_{X,Y}Y, X)$ and for X orthogonal to ξ , let $D(X) = B(X, \phi X)$. Also recall the tensor field P defined by

$$\begin{aligned} P(X, Y, Z, W) = & d\eta(X, Z)g(Y, W) - d\eta(X, W)g(Y, Z) \\ & - d\eta(Y, Z)g(X, W) + d\eta(Y, W)g(X, Z) \end{aligned}$$

in Section 7.1.

Proposition 7.8 *On a Sasakian manifold, for tangent vectors X and Y orthogonal to ξ we have*

$$\begin{aligned} B(X, Y) = \frac{1}{32} \Big(& 3D(X + \phi Y) + 3D(X - \phi Y) - D(X + Y) - D(X - Y) \\ & - 4D(X) - 4D(Y) - 24P(X, Y, X, \phi Y) \Big). \end{aligned}$$

Proof. Direct expansion and Lemma 7.1 give

$$\begin{aligned} \frac{1}{32} \Big(& 3D(X + \phi Y) + 3D(X - \phi Y) - D(X + Y) - D(X - Y) \\ & - 4D(X) - 4D(Y) - 24P(X, Y, X, \phi Y) \Big) \end{aligned}$$

$$\begin{aligned}
&= \frac{1}{32} \left(6g(R_{X \cdot Y}Y, X) + 6g(R_{\phi X \cdot \phi Y}\phi Y, \phi X) + 8g(R_{X \cdot \phi X}\phi Y, Y) \right. \\
&\quad + 12g(R_{X \cdot Y}\phi Y, \phi X) - 2g(R_{X \cdot \phi Y}\phi Y, X) - 2g(R_{\phi X \cdot Y}Y, \phi X) \\
&\quad \left. + 4g(R_{X \cdot \phi Y}Y, \phi X) - 24P(X, Y, X, \phi Y) \right) \\
&= g(R_{X \cdot Y}Y, X).
\end{aligned}$$

■

Proposition 7.9 *Let M be a Sasakian manifold and $\{X, Y\}$ an orthonormal pair in $T_m M$ with X and Y orthogonal to ξ . Set $g(X, \phi Y) = \cos \theta$, $0 \leq \theta \leq \pi$. Then the sectional curvature $K(X, Y)$ is given by*

$$\begin{aligned}
K(X, Y) &= \frac{1}{8} \left(3(1 + \cos \theta)^2 H(X + \phi Y) + 3(1 - \cos \theta)^2 H(X - \phi Y) \right. \\
&\quad \left. - H(X + Y) - H(X - Y) - H(X) - H(Y) + 6 \sin^2 \theta \right).
\end{aligned}$$

Proof. $K(X, Y) = B(X, Y)$ in the previous proposition, so we examine the terms in the expansion of $B(X, Y)$. Clearly $D(X) = g(X, X)^2 H(X)$ for any X orthogonal to ξ , and hence for the given pair $\{X, Y\}$, $g(X + \phi Y, X + \phi Y) = 2(1 + \cos \theta)$, $g(X - \phi Y, X - \phi Y) = 2(1 - \cos \theta)$, $g(X + Y, X + Y) = 2$ and $g(X - Y, X - Y) = 2$. Thus $D(X + \phi Y) = 4(1 + \cos \theta)^2 H(X + \phi Y)$, and so on. Finally, note that $P(X, Y, X, \phi Y) = -\sin^2 \theta$, completing the proof. ■

Theorem 7.18 *The ϕ -sectional curvatures of a Sasakian manifold determine the curvature completely.*

Proof. Since the sectional curvatures of a Riemannian manifold determine the curvature, it suffices to show that for an orthonormal pair $\{X, Y\}$, $K(X, Y)$ is determined uniquely by H and g . If X and Y are orthogonal to ξ , the previous proposition applies. If X or Y is ξ , $K(X, Y) = 1$. So suppose that $X = \eta(X)\xi + aZ$ and $Y = \eta(Y)\xi + bW$, where $\eta(X)$, $\eta(Y)$, $a = \sqrt{1 - \eta(X)^2}$, and $b = \sqrt{1 - \eta(Y)^2}$ are nonzero. Recall that on a Sasakian manifold, $R_{\xi Z}\xi = -Z$ and $R_{Z\xi}Z = -\xi$ for any unit vector orthogonal to ξ . Therefore

$$\begin{aligned}
K(X, Y) &= g(R_{\eta(X)\xi + aZ} \eta(Y)\xi + bW, \eta(X)\xi + aZ) \\
&= b^2 \eta(X)^2 - 2ab \eta(X) \eta(Y) g(Z, W) + a^2 \eta(Y)^2 \\
&\quad + a^2 b^2 g(R_{ZW}W, Z).
\end{aligned}$$

Now $g(Z, W) + \frac{1}{ab} g(X - \eta(X)\xi, Y - \eta(Y)\xi) = -\frac{1}{ab} \eta(X)\eta(Y)$, giving

$$\begin{aligned} g(R_{ZW}W, Z) &= (1 - g(Z, W)^2)K(Z, W) \\ &= \left(1 - \frac{1}{a^2b^2}\eta(X)^2\eta(Y)^2\right)K(Z, W). \end{aligned}$$

Thus

$$\begin{aligned} K(X, Y) &= \eta(X)^2(1 - \eta(Y)^2) + 2\eta(X)^2\eta(Y)^2 + \eta(Y)^2(1 - \eta(X)^2) \\ &\quad + ((1 - \eta(X)^2)(1 - \eta(Y)^2) - \eta(X)^2\eta(Y)^2)K(Z, W) \\ &= \eta(X)^2 + \eta(Y)^2 + (1 - \eta(X)^2 - \eta(Y)^2)K(Z, W), \end{aligned}$$

and $K(Z, W)$ is given by the previous proposition completing the proof. \blacksquare

Note that the above proof uses not only the values of the ϕ -sectional curvatures, but also the facts that on a Sasakian manifold $R_{\xi X}\xi = -X$ and $R_{X\xi}X = -\xi$ for any unit vector X orthogonal to ξ . Thus we have actually proved that any tensor field of type $(1, 3)$ on a Sasakian manifold that satisfies the symmetries of the curvature tensor, the first Bianchi identity, identity (a) of Lemma 7.1, $R_{\xi X}\xi = -X$, and $R_{X\xi}X = -\xi$ for any unit vector X orthogonal to ξ and that agrees with the values of the ϕ -sectional curvatures must be the curvature tensor. Therefore we can easily prove the following theorem of Ogiue [1964].

Theorem 7.19 *If the ϕ -sectional curvature at any point of a Sasakian manifold of dimension ≥ 5 is independent of the choice of ϕ -section at the point, then it is constant on the manifold and the curvature tensor is given by*

$$\begin{aligned} R_{XY}Z &= \frac{c+3}{4}(g(Y, Z)X - g(X, Z)Y) \\ &\quad + \frac{c-1}{4}\left(\eta(X)\eta(Z)Y - \eta(Y)\eta(Z)X + g(X, Z)\eta(Y)\xi \right. \\ &\quad \left. - g(Y, Z)\eta(X)\xi + \Phi(Z, Y)\phi X - \Phi(Z, X)\phi Y + 2\Phi(X, Y)\phi Z\right), \end{aligned}$$

where c is the constant ϕ -sectional curvature.

Proof. In view of the above remark, in order to see that the curvature tensor has the above form with c a function on the manifold one need

only check the necessary conditions, and this is easily done. The Ricci tensor ρ and the scalar curvature τ are given by

$$\rho(X, Y) = \frac{n(c+3) + c - 1}{2}g(X, Y) - \frac{(n+1)(c-1)}{2}\eta(X)\eta(Y)$$

and

$$\tau = \frac{1}{2}(n(2n+1)(c+3) + n(c-1)).$$

Now from the second Bianchi identity, $\nabla_\alpha \tau - 2\nabla_\beta \rho^\beta_\alpha = 0$, where ρ^β_α are the components of the Ricci tensor of type $(1, 1)$, and hence

$$(n-1)dc + (\xi c)\eta = 0.$$

Applying this to ξ , we have $\xi c = 0$ and hence $dc = 0$ for $n \neq 1$, as desired. ■

A Sasakian manifold of constant ϕ -sectional curvature c will be called a *Sasakian space form* and denoted by $M(c)$. Note that a Sasakian space form has constant scalar curvature and is η -Einstein. Also if $c < 1$ we have the following pinching of the sectional curvature, similar to that in the Kähler case:

$$c \leq K(X, Y) \leq \frac{c+3}{4};$$

if $c > 1$, the inequalities are reversed.

Th. Koufogiorgos [1997a] studied (κ, μ) -manifolds of dimension ≥ 5 for which the ϕ -sectional curvature at any point is independent of the choice of ϕ -section at the point. He proved that the ϕ -sectional curvature is constant and obtained the curvature tensor explicitly.

In the general context of contact metric manifolds J. T. Cho [2003] introduced the notion of a *contact Riemannian space form*. We get at this notion in the following way. In Mitric [1991] and Tanno [1992] it was shown that the tangent sphere bundle with its standard contact metric structure is a CR-manifold if and only if the base manifold is of constant curvature (see Theorem 9.9 below). Cho first computes the covariant derivative of h in this case, obtaining

$$(\nabla_X h)Y = g((h - h^2)\phi X, Y)\xi + \eta(Y)(h - h^2)\phi X - \mu\eta(X)h\phi Y,$$

where μ is a constant. He then abstracts this idea and defines the class \mathfrak{Q} of contact metric CR-manifolds for which the covariant derivative of

h satisfies the above condition. We remark that in the study of contact manifolds in general, lack of control of the covariant derivative of h is often an obstacle to further results.

Now for a contact metric manifold M^{2n+1} of class \mathfrak{Q} with $n > 1$ and for which the ϕ -sectional curvature is independent of the choice of ϕ -section, Cho shows that the ϕ -sectional curvature is constant on M^{2n+1} and he computes the curvature tensor explicitly. He then defines a *contact Riemannian space form* to be a complete, simply connected contact metric manifold of class \mathfrak{Q} of constant ϕ -sectional curvature. Cho also gives a number of non-Sasakian examples and shows that a contact Riemannian space form is locally homogeneous and is strongly locally ϕ -symmetric (see Section 7.9).

Another generalization of a Sasakian space form was introduced by P. Alegre, A. Carriazo and the author in [2004] and studied further by Alegre and Carriazo in [2008]. An almost contact metric manifold is a *generalized Sasakian space form* if its curvature tensor is of the form

$$\begin{aligned} R_{XY}Z = & f_1(g(Y, Z)X - g(X, Z)Y) \\ & + f_2(g(X, \phi Z))\phi Y - g(Y, \phi Z)\phi X + 2g(X, \phi Y)\phi Z \\ & + f_3(\eta(X)\eta(Z)Y - \eta(Y)\eta(Z)X + g(X, Z)\eta(Y)\xi \\ & - g(Y, Z)\eta(X)\xi), \end{aligned}$$

where the f_i 's are differentiable functions. There exist numerous examples, but the most interesting ones are on almost contact metric manifolds that are not contact metric manifolds. In particular, in dimensions ≥ 5 a contact metric generalized Sasakian space form is a Sasakian space form. For this reason we will not give a detailed discussion and refer the reader to the references. The special case of $f_2 = f_3$ with some additional conditions was considered by Bueken and Vanhecke [1988b].

7.8 Examples of Sasakian space forms

To begin, let (ϕ, ξ, η, g) be a contact metric structure and recall the notion of a *\mathcal{D} -homothetic deformation*:

$$\bar{\eta} = a\eta, \quad \bar{\xi} = \frac{1}{a}\xi, \quad \bar{\phi} = \phi, \quad \bar{g} = ag + a(a-1)\eta \otimes \eta,$$

where a is a positive constant. The deformed structure $(\bar{\phi}, \bar{\xi}, \bar{\eta}, \bar{g})$ is again a contact metric structure, and it enjoys many of the properties of the original structure as we remarked in Section 7.3. In particular, if (ϕ, ξ, η, g) is Sasakian, so is $(\bar{\phi}, \bar{\xi}, \bar{\eta}, \bar{g})$; if $M(c)$ is a Sasakian space form, then deforming the structure, we obtain the Sasakian space form $M(\bar{c})$ where $\bar{c} = \frac{c+3}{a} - 3$ (see Tanno [1968], [1969] for details). We will show that there exist Sasakian space forms $M(c)$ for every value of c . Moreover, we state the following theorem of Tanno and refer to [1969] for the proof.

Theorem 7.20 *Let $M(c)$ be a complete, simply connected Sasakian manifold with constant ϕ -sectional curvature c . Then $M(c)$ belongs one of the three families of examples listed below.*

7.8.1 S^{2n+1}

Let (ϕ, ξ, η, g) be the standard contact metric structure on the sphere S^{2n+1} constructed either as a hypersurface of \mathbb{C}^{n+1} (Example 4.5.2 and Section 6.3) or as a principal circle bundle over $\mathbb{C}P^n$ (Examples 4.5.4 and 6.7.2). Applying a \mathcal{D} -homothetic deformation to this structure, one obtains a Sasakian structure on S^{2n+1} with constant ϕ -sectional curvature $c = \frac{4}{a} - 3$. Note that from the remark on pinching in Section 7.7, these metrics on S^{2n+1} for $c > 0$ satisfy the condition of Goldberg's theorem [1967] referred to in Section 6.8.

7.8.2 \mathbb{R}^{2n+1}

In Examples 4.5.1 and 6.7.1 we saw that \mathbb{R}^{2n+1} with coordinates (x^i, y^i, z) , $i = 1, \dots, n$, admits the Sasakian structure

$$\eta = \frac{1}{2}(dz - \sum_{i=1}^n y^i dx^i), \quad g = \eta \otimes \eta + \frac{1}{4} \sum_{i=1}^n ((dx^i)^2 + (dy^i)^2).$$

With this metric, \mathbb{R}^{2n+1} is a Sasakian space form with $c = -3$ (cf. Okumura [1962b]) and often denoted by $\mathbb{R}^{2n+1}(-3)$.

7.8.3 $B^n \times \mathbb{R}$

Let B^n be a simply connected bounded domain in \mathbb{C}^n and (J, G) a Kähler structure with constant holomorphic sectional curvature $k < 0$. For such a structure, the fundamental 2-form Ω of the Kähler structure is exact

and hence $\Omega = d\omega$ for some real analytic 1-form ω . Now on $B^n \times \mathbb{R}$ let π denote the projection onto B^n and t the coordinate on \mathbb{R} . Then $B^n \times \mathbb{R}$ with $\eta = \pi^*\omega + dt$ and $g = \pi^*G + \eta \otimes \eta$ is a Sasakian manifold. Regarding η as a connection form on $B^n \times \mathbb{R}$, let $\tilde{\pi}X$ denote the horizontal lift of a vector field X on B^n . Also we denote by \underline{K} the sectional curvature of B^n . Then by direct computation (see Ogiue [1965] or use the Riemannian submersion technique of O'Neill [1966]), we have $K(\tilde{\pi}X, \tilde{\pi}Y) = \underline{K}(X, Y) - 3\eta(\nabla_{\tilde{\pi}X}\tilde{\pi}Y)^2$ where $\{X, Y\}$ is an orthonormal pair on B^n . Now $g(\nabla_{\tilde{\pi}X}\tilde{\pi}Y, \xi) = -g(\tilde{\pi}Y, \nabla_{\tilde{\pi}X}\xi) = g(\tilde{\pi}Y, \phi\tilde{\pi}X) = g(\tilde{\pi}Y, \tilde{\pi}JX)$. Therefore $B^n \times \mathbb{R}$ has constant ϕ -sectional curvature $c = k - 3$.

7.9 Locally ϕ -symmetric spaces

We have seen that the only locally symmetric Sasakian manifolds are locally isometric to $S^{2n+1}(1)$ and that the only locally symmetric contact metric manifolds are locally isometric to $S^{2n+1}(1)$ or to $E^{n+1} \times S^n(4)$. Certainly this can be regarded as saying that the idea of being locally symmetric is too strong. For this reason, T. Takahashi [1977] introduced the notion of a locally ϕ -symmetric space. A Sasakian manifold is said to be a *Sasakian locally ϕ -symmetric space* if


$$\phi^2(\nabla_V R)_{XY}Z = 0$$

for all vector fields V, X, Y, Z orthogonal to ξ ; such spaces were called *locally \mathcal{D} -symmetric spaces* by Shibuya [1982]. It is easy to check that Sasakian space forms are locally ϕ -symmetric spaces. In [1987a] L. Vanhecke and the author showed that a Sasakian manifold is locally ϕ -symmetric if and only if

$$g((\nabla_X R)_{X\phi X}X, \phi X) = 0$$

for all vector fields X orthogonal to ξ .

Note that on a Sasakian manifold M , or more generally on a K-contact manifold, a geodesic that is initially orthogonal to ξ remains orthogonal to ξ . We call such a geodesic a *ϕ -geodesic*. A local diffeomorphism s_m of M , $m \in M$, is a *ϕ -geodesic symmetry* if its domain contains a (possibly) smaller domain \mathcal{U} such that for every ϕ -geodesic $\gamma(s)$ parametrized by



arc length we have (i) $\gamma(0)$ is in the intersection of \mathcal{U} and the integral curve of ξ through m , and (ii)

$$(s_m \circ \gamma)(s) = \gamma(-s)$$

for all s with $\gamma(\pm s) \in \mathcal{U}$. Since the points of the integral curve of ξ through m are fixed, we see that setting $S = -I + 2\eta \otimes \xi$, we have

$$s_m = \exp_m \circ S_m \circ \exp_m^{-1}.$$

Takahashi [1977] defines a Sasakian manifold to be a *Sasakian globally ϕ -symmetric space* by requiring that any ϕ -geodesic symmetry can be extended to a global automorphism of the structure and that the Killing vector field ξ generate a 1-parameter group of global transformations.

Among the main results of Takahashi [1977] are the following four theorems.

Theorem 7.21 *A Sasakian locally ϕ -symmetric space is locally isometric to a Sasakian globally ϕ -symmetric space, and a complete, connected, simply connected Sasakian locally ϕ -symmetric space is globally ϕ -symmetric.*

Theorem 7.22 *A Sasakian manifold is locally ϕ -symmetric if and only if it admits a ϕ -geodesic symmetry at every point that is a local automorphism of the structure.*

Now suppose that \mathcal{U} is a neighborhood on M on which ξ is regular. Then since M is Sasakian, the projection $\pi : \mathcal{U} \rightarrow \mathcal{V} = \mathcal{U}/\xi$ gives a Kähler structure on \mathcal{V} . Furthermore, if $\underline{s}_{\pi(m)}$ denotes the geodesic symmetry on \mathcal{V} at $\pi(m)$, then $\underline{s}_{\pi(m)} \circ \pi = \pi \circ s_m$.

Theorem 7.23 *A Sasakian manifold is locally ϕ -symmetric if and only if each Kähler manifold that is the base of a local fibering is a Hermitian locally symmetric space.*

Following Okumura [1962a], we define on a Sasakian manifold a linear connection $\bar{\nabla}$ by $\bar{\nabla}_X Y = \nabla_X Y + T_X Y$, where

$$T_X Y = d\eta(X, Y)\xi - \eta(X)\phi Y + \eta(Y)\phi X,$$

and it is easy to see that the structure tensors are parallel with respect to this connection. Takahashi [1977] then proves the following.

Theorem 7.24 *A Sasakian manifold is a locally ϕ -symmetric space if and only if $\bar{\nabla}\bar{R} = 0$, equivalently,*

$$(\nabla_V R)_{XY}Z = -T_V R_{XY}Z + R_{TVXY}Z + R_X T_V YZ + R_{XY}T_V Z$$

for all X, Y, Z, V .

In the spirit of the fact that a Riemannian manifold is locally symmetric if and only if the local geodesic symmetries are isometries and in view of the above results of Takahashi, we state the following extension of Theorem 7.21 (sufficiency in the Sasakian case, Vanhecke and the author [1987b]; in the K-contact case, Bueken and Vanhecke [1989]).

Theorem 7.25 *On a Sasakian locally ϕ -symmetric space, local ϕ -geodesic symmetries are isometries. Conversely, if on a K-contact manifold the local ϕ -geodesic symmetries are isometries, the manifold is a Sasakian locally ϕ -symmetric space.*

Originally, the notion of a locally ϕ -symmetric space was for the most part explored only in the Sasakian context, and it was not clear what the corresponding notion should be for a general contact metric manifold. Without the K-contact property one loses the fact that a geodesic initially orthogonal to ξ remains orthogonal to ξ . We have just seen that in the Sasakian case local ϕ -symmetry is equivalent to reflections in the integral curves of the characteristic vector field being isometries. Boeckx and Vanhecke [1997] proposed this property as the definition for local ϕ -symmetry in the contact metric case and formalized two notions in Boeckx, Bueken and Vanhecke [1999]. We adopt this formulation here. A contact metric manifold is a *weakly locally ϕ -symmetric space* if it satisfies

$$\phi^2(\nabla_V R)_{XY}Z = 0$$

for all vector fields V, X, Y, Z orthogonal to ξ as in the Sasakian case. A contact metric manifold is a *strongly locally ϕ -symmetric space* if reflections in the integral curves of the characteristic vector field are isometries.

From Chen and Vanhecke [1989] (see also Boeckx, Bueken and Vanhecke [1999]) one sees that on a strongly locally ϕ -symmetric space,

$$\begin{aligned} g((\nabla_X^{2k} \dots_X R)_{XY} X, \xi) &= 0, \\ g((\nabla_X^{2k+1} R)_{XY} X, Z) &= 0, \\ g((\nabla_X^{2k+1} R)_{X\xi} X, \xi) &= 0, \end{aligned}$$

for all X, Y, Z orthogonal to ξ and all $k \in \mathbb{N}$. Conversely, on an analytic Riemannian manifold these conditions are sufficient for the contact metric manifold to be a strongly locally ϕ -symmetric space. In particular, taking $k = 0$ in the second condition, we note that a strongly locally ϕ -symmetric space is weakly locally ϕ -symmetric. Calvaruso, Perrone and Vanhecke [1999] showed that a 3-dimensional strongly locally ϕ -symmetric space is either K-contact with constant scalar curvature or is a (κ, μ) -manifold with $\kappa < 1$. They also showed that a 3-dimensional contact metric manifold is a strongly locally ϕ -symmetric space if and only if it is locally contact homogeneous, i.e., the pseudogroup of local automorphisms of the contact metric structure acts transitively on the manifold, and ξ is an eigenvector of the Ricci operator.

Examples of strongly locally ϕ -symmetric spaces include the non-Sasakian (κ, μ) -manifolds (Boeckx [1999]). Special cases of these are the non-abelian 3-dimensional unimodular Lie groups with left-invariant contact metric structures (Boeckx, Bueken and Vanhecke [1999]). Boeckx, Bueken and Vanhecke [1999] also gave an example of a non-unimodular Lie group with a weakly locally ϕ -symmetric contact metric structure which is not strongly locally ϕ -symmetric.

To see these last examples explicitly, we include the classification of simply connected homogeneous 3-dimensional contact metric manifolds as given by Perrone [1998]. Let $W = \frac{1}{8}(\tau - \text{Ric}(\xi) + 4)$ denote the Webster scalar curvature (cf. Section 10.4 below). The classification of 3-dimensional Lie groups and their left-invariant metrics was given by Milnor in [1976].

Theorem 7.26 *Let (M^3, η, g) be a simply connected homogeneous contact metric manifold. Then M is a Lie group G and both g and η are left invariant. More precisely, we have the following classification:*

(1) *If G is unimodular, then it is one of the following Lie groups:*

1. *The Heisenberg group when $W = |\mathcal{L}_\xi g| = 0$;*
2. *$SU(2)$ when $4\sqrt{2}W > |\mathcal{L}_\xi g|$;*
3. *the universal covering of the group of rigid motions of the Euclidean plane when $4\sqrt{2}W = |\mathcal{L}_\xi g| > 0$;*
4. *the universal covering of $SL(2, \mathbb{R})$ when $-|\mathcal{L}_\xi g| \neq 4\sqrt{2}W < |\mathcal{L}_\xi g|$;*
5. *the group of rigid motions of the Minkowski plane when $4\sqrt{2}W = -|\mathcal{L}_\xi g| < 0$.*

(2) *If G is non-unimodular, its Lie algebra is given by*

$$[e_1, e_2] = \alpha e_2 + 2\xi, \quad [e_1, \xi] = \gamma e_2, \quad [e_2, \xi] = 0,$$

where $\alpha \neq 0$, $e_1, e_2 = \phi e_1 \in \mathcal{D}$, and $4\sqrt{2}W < |\mathcal{L}_\xi g|$. Moreover, if $\gamma = 0$, the structure is Sasakian and $W = -\frac{\alpha^2}{4}$.

The structures on the unimodular Lie groups in this theorem satisfy the (κ, μ) -nullity condition, and hence they are strongly locally ϕ -symmetric. The weakly locally ϕ -symmetric contact metric structure of Boeckx, Bueken and Vanhecke [1999] that is not strongly locally ϕ -symmetric is the non-unimodular case with $\gamma = 2$.

Notice also in the unimodular case the role played by the invariant $p = 4\sqrt{2}W/|\mathcal{L}_\xi g|$. Moreover, $W = \frac{(2-\mu)}{4}$ and $|\mathcal{L}_\xi g| = 2\sqrt{2}\sqrt{1-\kappa}$; thus, $p = 2 - \mu/2\sqrt{1-\kappa}$, which is the invariant I_M of Boeckx discussed in Section 7.3.

Returning to Boeckx's observation [1999] that a non-Sasakian (κ, μ) -manifold is strongly locally ϕ -symmetric and also locally homogeneous, he recently, [2006], proved a converse.

Theorem 7.27 *Let M be a locally contact homogeneous contact metric manifold. If M is strongly locally ϕ -symmetric, then it is a (κ, μ) -manifold.*

J. Berndt [1997] studied the geometry of the complex Grassmannian $\mathbb{C}G_{2,m}$ of complex 2-planes in \mathbb{C}^{m+2} . Each point in his model of this

Grassmannian represents a closed geodesic in the focal set Q^{m+1} of $\mathbb{C}P^{m+1}$ in $\mathbb{H}P^{m+1}$, and he views the Riemannian submersion $Q^{m+1} \longrightarrow \mathbb{C}G_{2,m}$ as an analogue of the Hopf fibration $S^{2m+1} \longrightarrow \mathbb{C}P^m$. Among Berndt's observations [1997, p. 37] is that Q^{m+1} is a Sasakian globally ϕ -symmetric space and that Q^2 has constant ϕ -sectional curvature equal to 5.

Cho [1999] makes use of the generalized Tanaka connection ${}^*\nabla$ (see Section 10.4) and studies contact metric manifolds satisfying

$$({}^*\nabla_{\dot{\gamma}}R)(\cdot, \dot{\gamma})\dot{\gamma} = 0$$

for any unit ${}^*\nabla$ -geodesic γ (${}^*\nabla_{\dot{\gamma}}\dot{\gamma} = 0$). Cho shows that a (κ, μ) -manifold satisfying this condition is either Sasakian locally ϕ -symmetric, 3-dimensional with $\mu = 0$ and weakly locally ϕ -symmetric, or has $\mu = 2$ and is weakly locally ϕ -symmetric.

Returning to the Sasakian context, Vanhecke and the author [1987a] showed that complete, simply connected globally ϕ -symmetric spaces are naturally reductive homogeneous spaces. Watanabe [1980] showed that a Sasakian locally ϕ -symmetric space is locally homogeneous and analytic. Berndt and Vanhecke [1999] proved that a simply connected Sasakian ϕ -symmetric space is weakly symmetric, i.e., any two points can be interchanged by an isometry. Complete simply connected globally ϕ -symmetric spaces have been classified by Jiménez and Kowalski [1993]. In dimension 3, the classification is the unit sphere S^3 together with the universal covering of $SL(2, \mathbb{R})$, the Heisenberg group, and $SU(2)$, each with a special left-invariant metric (Vanhecke and the author [1987b]). In dimension 5, the classification was obtained by Kowalski and Wegrzynowski [1987]. As a corollary to this classification, Kowalski and Wegrzynowski in dimension 5 and Jiménez and Kowalski in general dimension gave a classification of the complete, simply connected Sasakian space forms. An isospectral problem for locally ϕ -symmetric spaces was studied by Shibuya [1982].

Watanabe [1980] also showed that a 3-dimensional Sasakian manifold with constant scalar curvature is locally ϕ -symmetric. Two extensions of this are the following: A compact regular Sasakian manifold with constant scalar curvature and nonnegative sectional curvature is locally ϕ -symmetric (Perrone and Vanhecke [1991]). A 3-dimensional contact metric manifold with $Q\phi = \phi Q$ is weakly locally ϕ -symmetric if and only

if it is of constant scalar curvature (Koufogiorgos, Sharma and the author [1990]).

We have remarked that the product metric on $S^3 \times S^2$ is not an associated metric. A family of Sasakian globally ϕ -symmetric structures on $S^3 \times S^2$ was constructed by Watanabe and Fujita [1988], and Perrone and Vanhecke [1991] proved that the only 5-dimensional compact, simply connected, homogeneous contact manifolds are diffeomorphic to S^5 or $S^3 \times S^2$.

On a Riemannian locally symmetric space, the local geodesic symmetries are isometries and hence volume-preserving. However, there exist Riemannian manifolds whose local geodesic symmetries are volume preserving but not locally symmetric (see, e.g., Kowalski and Vanhecke [1984]). In dimension 3 (Vanhecke and the author [1987b]; see also Watanabe [1980]) and dimension 5 (Vanhecke and the author [1987c]), Sasakian manifolds whose geodesic symmetries are volume-preserving are locally ϕ -symmetric and conversely.

J. C. González-Dávila, M. C. González-Dávila and L. Vanhecke [1995] considered metrics with respect to which a given vector field ξ is a unit Killing vector field, with emphasis on the case in which the dual form η is a contact form; this is again the context of the R-contact manifolds of Rukimbira [1993]. These authors study the case in which local reflections with respect to the flow lines are isometries; such spaces are called *locally Killing-transversally symmetric spaces*. In particular, they show that a Riemannian manifold with a unit Killing vector field ξ has a natural structure as a Sasakian locally ϕ -symmetric space if and only if it is a locally Killing-transversally symmetric space and $K(X, \xi) = 1$ for all X orthogonal to ξ .

There are a number of other geometric ideas surrounding the subjects of Sasakian space forms and locally ϕ -symmetric spaces, and we will mention a few of these with a few references. One area of interest is the study of reflections and other symmetries; in addition to some of the references already given, we mention Bueken and Vanhecke [1993] which studies reflections in a submanifold. Another area is the study of Jacobi vector fields and their use in the characterization of Sasakian space forms as well as in the study of symmetries. Here we will give only a few references on the use of Jacobi fields: Vanhecke and the author [1987d], Bueken and Vanhecke [1988a], and Vanhecke [1988] for a survey of the techniques involved.

8

Submanifolds of Kähler and Sasakian Manifolds

In this chapter we study submanifolds in both contact and Kähler geometry. These are extensive subjects in their own right, and we give only a few basic results.

8.1 Invariant submanifolds

For a submanifold M of a Riemannian manifold (\tilde{M}, \tilde{g}) we denote the induced metric by g . Then the Levi-Civita connection ∇ of g and the second fundamental form σ are related to the ambient Levi-Civita connection $\tilde{\nabla}$ by

$$\tilde{\nabla}_X Y = \nabla_X Y + \sigma(X, Y).$$

For a normal vector field ν we denote by A_ν the corresponding Weingarten map and we denote by ∇^\perp the connection in the normal bundle; in particular, A_ν and ∇^\perp are defined by

$$\tilde{\nabla}_X \nu = -A_\nu X + \nabla_X^\perp \nu.$$

The Gauss equation is

$$\begin{aligned} \tilde{R}(X, Y, Z, W) &= R(X, Y, Z, W) + \tilde{g}(\sigma(X, Z), \sigma(Y, W)) \\ &\quad - \tilde{g}(\sigma(Y, Z), \sigma(X, W)). \end{aligned}$$

Defining the covariant derivative of σ by $(\nabla'\sigma)(X, Y, Z) = \nabla_X^\perp \sigma(Y, Z) - \sigma(\nabla_X Y, Z) - \sigma(Y, \nabla_X Z)$, the Codazzi equation is

$$(\tilde{R}_{XY}Z)^\perp = (\nabla'\sigma)(X, Y, Z) - (\nabla'\sigma)(Y, X, Z).$$

Finally, for normal vector fields ν and ζ , the equation of Ricci–Kühne is

$$\tilde{R}(X, Y, \nu, \zeta) = R^\perp(X, Y, \nu, \zeta) - g([A_\nu, A_\zeta]X, Y).$$

For a general reference to submanifold theory see Chen [2000].

Let \tilde{M}^{2n} be an almost Hermitian manifold with structure tensors (\tilde{g}, \tilde{J}) . A submanifold M is said to be *invariant* if $\tilde{J}T_p M \subset T_p M$. It is well known that an invariant submanifold of a Kähler manifold is both Kähler and minimal.

For a contact metric manifold \tilde{M}^{2n+1} with structure tensors $(\tilde{\phi}, \tilde{\xi}, \tilde{\eta}, \tilde{g})$, a submanifold M is said to be *invariant* if $\tilde{\phi}T_p M \subset T_p M$. Some authors also require that $\tilde{\xi}$ be tangent to M , but this is a consequence. Clearly $\tilde{\xi}$ cannot be normal on any neighborhood \mathcal{U} of M , for then \mathcal{U} would be an integral submanifold of the contact subbundle \mathcal{D} and hence as we have seen (Section 5.1) would not be invariant by $\tilde{\phi}$. Now if $\tilde{\xi} = U + \nu$, where U is tangent and ν is normal, first note that $\tilde{\phi}\nu$ is normal, since $\tilde{g}(\tilde{\phi}\nu, X) = -\tilde{g}(\nu, \tilde{\phi}X) = 0$. Therefore $0 = \tilde{\phi}\tilde{\xi} = \tilde{\phi}U + \tilde{\phi}\nu$ and hence $\tilde{\phi}U = 0$ and $\tilde{\phi}\nu = 0$. As a result, $U = \tilde{\eta}(U)\tilde{\xi}$ and $\nu = \tilde{\eta}(\nu)\tilde{\xi}$, but both U and ν cannot be collinear with $\tilde{\xi}$.

Clearly an invariant submanifold inherits a contact metric structure by restriction. Moreover, for the induced structure (ϕ, ξ, η, g) we have $h = \tilde{h}|_M$ as well. Also for the second fundamental form we have

$$\sigma(\xi, X) = \tilde{\nabla}_X \xi - \nabla_X \xi = -\tilde{\phi}X - \tilde{\phi}\tilde{h}X - (-\phi X - \phi hX) = 0.$$

Our first result is a theorem of Chinea [1985] and independently of Endo [1985]; we give the proof of Chinea.

Theorem 8.1 *An invariant submanifold M of a contact metric manifold is minimal.*

Proof. By Lemma 7.3 we have

$$(\tilde{\nabla}_X \tilde{\phi})Y + (\tilde{\nabla}_{\tilde{\phi}X} \tilde{\phi})\tilde{\phi}Y = 2\tilde{g}(X, Y)\tilde{\xi} - \tilde{\eta}(Y)(X + \tilde{h}X + \tilde{\eta}(X)\tilde{\xi}),$$

from which

$$\begin{aligned} \nabla_X \phi Y + \sigma(X, \phi Y) - \phi \nabla_X Y - \tilde{\phi} \sigma(X, Y) \\ + \nabla_{\phi X} (-Y + \eta(Y)\xi) - \sigma(\phi X, Y) - \phi \nabla_{\phi X} \phi Y - \tilde{\phi} \sigma(\phi X, \phi Y) \\ = 2\tilde{g}(X, Y)\tilde{\xi} - \tilde{\eta}(Y)(X + \tilde{h}X + \tilde{\eta}(X)\tilde{\xi}). \end{aligned}$$

Taking the normal part, we have

$$\sigma(X, \phi Y) - \sigma(\phi X, Y) - \tilde{\phi}(\sigma(X, Y) + \sigma(\phi X, \phi Y)) = 0.$$

Interchanging X and Y now yields

$$\sigma(\phi X, \phi Y) = -\sigma(X, Y)$$

and hence that M is minimal. ■

Note also that $\sigma(X, \phi Y) = \sigma(\phi X, Y)$. Thus we have $g(A_\nu X, \phi Y) = \tilde{g}(\sigma(X, \phi Y), \nu) = \tilde{g}(\sigma(\phi X, Y), \nu) = g(A_\nu \phi X, Y)$ and hence

$$A_\nu \phi + \phi A_\nu = 0.$$

Moreover, a ϕ -section on M is a $\tilde{\phi}$ -section on \tilde{M} ; the Gauss equation and $\sigma(\phi X, \phi Y) = -\sigma(X, Y)$ yield $H(X) \leq \tilde{H}(X)$, with equality holding everywhere if and only if M is totally geodesic (Endo [1986]).

Theorem 8.2 *If \tilde{M} is a K -contact (resp. Sasakian) manifold and M an invariant submanifold, then M is also K -contact (resp. Sasakian).*

Proof. We have already noted that $h = \tilde{h}|_M$ and hence the K -contact result. Now again as in the last proof,

$$(\tilde{\nabla}_X \tilde{\phi})Y = \nabla_X \phi Y + \sigma(X, \phi Y) - \phi \nabla_X Y - \tilde{\phi} \sigma(X, Y),$$

so if $(\tilde{\nabla}_X \tilde{\phi})Y = \tilde{g}(X, Y)\tilde{\xi} - \tilde{\eta}(Y)X$, we have $(\nabla_X \phi)Y = g(X, Y)\xi - \eta(Y)X$. ■

In [1973a] Harada showed that invariant submanifolds of a compact regular Sasakian manifold respect the Boothby–Wang fibration. For a compact regular Sasakian manifold M , let us denote by M/ξ the base manifold of the Boothby–Wang fibration, which, as we have seen, carries a natural Kähler structure. We can now state Harada’s result.

Theorem 8.3 *Let M be a compact invariant submanifold of a compact regular Sasakian manifold \tilde{M} . Then M is regular and M/ξ is a Kähler (invariant) submanifold of \tilde{M}/ξ .*

For example, consider the complex quadric Q^{n-1} in $\mathbb{C}P^n$ together with the Hopf fibration $S^{2n+1} \longrightarrow \mathbb{C}P^n$. Then the set of fibers over Q^{n-1} form a codimension 2 invariant submanifold of the Sasakian structure on S^{2n+1} ; see Kenmotsu [1969], Kon [1976].

Let us now very briefly recall results of Simons [1968] and of Chern, do Carmo and Kobayashi [1970] on submanifolds of a sphere. The Simons result is that if M^n is a closed minimal submanifold of the sphere $S^{n+p}(1)$ and if $|\sigma|^2 < n/(2 - \frac{1}{p})$, then M^n is totally geodesic. Chern, do Carmo and Kobayashi proved that the only minimal submanifolds of the sphere $S^{n+p}(1)$ with $|\sigma|^2 = n/(2 - \frac{1}{p})$ are pieces of the Clifford minimal hypersurfaces and the Veronese surface. The proofs of these center on the computation of the Laplacian of $|\sigma|^2$. In [1972] Ogiue used these ideas to study complete invariant submanifolds M^{2n} in $\mathbb{C}P^{n+p}(1)$, the complex projective space with the Fubini–Study metric of constant holomorphic curvature 1. Ogiue’s results of [1972] may be summarized as follows: Let M^{2n} be a complete Kähler submanifold of $\mathbb{C}P^{n+p}(1)$. If the holomorphic curvature H of M^{2n} is greater than $\frac{1}{2}$ and the scalar curvature of M^{2n} is constant or if $H > 1 - \frac{n+2}{2(n+2p)}$, then M^{2n} is totally geodesic.

In his excellent survey article [1974], Ogiue obtained other results, posed several conjectures and open problems, and continued this theme in [1976a]. For example, in the context of the above results Ogiue proved in [1976a] that if the sectional curvature K of M^{2n} exceeds $\frac{n+3}{8n}$ or if $K > \frac{1}{8}$ and $H > \frac{1}{2}$, then M^{2n} is totally geodesic. Ogiue’s conjectures from [1974] and [1976a] included that (i) $H > \frac{1}{2}$ or (ii) $K > \frac{1}{8}$ and $n \geq 2$ imply that M^{2n} is totally geodesic. In [1985a] A. Ros introduced a new technique to attack this kind of problem for M^{2n} compact, viz. for unit tangent vectors V , regard $|\sigma(V, V)|^2$ as a function of the unit tangent bundle of M^{2n} and study its behavior at its maximum point. The result of Ros in [1985a] is that Ogiue’s conjecture (i) is true, and using the same technique, Ros and Verstraelen in [1984] proved conjecture (ii). Notice that the Calabi (Veronese) embeddings of $\mathbb{C}P^n(\frac{1}{2})$ into $\mathbb{C}P^{\frac{n(n+3)}{2}}(1)$ show that these conjectures are best possible (see Section 12.7). In fact, in [1985b] Ros gave a complete classification of compact Kähler submanifolds of $\mathbb{C}P^{n+p}(1)$ with $H \geq \frac{1}{2}$; there are seven interesting cases. As an illustration of the

Ros technique we will give in the next section the proof of a theorem of Urbano [1985] on Lagrangian submanifolds of $\mathbb{C}P^n$.

The corresponding problem in Sasakian geometry is to study a compact invariant submanifold M^{2n+1} of a Sasakian space form $\tilde{M}^{2(n+p)+1}(\tilde{c})$ with constant ϕ -sectional curvature $\tilde{c} > -3$. This was first taken up by Harada [1973a], [1973b], Kon [1976], and later by VanLindt, Verheyen and Verstraelen [1986]. Again the later results are the stronger ones and use the Ros technique; in particular, VanLindt, Verheyen and Verstraelen prove that if the ϕ -sectional curvature of M^{2n+1} exceeds $\frac{\tilde{c}+3}{2}$ or if the sectional curvature of M^{2n+1} exceeds $\frac{\tilde{c}+3}{8}$, then M^{2n+1} is totally geodesic.

A well-known classical characterization of complex space forms is the following theorem of Yano and Mogi [1955].

Theorem 8.4 *A Kähler manifold M^{2n} , $n \geq 2$, is a complex space form if and only if for every point and every holomorphic section at the point, there exists a unique totally geodesic holomorphic curve tangent to the given holomorphic section at the point.*

The condition that for every point and for every holomorphic section at the point there exists a totally geodesic surface through the point and tangent to the section is known as the *axiom of holomorphic planes*, and this result has many generalizations. For example, instead of planes one can consider $2k$ -dimensional holomorphic subspaces, $1 \leq k < n$, and instead of totally geodesic submanifolds one can consider umbilical submanifolds (*axiom of holomorphic $2k$ -spheres*), Kassabov [1982] (see also Goldberg and Moskal [1976], where the umbilical submanifolds are assumed to have parallel mean curvature vector).

In Sasakian geometry one has the *axiom of ϕ -holomorphic planes*, which requires for every point and every ϕ -section at the point the existence of a totally geodesic surface tangent to the ϕ -section at the point. K. Ogiue [1964] showed that a Sasakian manifold is a Sasakian space form if and only if it satisfies the axiom of ϕ -holomorphic planes. For the idea of an axiom of ϕ -holomorphic 2-spheres, see Harada [1974a].

8.2 Lagrangian and integral submanifolds

We have already seen in Section 1.2 that the maximum dimension of an isotropic submanifold of a symplectic manifold M^{2n} is n and that an n -dimensional isotropic submanifold is called a Lagrangian submanifold.

In Section 5.1 we saw that the maximum dimension of an integral submanifold of a contact manifold M^{2n+1} is also n . In almost Hermitian geometry, isotropic submanifolds are known as *totally real submanifolds*, since they are characterized by the fact that the almost complex structure maps the tangent space at any point into the normal space at the point.

In the spirit of the results of the last section we mention a few results on compact minimal Lagrangian submanifolds M^n of a complex space form $\tilde{M}^{2n}(c)$. In [1974] Yau considered a totally real minimal surface M^2 in a Kähler surface of constant holomorphic curvature c and proved the following results: (1) If M^2 has genus zero, then M^2 is the standard embedding of $\mathbb{R}P^2$ in $\mathbb{C}P^2$. (2) If M^2 is complete and nonnegatively curved, then it is totally geodesic or flat. (3) If M^2 is complete and nonpositively curved with Gaussian curvature K and if $\frac{c}{4} - K \geq a > 0$ for some constant a , then M^2 is totally geodesic or flat. In [1973] Houh proved that if a totally real minimal surface M^2 in $\mathbb{C}P^2$ has constant scalar normal curvature, it is totally geodesic or is nonpositively curved, and combining this with Yau's result, Houh showed that if in addition M^2 is complete, it is totally geodesic or flat. The flat case is realized by the torus, T^2 , embedded in $\mathbb{C}P^2$ as a flat minimal Lagrangian submanifold; Ludden, Okumura and Yano [1975].

In [1976b] Ogiue showed that if $c > 0$ and the sectional curvature K of M^n satisfies $K > \frac{(n-2)c}{4(2n-1)}$, then M^n is totally geodesic. This was extended by Chen and Houh [1979], who showed that if $c > 0$ and $K \geq \frac{(n-2)c}{4(2n-1)}$, then either M^n is totally geodesic or $n = 2$ and the surface M^2 is flat. If a minimal Lagrangian submanifold in $\tilde{M}^{2n}(c)$ is itself of constant curvature k , then it was shown by Chen and Ogiue [1974a] that the submanifold is either totally geodesic or $k \leq 0$; this was improved upon by Ejiri [1982], who showed that the submanifold is either totally geodesic or flat. Finally, to give a stronger result and to illustrate the technique of A. Ros, we give the following theorem of Urbano [1985] with proof.

Theorem 8.5 *Let M be a compact Lagrangian submanifold minimally immersed in $\mathbb{C}P^n(c)$. If the sectional curvature K of M is greater than 0, then M is totally geodesic.*

Proof. We denote by (J, \tilde{g}) the almost Hermitian structure on $\mathbb{C}P^n(c)$, where \tilde{g} is the Fubini–Study metric of constant holomorphic curvature c .

For tangent vectors X and Y we have readily

$$0 = (\tilde{\nabla}_X J)Y = \tilde{\nabla}_X JY - J\tilde{\nabla}_X Y = -A_{JY}X + \nabla_X^\perp JY - J\nabla_X Y - J\sigma(X, Y),$$

giving

$$A_{JY}X = -J\sigma(X, Y), \quad \nabla_X^\perp JY = J\nabla_X Y.$$

The equations of Ricci–Kühne and Gauss give

$$\begin{aligned} R^\perp(X, Y, JZ, JW) &= \tilde{g}(\tilde{R}_{XY}JZ, JW) + g([A_{JZ}, A_{JW}]X, Y) \\ &= g(R_{XY}Z, W) - \tilde{g}(\sigma(Y, Z), \sigma(X, W)) \\ &\quad + \tilde{g}(\sigma(X, Z), \sigma(Y, W)) + g(A_{JW}X, A_{JZ}Y) \\ &\quad - g(A_{JZ}X, A_{JW}Y) \\ &= g(R_{XY}Z, W). \end{aligned}$$

Since the ambient space is of constant holomorphic curvature, $\tilde{R}_{XY}Z = \frac{c}{4}(g(Y, Z)X - g(X, Z)Y - \Omega(Y, Z)JX + \Omega(X, Z)JY + 2\Omega(X, Y)JZ)$ and hence $(\tilde{R}_{XY}Z)^\perp = 0$. Thus by the Codazzi equation, $(\nabla'\sigma)(X, Y, Z) = \nabla_X^\perp \sigma(Y, Z) - \sigma(\nabla_X Y, Z) - \sigma(Y, \nabla_X Z)$ is symmetric. Defining $\nabla'^2\sigma$ by

$$\begin{aligned} (\nabla'^2\sigma)(X, Y, Z, W) &= \nabla_X^\perp((\tilde{\nabla}\sigma)(Y, Z, W)) \\ &\quad - (\nabla'\sigma)(\nabla_X Y, Z, W) - (\nabla'\sigma)(Y, \nabla_X Z, W) \\ &\quad - (\nabla'\sigma)(Y, Z, \nabla_X W) \end{aligned}$$

we have by the symmetry of $(\nabla'\sigma)(X, Y, Z)$ that

$$\begin{aligned} (\nabla'^2\sigma)(X, Y, Z, W) &= (\nabla'^2\sigma)(Y, X, Z, W) + R_{XY}^\perp \sigma(Z, W) \\ &\quad - \sigma(R_{XY}Z, W) - \sigma(Z, R_{XY}W). \end{aligned}$$

Define a real-valued function on the unit tangent bundle T_1M by $f(V) = \tilde{g}(\sigma(V, V), JV)$. Since T_1M is compact, f attains its maximum at a unit vector V tangent to M at a point p . For any unit tangent vector U at p , let $\gamma(t)$ be the geodesic in M with $\gamma(0) = p$ and $\gamma'(0) = U$. Let $V(t)$ be the parallel vector field along γ with $V(0) = V$. Then

$$0 = \frac{d}{dt}f(V(t))\Big|_{t=0} = \tilde{g}((\nabla'\sigma)(U, V, V), JV).$$

Also

$$\begin{aligned}
0 &\geq \frac{d^2}{dt^2} f(V(t)) \Big|_{t=0} = \tilde{g}((\nabla'^2 \sigma)(U, U, V, V), JV) \\
&= \tilde{g}((\nabla'^2 \sigma)(U, V, V, U), JV) \\
&= \tilde{g}((\nabla'^2 \sigma)(V, U, V, U) + R_U^\perp \sigma(V, U) - \sigma(R_U V, U) \\
&\quad - \sigma(R_U V, U, V), JV) \\
&= \tilde{g}((\nabla'^2 \sigma)(V, V, U, U), JV) + g(R_U V, J\sigma(U, V)) \\
&\quad - g(A_{JU} R_U V, V) - g(A_{JV} R_U V, V) \\
&= \tilde{g}((\nabla'^2 \sigma)(V, V, U, U), JV) + 2g(R_U V, J\sigma(U, V)) \\
&\quad + g(R_U V, J\sigma(V, V)).
\end{aligned}$$

On the other hand, restricting f to the fiber of $T_1 M$ at p and taking U orthogonal to V , we have

$$\begin{aligned}
0 &= Uf = 2\tilde{g}(\sigma(U, V), JV) + \tilde{g}(\sigma(V, V), JU) \\
&= 2g(A_{JV} V, U) + \tilde{g}(\sigma(V, V), JU) = 3\tilde{g}(\sigma(V, V), JU).
\end{aligned}$$

Therefore, since U is any unit vector orthogonal to V ,

$$\sigma(V, V) = f(V)JV \quad \text{or} \quad A_{JV} V = f(V)V,$$

i.e., V is an eigenvector of A_{JV} with eigenvalue $f(V)$. Furthermore,

$$\begin{aligned}
0 &\geq U^2 f = 6\tilde{g}(\sigma(U, V), JU) + 3\tilde{g}(\sigma(V, V), J(-V)) \\
&= 6\tilde{g}(\sigma(U, U), JV) - 3f(V) \\
&= 6g(A_{JV} U, U) - 3f(V).
\end{aligned}$$

Thus if $\{U_1, \dots, U_n\}$ is an orthonormal eigenvector basis of A_{JV} with $U_n = V$ and if $\lambda_i, i = 1, \dots, n-1$, are the other eigenvalues, then

$$f(V) - 2\lambda_i \geq 0.$$

Now from the above and the minimality, we have

$$\begin{aligned}
 0 &\geq \sum_{i=1}^n \left\{ \tilde{g}((\tilde{\nabla}^2 \sigma)(V, V, U_i, U_i), JV) \right. \\
 &\quad \left. + 2g(R_{U_i V} V, J\sigma(U_i, V)) + g(R_{U_i V} U_i, J\sigma(V, V)) \right\} \\
 &= \sum_{i=1}^{n-1} \left\{ -2\lambda_i K(V, U_i) + f(V)K(V, U_i) \right\} \\
 &= \sum_{i=1}^{n-1} K(V, U_i) \{f(V) - 2\lambda_i\}.
 \end{aligned}$$

Finally, since the sectional curvature is positive, we conclude that each λ_i equals $\frac{1}{2}f(V)$ and hence $\text{tr} A_{JV} = \frac{(n+1)f(V)}{2} = 0$, giving $f(V) = 0$. Now $f(-U) = -f(U)$ and V was the maximum point of f , so $f = 0$. Thus $\sigma(V, V)$ is orthogonal to JV and to JU for any $U \perp V$, and hence $\sigma(V, V) = 0$ for any vector V , and so M must be totally geodesic. ■

In Section 1.2 and Example 5.3.3 we remarked that there is no topological embedding of a sphere as a Lagrangian submanifold of \mathbb{C}^n and no umbilical, non-totally-geodesic, Lagrangian submanifolds of a complex space-form. The immersed Whitney sphere was the closest candidate with only one double point. Its second fundamental form σ was given by

$$\sigma(X, Y) = \frac{n}{n+2} (\tilde{g}(X, Y) \mathbf{H} + \tilde{g}(JX, \mathbf{H}) JY + \tilde{g}(JY, \mathbf{H}) JX),$$

where \mathbf{H} denotes the mean curvature vector (Borrelli, Chen and Morvan [1995], Ros and Urbano [1998]).

More generally, a Lagrangian submanifold of a Kähler manifold is said to be *Lagrangian H-umbilical* if the second fundamental form σ is of the form

$$\begin{aligned}
 \sigma(X, Y) &= \alpha \tilde{g}(JX, \mathbf{H}) \tilde{g}(JY, \mathbf{H}) \mathbf{H} \\
 &\quad + \beta \tilde{g}(\mathbf{H}, \mathbf{H}) (\tilde{g}(X, Y) \mathbf{H} + \tilde{g}(JX, \mathbf{H}) JY + \tilde{g}(JY, \mathbf{H}) JX)
 \end{aligned}$$

for suitable functions α and β . This notion was introduced and a classification of such submanifolds in complex space forms was given by Chen in a series of papers [1997a,b], [1998].

In the contact case, integral submanifolds are often called *C-totally real submanifolds*, since ϕ maps the tangent space at any point into the

normal space. However, we will not adopt this term here. As a reminder, since $\eta(X) = 0$ for any vector X tangent to the integral submanifold, ξ is a normal vector field.

Examples that we might mention here are $S^n \subset S^{2n+1}$ as a totally geodesic integral submanifold as was described in Example 5.3.1 and $T^2 \subset S^5$ as a flat minimal integral submanifold as was described in Example 5.3.2.

We begin our discussion of integral submanifolds with the following lemma.

Lemma 8.1 *Let M be an integral submanifold of a K -contact manifold \tilde{M} . Then $A_\xi = 0$.*

Proof.

$$g(A_\xi X, Y) = \tilde{g}(\sigma(X, Y), \xi) = \tilde{g}(\tilde{\nabla}_X Y, \xi) = -\tilde{g}(Y, \tilde{\nabla}_X \xi) = \tilde{g}(Y, \phi X) = 0.$$

■

Now for an integral submanifold M^n of a Sasakian manifold \tilde{M}^{2n+1} , let $\{e_1, \dots, e_n\}$ be a local orthonormal basis on M^n . Then $\{\phi e_1, \dots, \phi e_n, \xi\}$ is an orthonormal basis of the normal space at each point of the local domain. For simplicity we write A_i for $A_{\phi e_i}$.

Lemma 8.2 *Let M be an integral submanifold of a Sasakian manifold \tilde{M} . Then $A_i e_j = A_j e_i$.*

Proof.

$$\begin{aligned} g(A_i e_j, e_k) &= \tilde{g}(\sigma(e_j, e_k), \phi e_i) = \tilde{g}(\tilde{\nabla}_{e_k} e_j, \phi e_i) \\ &= -\tilde{g}(e_j, (\tilde{\nabla}_{e_k} \phi) e_i + \phi \tilde{\nabla}_{e_k} e_i) = \tilde{g}(\tilde{\nabla}_{e_k} e_i, \phi e_j) = g(A_j e_i, e_k). \end{aligned}$$

■

Now let M^n be an integral submanifold of a Sasakian space form $\tilde{M}(c)$; the Gauss equation yields

$$\begin{aligned} g(R_{XY}Z, W) &= \frac{c+3}{4}(g(X, W)g(Y, Z) - g(X, Z)g(Y, W)) \\ &\quad + \sum_{i=1}^n (g(A_i X, W)g(A_i Y, Z) - g(A_i X, Z)g(A_i Y, W)) \end{aligned}$$

by virtue of Lemma 8.1. In turn, the sectional curvature $K(X, Y)$ of M^n determined by an orthonormal pair $\{X, Y\}$ is given by

$$K(X, Y) = \frac{c+3}{4} + \sum_{i=1}^n (g(A_i X, X)g(A_i Y, Y) - g(A_i X, Y)^2).$$

Moreover, the Ricci tensor ρ and the scalar curvature τ of M^n are given by

$$\begin{aligned} \rho(X, Y) &= \frac{n-1}{4}(c+3)g(X, Y) + \sum_{i=1}^n (\operatorname{tr} A_i)g(A_i X, Y) - \sum_{i=1}^n g(A_i X, A_i Y), \\ \tau &= \frac{n(n-1)}{4}(c+3) + \sum_{i=1}^n (\operatorname{tr} A_i)^2 - |\sigma|^2. \end{aligned}$$

From these expressions the following proposition is not difficult, and we omit the proof.

Proposition 8.1 *Let M^n be an integral submanifold of a Sasakian space form $\tilde{M}^{2n+1}(c)$ that is minimally immersed. Then the following are equivalent:*

- (a) M^n is totally geodesic.
- (b) M^n is of constant curvature $\frac{c+3}{4}$.
- (c) $\rho = \frac{n-1}{4}(c+3)g$.
- (d) $\tau = \frac{n(n-1)}{4}(c+3)$.

Similar to the result of Chen and Ogiue [1974a] in the Kähler case, Yamaguchi, Kon and Ikawa [1976] proved the following result.

Theorem 8.6 *Let M^n be a minimal integral submanifold of a Sasakian space form $\tilde{M}^{2n+1}(c)$. If M^n has constant curvature k , then either M^n is totally geodesic or $k \leq 0$.*

For the standard Sasakian structure on $S^5(1)$ we give the following theorem of Yamaguchi, Kon and Miyahara [1976], and we include a proof, since its techniques, though not new, are different from those presented so far in this book.

Theorem 8.7 *Let M^2 be a complete integral surface of $S^5(1)$ that is minimally immersed. If the Gaussian curvature K of M^2 is ≤ 0 , then M^2 is flat.*

Proof. Choose a system of isothermal coordinates (x^1, x^2) so that the induced metric g is given by $g = E((dx^1)^2 + (dx^2)^2)$. Let $X_i = \frac{\partial}{\partial x^i}$ and $\sigma_{ij} = \sigma(X_i, X_j)$. Recall the standard formulas for the induced connection:

$$\nabla_{X_1} X_1 = -\nabla_{X_2} X_2 = \frac{X_1 E}{2E} X_1 - \frac{X_2 E}{2E} X_2, \quad \nabla_{X_1} X_2 = \frac{X_2 E}{2E} X_1 + \frac{X_1 E}{2E} X_2.$$

Using the minimality and the Codazzi equation, one readily obtains

$$\nabla_{X_1}^\perp \sigma_{12} - \nabla_{X_2}^\perp \sigma_{11} = 0, \quad \nabla_{X_1}^\perp \sigma_{11} + \nabla_{X_2}^\perp \sigma_{12} = 0.$$

Now define a complex-valued function F by

$$F = \tilde{g}(\sigma_{11}, \phi X_1) - i\tilde{g}(\sigma_{12}, \phi X_1).$$

Note that F is nowhere zero on M^2 , for if $F = 0$ at some point m , then $\tilde{g}(\sigma_{11}, \phi X_1)$ and $\tilde{g}(\sigma_{12}, \phi X_1)$ vanish at m , but by the minimality $\tilde{g}(\sigma_{22}, \phi X_1) = -\tilde{g}(\sigma_{11}, \phi X_1) = 0$ and by Lemma 8.2 $\tilde{g}(\sigma_{12}, \phi X_2) = \tilde{g}(\sigma_{22}, \phi X_1) = 0$ and $-\tilde{g}(\sigma_{22}, \phi X_2) = \tilde{g}(\sigma_{11}, \phi X_2) = \tilde{g}(\sigma_{12}, \phi X_1) = 0$. Thus σ vanishes at m , and so by the Gauss equation, the Gaussian curvature at m is $+1$, contradicting the hypothesis $K \leq 0$.

Differentiating the real part of F with respect to X_1 , we have

$$\begin{aligned} X_1 \Re F &= \tilde{g}(\nabla_{X_1}^\perp \sigma_{11}, \phi X_1) + \tilde{g}\left(\sigma_{11}, E\xi + \frac{X_1 E}{2E} \phi X_1 - \frac{X_2 E}{2E} \phi X_2\right) \\ &= -\tilde{g}(\nabla_{X_2}^\perp \sigma_{12}, \phi X_1) + \tilde{g}\left(\sigma_{11}, \frac{X_1 E}{2E} \phi X_1 - \frac{X_2 E}{2E} \phi X_2\right). \end{aligned}$$

Differentiating with respect to X_2 and making similar calculations for the imaginary part of F , Lemma 8.2 and the minimality yield $X_1 \Re F = X_2 \Im F$ and $X_2 \Re F = -X_1 \Im F$. Thus F is analytic and therefore $\log |F|^2$ is harmonic.

Now $|F|^2 = \tilde{g}(\sigma_{11}, \phi X_1)^2 + \tilde{g}(\sigma_{12}, \phi X_1)^2$. On the other hand, the Gauss equation gives the Gaussian curvature K as

$$K = 1 + \frac{1}{E^2}(\tilde{g}(\sigma_{11}, \sigma_{22}) - \tilde{g}(\sigma_{12}, \sigma_{12})) = 1 - \frac{2}{E^3}|F|^2.$$

Thus $|F|^2 = E^3 \left(\frac{1-K}{2}\right)$. Note also the classical formula for the Gaussian curvature of g , namely $K = \frac{-1}{6E} \Delta \log E^3$.

Suppose now that the Gaussian curvature of M^2 is nonpositive. Then

$$\Delta \log \frac{|F|^2}{E^3} = -\Delta \log E^3 = 6EK \leq 0 \quad (*)$$

and

$$\log \frac{|F|^2}{E^3} = \log \frac{1-K}{2} \geq \log \frac{1}{2}.$$

Thus $-\log \frac{|F|^2}{E^3}$ is a subharmonic function which is bounded above.

Now define a metric g^* on M^2 by $g^* = |F|((dx^1)^2 + (dx^2)^2)$; its Gaussian curvature is $-\frac{1}{4|F|} \Delta \log |F|^2 = 0$. That is, g^* is a flat metric on M^2 that is conformally equivalent to g , and hence the universal covering surface \tilde{M} of M^2 is conformally equivalent to the Euclidean plane. Thus \tilde{M} is a parabolic surface; but every subharmonic function that is bounded above on a parabolic surface is a constant. Therefore $-\log \frac{|F|^2}{E^3}$, lifted to \tilde{M} , is a constant, and hence it is constant on M^2 . Equation (*) now gives $K = 0$. ■

Combining Theorems 8.6 and 8.7, we have the following corollary.

Corollary 8.1 *A complete integral surface of $S^5(1)$ with constant curvature that is minimally immersed has constant curvature 0 or +1.*

Turning to curvature conditions in the spirit of the Urbano result above, we have the following result of VanLindt, Verheyen and Verstraelen [1986].

Theorem 8.8 *Let M^n be a compact integral submanifold minimally immersed in a Sasakian space form $M^{2n+1}(c)$ with $c > -3$. If $K > 0$, then M^n is totally geodesic.*

If the sectional curvature is only ≥ 0 , one can do better in dimension 7; namely, we have the following result of Dillen and Vrancken [1989].

Theorem 8.9 *Let M^3 be a compact integral submanifold of the standard Sasakian structure on $S^7(1)$ that is minimally immersed. If $K \geq 0$, then either M^3 is totally geodesic, M^3 is a covering of the 3-torus or M^3 is a covering of $S^1(\sqrt{3}) \times S^2(\frac{\sqrt{3}}{2})$.*

In dimension 5 the third case does not have an analogue, and the corresponding result was given by Verstraelen and Vrancken [1988]. For the

example of $S^1(\sqrt{3}) \times S^2(\frac{\sqrt{3}}{2})$ in the above theorem, the sectional curvatures satisfy $0 \leq K \leq \frac{4}{3}$, where both extremal values are attained (Dillen and Vrancken [1990]). Restricting the curvature to $0 \leq K \leq 1$, Dillen and Vrancken proved in [1990] the following theorem.

Theorem 8.10 *If M^n is a compact minimal integral submanifold of $S^{2n+1}(1)$ and if $0 \leq K \leq 1$, then K is identically 0 or 1.*

Other conditions on integral submanifolds of Sasakian space forms that have been considered include the notions of the mean curvature vector \mathbf{H} and second fundamental form being C -parallel. That is, $\nabla_X^\perp \mathbf{H}$ is parallel to ξ for all tangent vectors X , and respectively, $(\nabla' \sigma)(X, Y, Z)$ is parallel to ξ for all tangent vectors X, Y, Z . Recall also that a curve $\gamma(s)$ parametrized by arc length in a Riemannian manifold is a *Frenet curve* of osculating order r if there exist orthonormal vector fields E_1, E_2, \dots, E_r along γ , such that

$$\dot{\gamma} = E_1, \quad \nabla_{\dot{\gamma}} E_1 = k_1 E_2, \quad \nabla_{\dot{\gamma}} E_2 = -k_1 E_1 + k_2 E_3,$$

$$\nabla_{\dot{\gamma}} E_{r-1} = -k_{r-2} E_{r-2} + k_{r-1} E_r, \quad \nabla_{\dot{\gamma}} E_r = -k_{r-1} E_{r-1},$$

where k_1, k_2, \dots, k_{r-1} are positive C^∞ functions of s . The function k_j is called the j th curvature of γ . So, for example, a geodesic is a Frenet curve of osculating order 1, a circle is a Frenet curve of osculating order 2 with k_1 a constant; a helix of order r is a Frenet curve of osculating order r such that k_1, k_2, \dots, k_{r-1} are constants. With these ideas in mind, C. Baikoussis and the author [1992] proved the following theorem.

Theorem 8.11 *Let M^2 be an integral surface of a Sasakian space form $\tilde{M}^5(c)$. If the mean curvature vector \mathbf{H} is C -parallel, then either M^2 is minimal or locally the Riemannian product of two curves as follows: (i) a helix of order 4 and a geodesic or helix of order 3, (ii) a helix of order 3 and a geodesic or helix of order 3, or (iii) a circle and a geodesic or helix of order 3.*

On the other hand, Baikoussis, Koufogiorgos and the author [1995] obtained the following result.

Theorem 8.12 *Let M^3 be an integral submanifold of a Sasakian space form $\tilde{M}^7(c)$ with C -parallel second fundamental form. Then locally M^3 is either flat or totally geodesic or a product $\gamma \times M^2$, where γ is a curve and M^2 is a surface of constant curvature and also has C -parallel second fundamental form.*

This paper of Baikoussis, Koufogiorgos and the author includes for the unit sphere, $S^7(1)$, an explicit representation of the flat case. Recently, Fetcu and Oniciuc [to appear] obtained an explicit expression of the flat case for $S^7(c)$, the Sasakian space form with $-3 < c < 1$ (see Example 7.8.1).

We now give an example of this theorem in the case of ϕ -sectional curvature $c < -3$. In Example 7.8.3 we saw that the product $B^n \times \mathbb{R}$, where B^n is a simply connected bounded domain in \mathbb{C}^n , has a Sasakian structure of constant ϕ -sectional curvature $c < -3$. Now take B^n to be the unit ball in \mathbb{C}^n with the metric

$$d\bar{s}^2 = 4 \frac{(1 - \sum |z_i|^2)(\sum dz_i d\bar{z}_i) + (\sum \bar{z}_i dz_i)(\sum z_i d\bar{z}_i)}{(1 - \sum |z_i|^2)^2}.$$

This metric has constant holomorphic curvature -1 , and the fundamental 2-form Ω of the Kähler structure is given by

$$\begin{aligned} \Omega = d\omega &= -4i \frac{(1 - \sum |z_j|^2)(\sum dz_j \wedge d\bar{z}_j) + (\sum \bar{z}_j dz_j) \wedge (\sum z_j d\bar{z}_j)}{(1 - \sum |z_j|^2)^2} \\ &= d \left(Re \frac{4i \sum \bar{z}_j dz_j}{1 - \sum |z_j|^2} \right). \end{aligned}$$

Then $B^n \times \mathbb{R}$ with contact form $\eta = \omega + dt$ and metric $ds^2 = d\bar{s}^2 + \eta \otimes \eta$ is a Sasakian space form with constant ϕ -sectional curvature $c = -4$.

It is well known that setting the imaginary part of z_j equal to zero gives an embedding of real hyperbolic space H^n of constant curvature $-\frac{1}{4}$ as a totally real, totally geodesic submanifold of B^n . To construct a nontrivial example of M^3 as an integral submanifold of $\bar{M}^7(-4) = B^3 \times \mathbb{R}$ with C -parallel second fundamental form, we consider an umbilical surface in H^3 , and as t varies in \mathbb{R} we rotate H^3 so that the surface will trace out the desired M^3 . For this purpose it will be convenient to use the polar form $r_j e^{i\theta_j}$ of z_j . Then η, ξ and ds^2 become

$$\eta = 4 \frac{\sum r_j^2 d\theta_j}{1 - \sum r_j^2} + dt, \quad \xi = \frac{\partial}{\partial t},$$

and

$$ds^2 = \frac{(1 - \sum r_j^2)(\sum(dr_j^2 + r_j^2 d\theta_j^2)) + (\sum r_j dr_j)^2 + (\sum r_j^2 d\theta_j)^2}{(1 - \sum r_j^2)^2} + 16 \frac{(\sum r_j^2 d\theta_j)^2}{(1 - \sum r_j^2)^2} + 8 \frac{\sum r_j^2 d\theta_j dt}{1 - \sum r_j^2} + dt.$$

We remark that the induced metric on H^3 in B^3 defined by $\theta_j =$ constant is the Beltrami–Klein model and not the Poincaré model of hyperbolic space. In this model an umbilical submanifold is in general an ellipsoid. With this in mind we construct our example as follows. On the Sasakian space form $\bar{M}^7(-4) = B^3 \times \mathbb{R}$ we designate the coordinates by $(x_1, \dots, x_7) = (r_1, r_2, r_3, \theta_1, \theta_2, \theta_3, t)$. Now define $\iota : M \rightarrow \bar{M}^7(-4)$ by

$$(r_1, r_2, t) \mapsto \left(r_1, r_2, b\sqrt{1 - r_1^2 - r_2^2}, 0, 0, -\frac{1 - b^2}{4b^2}t, t \right),$$

where b is a constant such that $b^2 < 1$. It is easy to see that $\eta(\iota_*\partial_1) = \eta(\iota_*\partial_2) = \eta(\iota_*\partial_7) = 0$. Thus M is a 3-dimensional integral submanifold of the Sasakian space form $\bar{M}^7(-4)$. Direct computation then shows that M has C -parallel second fundamental form (see Baikoussis, Koufogiorgos and the author [1995] for details).

In [1973] Chen and Ogiue introduced the *axiom of antiholomorphic k -planes* on a Kähler manifold M^{2n} , $n \geq 2$. The axiom requires that for every point and every totally real k -plane at the point, there exists a k -dimensional totally geodesic submanifold tangent to the k -plane at the point. The main result of Chen and Ogiue is that a Kähler manifold M^{2n} , $n \geq 3$, is a complex space form if and only if it satisfies the axiom of antiholomorphic 2-planes.

In [2001] Kirichenko took a somewhat different approach to this problem and introduced the *axiom of totally real k -planes* for an almost Hermitian manifold M^{2n} . The axiom requires that for every point and every totally real k -plane at the point, there exists a k -dimensional totally real, totally geodesic submanifold tangent to the k -plane at the point. Kirichenko is then able to extend the axiomatic characterization of complex space forms to the almost Kähler (symplectic) setting.

Theorem 8.13 *An almost Kähler manifold M^{2n} , $n \geq 3$, is a complex space form if and only if it satisfies the axiom of totally real n -planes (Lagrangian).*

Similarly, in contact geometry one has these two settings. A Sasakian manifold satisfies the *axiom of anti- ϕ -holomorphic planes* if for every point and every 2-plane at the point that is annihilated by η and $d\eta$, there exists a totally geodesic surface tangent to the plane at the point (Harada [1974b]). Harada proves that a Sasakian manifold of dimension ≥ 7 is a Sasakian space form if and only if it satisfies the axiom of anti- ϕ -holomorphic planes. On the other hand, Kirichenko and Borisovski [1998] introduced an axiom called the *geodesic integrability* (of the contact subbundle) for a contact metric manifold M^{2n+1} . The idea is to require that for every point and every n -dimensional linear subspace at the point that is annihilated by η and $d\eta$, there exists a unique n -dimensional, totally geodesic, integral submanifold tangent to the subspace at the point. With this notion Kirichenko and Borisovski can extend the Harada idea to the K-contact case. Specifically, they prove that a K-contact manifold of dimension ≥ 7 is a Sasakian space form if and only if it is geodesically integrable.

9

Tangent Bundles and Tangent Sphere Bundles

In the first two sections of this chapter we discuss the geometry of the tangent bundle and the tangent sphere bundle. In Section 3 we briefly present a more general construction on vector bundles and in Section 4 specialize to the case of the normal bundle of a submanifold. The formalism for the tangent bundle and the tangent sphere bundle is of sufficient importance to warrant its own development, rather than specializing from the vector bundle case. In Section 5 we discuss briefly a contact structure on the projectivized tangent bundle of the hyperbolic plane and the geodesic flow on this bundle rather than on the unit tangent bundle, which is its usual setting.

9.1 Tangent bundles

In Chapter 1 we saw that the cotangent bundle of a manifold has a natural symplectic structure, and we will see here that the same is true of the tangent bundle of a Riemannian manifold. Moreover, the tangent bundle carries a natural Riemannian metric, called the Sasaki metric. We give the connection and curvature of this metric and the implication of the metric being locally symmetric or conformally flat.

Let M be an $(n+1)$ -dimensional C^∞ manifold and $\bar{\pi} : TM \longrightarrow M$ its tangent bundle. If (x^1, \dots, x^{n+1}) are local coordinates on M , set $q^i = x^i \circ \bar{\pi}$; then (q^1, \dots, q^{n+1}) together with the fiber coordinates (v^1, \dots, v^{n+1}) form local coordinates on TM .

If X is a vector field on M , its *vertical lift* X^V on TM is the vector field defined by $X^V \omega = \omega(X) \circ \bar{\pi}$, where ω is a 1-form on M , which on the left side of this equation is regarded as a function on TM .

For an affine connection D on M , the *horizontal lift* X^H of X is defined by $X^H \omega = D_X \omega$. The covariant derivative $D_X \omega$ has local expression $(X^i \frac{\partial \omega_j}{\partial x^i} - X^i \omega_k \Gamma_{ij}^k) dx^j$, where the Γ_{ij}^k 's are the connection coefficients. If we evaluate $D_X \omega$ on a vector $t = v^l \frac{\partial}{\partial x^l}$, we have easily

$$(D_X \omega)(t) = v^j X^i \frac{\partial \omega_j}{\partial x^i} - X^i v^j \Gamma_{ij}^k \omega_k = \left(X^i \frac{\partial}{\partial q^i} - X^i v^j \Gamma_{ij}^k \frac{\partial}{\partial v^k} \right) \omega_l v^l.$$

Thus the local expression for X^H is

$$X^H = X^i \frac{\partial}{\partial q^i} - X^i v^j \Gamma_{ij}^k \frac{\partial}{\partial v^k}.$$

The span of the horizontal lifts at $t \in TM$ is called the *horizontal subspace* of $T_t TM$. The *connection map* $K : TTM \longrightarrow TM$ is defined by

$$KX^H = 0, \quad KX_t^V = X_{\bar{\pi}(t)}, \quad t \in TM.$$

The connection map K may also be defined in the following way. Given $X \in T_t TM$, let γ be a smooth curve in TM with tangent vector X at $t = \gamma(0)$. Let $\alpha = \bar{\pi} \circ \gamma$ be the projection of the curve to M . Then

$$KX = D_{\dot{\alpha}} \gamma|_0,$$

and the curve γ in TM is horizontal if γ , viewed as a vector field along α , is parallel. The horizontal subspace can also be defined by

$$\{(s \circ \alpha)_*(0) \in T_t TM \mid \alpha \text{ path in } M, s \text{ section of } TM, s(\alpha(0)) = t, D_{\dot{\alpha}(0)} s = 0\}.$$

It is immediate that $[X^V, Y^V] \omega = 0$. Furthermore,

$$\begin{aligned} [X^H, Y^V] &= \left[X^i \frac{\partial}{\partial q^i} - X^i v^j \Gamma_{ij}^k \frac{\partial}{\partial v^k}, Y^l \frac{\partial}{\partial v^l} \right] \\ &= \left(X^i \frac{\partial Y^k}{\partial x^i} + X^i Y^l \Gamma_{il}^k \right) \frac{\partial}{\partial v^k} = (D_X Y)^V. \end{aligned}$$

Similarly, denoting the curvature tensor of D on M by \mathbf{R} we have at the point $t \in TM$,

$$[X^H, Y^H]_t = [X, Y]_t^H - (\mathbf{R}_{X \cdot Y} t)^V.$$

TM admits an almost complex structure J defined by

$$JX^H = X^V, \quad JX^V = -X^H.$$

Using the above expressions for the Lie brackets in the Nijenhuis torsion of J one can easily see that J is integrable if and only if D has vanishing curvature and torsion (Hsu [1960], Dombrowski [1962]).

If now G is a Riemannian metric on M and D its Levi-Civita connection, we define a Riemannian metric \bar{g} on TM called the *Sasaki metric*, Sasaki [1958] (not to be confused with a Sasakian structure), by

$$\bar{g}(X, Y) = (G(\bar{\pi}_* X, \bar{\pi}_* Y) + G(KX, KY)) \circ \bar{\pi},$$

where X and Y are vector fields on TM . Since $\bar{\pi}_* \circ J = -K$ and $K \circ J = \bar{\pi}_*$, \bar{g} is Hermitian for the almost complex structure J .

On TM define the *Liouville form* β by $\beta(X)_t = G(t, \bar{\pi}_* X)$, $t \in TM$, or equivalently by the local expression $\beta = \sum G_{ij} v^i dq^j$. Then $d\beta$ is a symplectic structure on TM , and in particular, $2d\beta$ is the fundamental 2-form of the almost Hermitian structure (J, \bar{g}) . To see this, first note that since $\beta(X^V) = 0$ and $[X^V, Y^V] = 0$, $2d\beta(X^V, Y^V) = 0$. Similarly

$$\begin{aligned} 2d\beta(X^V, Y^H)_t &= X^V(G(t, \bar{\pi}_* Y^H) \circ \bar{\pi}) = X^V \left(G \left(v^l \frac{\partial}{\partial x^l}, Y \right) \circ \bar{\pi} \right) \\ &= G(X, Y) = \bar{g}(X^V, Y^V) = \bar{g}(X^V, JY^H). \end{aligned}$$

Now choose a vector field Z on M such that $Z_m = t$ and $(D_X Z)_m = 0$ for all X . Then

$$\begin{aligned} 2d\beta(X^H, Y^H)_t &= (X^H(G(Z, Y) \circ \bar{\pi}) - Y^H(G(Z, X) \circ \bar{\pi}) - G(t, [X, Y]))_m \\ &= G(t, D_X Y)_m - G(t, D_Y X)_m - G(t, [X, Y])_m = 0. \end{aligned}$$

Thus TM has an almost Kähler structure which is Kählerian if and only if (M, G) is flat (Tachibana and Okumura [1962]).

As before, we let \mathbf{R} denote the curvature tensor of D , which is now the Levi-Civita connection of G . The Levi-Civita connection $\bar{\nabla}$ of \bar{g} and

its curvature tensor \bar{R} were computed by Kowalski [1971]. These are given by the following formulas, and we will give a partial proof as an illustration:

$$\begin{aligned}(\bar{\nabla}_{X^H} Y^H)_t &= (D_X Y)_t^H - \frac{1}{2}(\mathbf{R}_{X^H Y}^H)_t^V, \\(\bar{\nabla}_{X^H} Y^V)_t &= \frac{1}{2}(\mathbf{R}_{t Y} X)_t^H + (D_X Y)_t^V, \\(\bar{\nabla}_{X^V} Y^H)_t &= \frac{1}{2}(\mathbf{R}_{t X} Y)_t^H, \\ \bar{\nabla}_{X^V} Y^V &= 0.\end{aligned}$$

For example, from $2\bar{g}(\bar{\nabla}_X Y, W) = X\bar{g}(Y, W) + Y\bar{g}(X, W) - W\bar{g}(X, Y) + \bar{g}([X, Y], W) + \bar{g}([W, X], Y) - \bar{g}([Y, W], X)$ we have

$$\begin{aligned}2\bar{g}(\bar{\nabla}_{X^H} Y^V, W^V) &= X^H \bar{g}(Y^V, W^V) + \bar{g}((D_X Y)^V, W^V) \\ &\quad - \bar{g}((D_X W)^V, Y^V) \\ &= \bar{g}((D_X Y)^V, W^V) + G(D_X Y, W) \circ \bar{\pi} \\ &= 2\bar{g}((D_X Y)^V, W^V)\end{aligned}$$

and

$$\begin{aligned}2\bar{g}(\bar{\nabla}_{X^H} Y^V, W^H)_t &= Y^V \bar{g}(X^H, W^H) + \bar{g}((\mathbf{R}_{X^H W})^V, Y^V) \\ &= G(\mathbf{R}_{t Y} X, W) \circ \bar{\pi} = \bar{g}((\mathbf{R}_{t Y} X)^H, W^H),\end{aligned}$$

giving the formula for $(\bar{\nabla}_{X^H} Y^V)_t$.

Turning to the curvature, we have

$$\begin{aligned}\bar{R}_{X^V Y^V} Z^V &= 0, \\(\bar{R}_{X^V Y^V} Z^H)_t &= \left(\mathbf{R}_{X^H Y} Z + \frac{1}{4} \mathbf{R}_{t X} \mathbf{R}_{t Y} Z - \frac{1}{4} \mathbf{R}_{t Y} \mathbf{R}_{t X} Z \right)_t^H, \\(\bar{R}_{X^H Y^V} Z^V)_t &= - \left(\frac{1}{2} \mathbf{R}_{Y Z} X + \frac{1}{4} \mathbf{R}_{t Y} \mathbf{R}_{t Z} X \right)_t^H, \\(\bar{R}_{X^H Y^V} Z^H)_t &= \frac{1}{2}((D_X \mathbf{R})_{t Y} Z)_t^H + \left(\frac{1}{2} \mathbf{R}_{X Z} Y + \frac{1}{4} \mathbf{R}_{\mathbf{R}_{t Y} Z X} t \right)_t^V,\end{aligned}$$

$$\begin{aligned}
(\bar{R}_{X^H Y^H} Z^V)_t &= \frac{1}{2}((D_X \mathbf{R})_t Z^Y - (D_Y \mathbf{R})_t Z^X)_t^H \\
&\quad + \left(\mathbf{R}_{X Y} Z + \frac{1}{4} \mathbf{R}_{\mathbf{R}_t Z Y} X^t - \frac{1}{4} \mathbf{R}_{\mathbf{R}_t Z X} Y^t \right)_t^V, \\
(\bar{R}_{X^H Y^H} Z^H)_t &= \left(\mathbf{R}_{X Y} Z + \frac{1}{4} \mathbf{R}_{\mathbf{R}_t Z Y} X + \frac{1}{4} \mathbf{R}_{\mathbf{R}_t Z X} Y + \frac{1}{2} \mathbf{R}_{\mathbf{R}_t X Y} Z \right)_t^H \\
&\quad + \frac{1}{2}((D_Z \mathbf{R})_{X Y} t)_t^V.
\end{aligned}$$

We will prove the fourth of these formulas. Recall that we may write the point t as $v^l \frac{\partial}{\partial x^l}$; this is important when we have to differentiate with respect to position. Also we abbreviate $\frac{\partial}{\partial x^i}$ by ∂_i :

$$\begin{aligned}
\bar{R}_{X^H Y^V} Z^H &= \bar{\nabla}_{X^H} \frac{1}{2} v^l (\mathbf{R}_{\partial_l Y} Z)^H \\
&\quad - \bar{\nabla}_{Y^V} ((D_X Z)^H - \frac{1}{2} v^i (\mathbf{R}_{X Z} \partial_i)^V) - \bar{\nabla}_{(D_X Y)^V} Z^H.
\end{aligned}$$

Therefore

$$\begin{aligned}
(\bar{R}_{X^H Y^V} Z^H)_t &= -\frac{1}{2} X^i v^j \Gamma_{ij}^k (\mathbf{R}_{\partial_k Y} Z)^H \\
&\quad + \frac{1}{2} v^l ((D_X \mathbf{R}_{\partial_l Y} Z)_t^H - \frac{1}{2} (\mathbf{R}_{X \mathbf{R}_{\partial_l Y} Z} t)_t^V) \\
&\quad - \frac{1}{2} (\mathbf{R}_{t Y} D_X Z)_t^H + \frac{1}{2} (\mathbf{R}_{X Z} Y)_t^V - \frac{1}{2} (\mathbf{R}_{t D_X Y} Z)_t^H,
\end{aligned}$$

from which the fourth curvature formula readily follows.

The main result of Kowalski [1971] is the following theorem.

Theorem 9.1 *The tangent bundle TM with the Sasaki metric \bar{g} is locally symmetric if and only if the base manifold (M, G) is flat, in which case (TM, \bar{g}) is flat.*

Proof. Clearly if the base manifold is flat, then so is (TM, \bar{g}) ; so we have only the necessity to prove. Using the above formulas for the connection

and curvature of \bar{g} , we have

$$\begin{aligned}
 ((\bar{\nabla}_{W^H} \bar{R})_{X^H Y^V} Z^V)_t &= \bar{\nabla}_{W^H} \left(-\frac{1}{2} \mathbf{R}_{Y Z} X - \frac{1}{4} \mathbf{R}_{t Y} \mathbf{R}_{t Z} X \right)_t^H \\
 &\quad - \bar{R}_{((D_W X)_t^H - \frac{1}{2} (\mathbf{R}_{W X} t)_t^V) Y^V} Z^V \\
 &\quad - \bar{R}_{X^H (\frac{1}{2} (\mathbf{R}_{t Y} W)_t^H + (D_W Y)_t^V)} Z^V \\
 &\quad - \bar{R}_{X^H Y^V} \left(\frac{1}{2} (\mathbf{R}_{t Z} W)_t^H + (D_W Z)_t^V \right).
 \end{aligned}$$

Using the hypothesis and taking the vertical part, we obtain

$$\begin{aligned}
 0 &= \frac{1}{2} \mathbf{R}_W (\frac{1}{2} \mathbf{R}_{Y Z} X + \frac{1}{4} \mathbf{R}_{t Y} \mathbf{R}_{t Z} X) t \\
 &\quad - \left(\mathbf{R}_X \frac{1}{2} \mathbf{R}_{t Y} W Z + \frac{1}{4} \mathbf{R}_{\mathbf{R}_{t Z} \frac{1}{2} \mathbf{R}_{t Y} W} X t - \frac{1}{4} \mathbf{R}_{\mathbf{R}_{t Z} X} \frac{1}{2} \mathbf{R}_{t Y} W t \right) \\
 &\quad - \left(\frac{1}{4} \mathbf{R}_{\mathbf{R}_{t Y} \frac{1}{2} \mathbf{R}_{t Z} W} X t + \frac{1}{2} \mathbf{R}_X \frac{1}{2} \mathbf{R}_{t Z} W Y \right).
 \end{aligned}$$

In this expression set $Y = t$ and $Z = t$ to get respectively

$$\begin{aligned}
 \mathbf{R}_W \mathbf{R}_{t Z} X t - \mathbf{R}_X \mathbf{R}_{t Z} W t &= 0, \\
 \mathbf{R}_W \mathbf{R}_{Y t} X t - 2 \mathbf{R}_X \mathbf{R}_{t Y} W t &= 0.
 \end{aligned}$$

Now replace Y by Z in the second of these equations and compare with the first to obtain

$$\mathbf{R}_X \mathbf{R}_{t Z} W t = 0.$$

Setting $W = X$ and taking the inner product with Z yields $|\mathbf{R}_{t Z} X|^2 = 0$ and hence that (M, G) is flat. \blacksquare

K. Bang [1994] obtained the corresponding result for (TM, \bar{g}) being conformally flat as a corollary to his result on conformally flat vector bundles, Theorem 9.13.

Theorem 9.2 *The tangent bundle TM with the Sasaki metric \bar{g} is conformally flat if and only if the base manifold (M, G) is flat, in which case (TM, \bar{g}) is flat.*

9.2 Tangent sphere bundles

We have seen that principal circle bundles over symplectic manifolds form a large class of examples of contact manifolds; they have K-contact structures which are Sasakian when the base manifolds are Kählerian. These examples together with the standard structure on \mathbb{R}^{2n+1} (Examples 3.2.1, 4.5.1, 6.7.1, 7.8.2) show that Sasakian manifolds form a large and important class of contact manifolds. However, despite the example of $T_1S^2 \cong \mathbb{R}P^3$ and more generally Theorem 9.3 below, the tangent and cotangent sphere bundles (Example 3.2.4) are not, in general, K-contact, even though they are classically an important class of contact manifolds. Boothby and Wang [1958] proved that a compact, simply connected, homogeneous contact manifold M of dimension $4r + 1$ with $r > 1$ is homeomorphic to a tangent sphere bundle only when M is the Stiefel manifold $V_{2r+2,2}$. In [1978] the author showed that the standard contact structure on the tangent sphere bundle of a compact Riemannian manifold of nonpositive constant curvature cannot be regular.

We will regard the tangent sphere bundle of a Riemannian manifold as the bundle of unit tangent vectors, even though, owing to the factor $\frac{1}{2}$ in the coboundary formula for $d\eta$, a homothetic change of metric will be made. (If one adopts the convention that the $\frac{1}{2}$ does not appear in the coboundary formula, this change is not necessary. However, to be consistent, the odd-dimensional sphere as a standard example of a Sasakian manifold should then be taken as a sphere of radius 2 (cf. Tashiro [1969] and Sasaki and Hatakeyama [1962]).)

The tangent sphere bundle, $\pi : T_1M \longrightarrow M$, is the hypersurface of TM defined by $\sum G_{ij}v^i v^j = 1$. The vector field $\nu = v^i \frac{\partial}{\partial v^i}$ is a unit normal as well as the position vector for a point $t \in T_1M$. We denote by g' the Riemannian metric induced on T_1M from the Sasaki metric \bar{g} on TM and by $\bar{\nabla}$ its Levi-Civita connection. We can easily find the Weingarten map of a hypersurface. For any vertical vector field U tangent to T_1M ,

$$\bar{\nabla}_U \nu = U v^i \frac{\partial}{\partial v^i} + v^i \bar{\nabla}_U \left(\frac{\partial}{\partial x^i} \right)^V = U.$$

For a horizontal tangent vector field X , we may suppose that X is the restriction of a horizontal lift. Then

$$\begin{aligned} (\bar{\nabla}_{(\partial_j)^H} \nu)_t &= ((\partial_j)^H v^i) \frac{\partial}{\partial v^i} + \frac{1}{2} v^i (\mathbf{R}_{t \partial_i} \partial_j)_t^H + v^i (D_{\partial_j} \partial_i)_t^V \\ &= \frac{1}{2} (\mathbf{R}_{t t} \partial_j)_t^H = 0, \end{aligned}$$

where we have again abbreviated $\frac{\partial}{\partial x^j}$ by ∂_j . Thus the Weingarten map A of T_1M with respect to the normal ν is given by $AU = -U$ for any vertical vector U and $AX = 0$ for any horizontal vector X .

With the simple form for the Weingarten map just obtained, many computations on T_1M can be done on TM . Yampol'skii [1985] (see also Borisenko and Yampol'skii [1987a], Tanno [1992]) computed the Levi-Civita connection and the curvature of g' .

While this is enough formalism to proceed, since vertical lifts need not be tangent to T_1M , Boeckx and Vanhecke [1997] introduced the notion of a tangential lift. For $X \in T_mM$, the *tangential lift* X^T of X to $(m, t) \in T_1M$ is given by

$$X^T = X^V - G(X, t)\nu.$$

The metric g' on T_1M at a point (m, t) is then given by

$$\begin{aligned} g'(X^H, Y^H) &= \bar{g}(X^H, Y^H) = G(X, Y), \quad g'(X^H, Y^T) = 0, \\ g'(X^T, Y^T) &= \bar{g}(X^T, Y^T) = G(X, Y) - G(X, t)G(Y, t). \end{aligned}$$

The Levi-Civita connection of g' is given by $\nabla_{X^T}Y^T = -G(Y, t)X^T$, while the formulas for $\nabla_{X^H}Y^H$, $\nabla_{X^H}Y^T$ and $\nabla_{X^T}Y^H$ are given by formulas for $\bar{\nabla}$ in the last section with vertical lifts replaced by tangential lifts, e.g.,

$$\nabla_{X^H}Y^H = (D_X Y)^H - \frac{1}{2}(\mathbf{R}_{X^H} t)^T,$$

since $G(\mathbf{R}_{\pi_* X} t, t) = 0$.

We know that as a hypersurface of the almost Kähler manifold TM , T_1M inherits an almost contact metric structure. Following the usual procedure (Example 4.5.2), we define ϕ' , ξ' , and η' by on T_1M by

$$\xi' = -J\nu = -v^i J \left(\frac{\partial}{\partial x^i} \right)^V = v^i \left(\frac{\partial}{\partial x^i} \right)^H, \quad JX = \phi'X + \eta'(X)\nu.$$

Then (ϕ', ξ', η', g') is an almost contact metric structure. Moreover, η' is the form on T_1M induced from the Liouville form β on TM , for

$$\begin{aligned} \eta'(X) &= \bar{g}(\nu, JX) = 2d\beta(\nu, X) = 2 \sum (d(G_{ij}v^j) \wedge dq^i) \left(v^k \frac{\partial}{\partial v^k}, X \right) \\ &= \sum G_{ij}v^j dq^i(X) = \beta(X). \end{aligned}$$

However, $g'(X, \phi'Y) = 2d\eta'(X, Y)$, so strictly speaking, (ϕ', ξ', η', g') is not a contact metric structure. Of course the difficulty is easily rectified and

$$\eta = \frac{1}{2}\eta', \quad \xi = 2\xi', \quad \phi = \phi', \quad g = \frac{1}{4}g'$$

is taken as the standard contact metric structure on T_1M . In local coordinates,

$$\xi = 2v^i \left(\frac{\partial}{\partial x^i} \right)^H;$$

the vector field $v^i \left(\frac{\partial}{\partial x^i} \right)^H$ is the well-known *geodesic flow* on T_1M .

Before proceeding to our theorems we obtain explicitly the covariant derivatives of ξ and ϕ . For a horizontal tangent vector field we may again use a horizontal lift. Then

$$\begin{aligned} (\nabla_{X^H} \xi)_t &= (\bar{\nabla}_{X^H} \xi)_t = (X^H 2v^i)(\partial_i)_t^H + 2v^i(D_X \partial_i)_t^H - (\mathbf{R}_{Xt})_t^V \\ &= -(\mathbf{R}_{Xt})_t^V, \end{aligned}$$

and hence for any horizontal vector X at $(m, t) \in T_1M$ we have

$$(\nabla_X \xi) = -(\mathbf{R}_{\pi_* X t})^V = -(\mathbf{R}_{\pi_* X t})^T.$$

For a vertical vector field U tangent to T_1M we have

$$(\nabla_U \xi)_t = (\bar{\nabla}_U \xi)_t = (U 2v^i)(\partial_i)_t^H - v^i(\mathbf{R}_{KU t}(\partial_i))_t^H = -2(\phi U)_t - (\mathbf{R}_{KU t})_t^H,$$

since $(\partial_i)^H = -J(\partial_i)^V$, or in terms of tangential lifts of a vector X on M ,

$$\nabla_{X^T} \xi = -2\phi X^T - (\mathbf{R}_{Xt})^H.$$

Comparing with $\nabla_X \xi = -\phi X - \phi hX$, we have for X horizontal and orthogonal to ξ and for U vertical,

$$hX = -X + (\mathbf{R}_{\pi_* X t})^H, \quad hU = U - (\mathbf{R}_{KU t})^T.$$

One can approach the differentiation of ϕ either in terms of lifts or by considering T_1M as a hypersurface in TM . First note that

$$\phi X^H = X^T, \quad \phi X^T = -X^H + \frac{1}{2}G(X, t)\xi$$

and that for any tangent vector fields X and Y ,

$$\begin{aligned} (\nabla_X \phi)Y &= \bar{\nabla}_X JY - (\nabla_X \eta')(Y)\nu + \eta'(Y)AX \\ &\quad - g'(X, A\phi Y)\nu - J\bar{\nabla}_X Y - g'(X, AY)\xi'. \end{aligned}$$

We present two computations, one done in each manner.

As before, for X, Y horizontal we suppose that they are horizontal lifts, and we have

$$\begin{aligned} (\nabla_{X^H} \phi)Y^H &= \nabla_{X^H} Y^T - \phi \nabla_{X^H} Y^H \\ &= \frac{1}{2}(\mathbf{R}_{t,Y} X)^H + (D_X Y)^T - \phi((D_X Y)^H - \frac{1}{2}(\mathbf{R}_{X,Y} t)^T) \\ &= \frac{1}{2}(\mathbf{R}_{t,X} Y)^H, \end{aligned}$$

where we have used the first Bianchi identity.

For $U = U^i \frac{\partial}{\partial v^i}$ a vertical vector field tangent to $T_1 M$ and X a horizontal tangent vector we have

$$\begin{aligned} ((\nabla_X \phi)U)_t &= -(XU^i)(\partial_i)_t^H - U^i(D_{\pi_* X} \partial_i)_t^H + \frac{1}{2}(\mathbf{R}_{\pi_* X} KU)_t^V \\ &\quad - (\nabla_X \eta')(U)\nu - J(XU^i)(\partial_i)_t^V - U^i J(D_{\pi_* X} \partial_i)_t^V \\ &\quad + \frac{1}{2}J(\mathbf{R}_{KU} \pi_* X)_t^H \\ &= \frac{1}{2}(\mathbf{R}_{\pi_* X} KU)_t^T, \end{aligned}$$

where we have used $(\nabla_X \eta')(U) = g'(\nabla_X \xi', U) = -\frac{1}{2}G((\mathbf{R}_{\pi_* X} t, KU) = \frac{1}{2}\bar{g}((\mathbf{R}_{\pi_* X} t, KU)_t^V, \nu)$.

Similarly one obtains

$$\begin{aligned} (\nabla_U \phi)X &= -2\eta(X)U + \frac{1}{2}(\mathbf{R}_{KU} \pi_* X)^T, \\ (\nabla_U \phi)W &= 2g(U, W)\xi + \frac{1}{2}(\mathbf{R}_{KU} KW)^H, \end{aligned}$$

where W is also a vertical vector field tangent to $T_1 M$.

We now prove a theorem of Tashiro [1969] which shows that the contact metric structure on the tangent sphere bundle is almost never Sasakian.

Theorem 9.3 *The contact metric structure (ϕ, ξ, η, g) on $T_1 M$ is K -contact if and only if the base manifold (M, G) has positive constant curvature $+1$, in which case $T_1 M$ is Sasakian.*

Proof. If (ϕ, ξ, η, g) is a K-contact structure, then $\nabla_X \xi = -\phi X$, as we have seen, but for a horizontal lift we have $(\nabla_{X^H} \xi)_t = -(\mathbf{R}_{X^H} t)^V$ and hence $(\phi X^H)_t = (\mathbf{R}_{X^H} t)^V$. Now for X orthogonal to t , $\phi X^H = X^V$ and therefore $\mathbf{R}_{X^H} t = X$ for all orthogonal pairs $\{X, t\}$ on (M, G) from which we have $\mathbf{R}_{X^H} Z = G(Y, Z)X - G(X, Z)Y$.

Conversely, by the formulas above for the covariant derivative of ϕ , the condition $\mathbf{R}_{X^H} Z = G(Y, Z)X - G(X, Z)Y$ on (M, G) gives us on $T_1 M$

$$(\nabla_X \phi)Y = \frac{1}{2}(G(\pi_* X, \pi_* Y)t - G(\pi_* Y, t)\pi_* X)^H = g(X, Y)\xi - \eta(Y)X,$$

$$(\nabla_X \phi)U = \frac{1}{2}(G(KU, t)\pi_* X - G(\pi_* X, KU)t)^T = 0,$$

$$(\nabla_U \phi)X = -2\eta(X)U + \frac{1}{2}(G(t, \pi_* X)KU - G(KU, \pi_* X)t)^T = -\eta(X)U,$$

and

$$(\nabla_U \phi)W = 2g(U, W)\xi + \frac{1}{2}(G(t, KW)KU - G(KU, KW)t)^H = g(U, W)\xi,$$

showing that $T_1 M$ is Sasakian. \blacksquare

In particular, the contact metric structure on the tangent sphere bundle of a unit sphere is η -Einstein, as shown by Tanno [1987b], and can be deformed by a \mathcal{D} -homothetic deformation to an Einstein metric. Thus $T_1 S^3(1) \sim S^3 \times S^2$ admits an Einstein metric other than a Riemannian product of constant curvature metrics. If g_0 denotes the standard metric on the sphere, then $(S^3, 2g_0) \times (S^2, g_0)$ is Einstein but clearly not an associated metric for any contact structure in view of Theorem 7.15 on locally symmetric contact metric manifolds. Recall also our discussion in Section 7.4 of the many Sasakian Einstein structures on $S^3 \times S^2$.

The symmetric operator l was defined by $lX = R_X \xi$ (Section 7.6). For a K-contact structure we have $l = I - \eta \otimes \xi$, and there exist many contact metric manifolds for which $l = 0$, as we will see in Section 9.4, Theorem 9.16. More strongly, we have seen that $R_{X^H} \xi = 0$ implies that the contact metric manifold M^{2n+1} is locally isometric to $E^{n+1} \times S^n(4)$ (Theorem 7.5), which is the tangent sphere bundle of Euclidean space. In the case of the contact metric structure on $T_1 M$ we have the following result (the author [1977]).

Theorem 9.4 *The standard contact metric structure (ϕ, ξ, η, g) on $T_1 M$ satisfies $lU = 0$ for vertical vector fields U if and only if the base manifold (M, G) is flat, in which case we have $R_{X^H} \xi = 0$ for all X and Y on $T_1 M$.*

Proof. First note that for a vertical vector field U tangent to T_1M , $R_U\xi\xi = 0$ implies $\bar{R}_U\xi\xi = 0$. Now recall that

$$K(\bar{R}_{X^H Y^V} Z^H)_t = \frac{1}{2}\mathbf{R}_{X Z} Y + \frac{1}{4}\mathbf{R}_{\mathbf{R}_{tY} Z X} t.$$

Since $\xi = 2v^i(\frac{\partial}{\partial x^i})^H$, combining these facts gives

$$\mathbf{R}_{\mathbf{R}_{KU} t t t} = 0,$$

i.e., $\mathbf{R}_{\mathbf{R}_{YX} X X} = 0$ for any orthonormal pair on (M, G) . Taking the inner product with Y gives $|\mathbf{R}_{YX} X|^2 = 0$ from which we see that (M, G) is flat.

Conversely, if (M, G) is flat, the Gauss equation for T_1M as a hypersurface of TM gives $g'(R'_{XY}\xi, Z) = 0$ and hence that $R_{XY}\xi = 0$. ■

We remark at this point that if the base manifold (M, G) has constant curvature -1 , then the sectional curvature $K(U, \xi) = 1$ for any vertical vector U tangent to T_1M and $K(X, \xi) = -7$ for any horizontal vector tangent to T_1M .

In Section 7.5 we studied the classification of all locally symmetric contact metric manifolds. Since these are locally isometric to $E^{n+1} \times S^n(4)$ or of constant curvature $+1$, the only possible cases for the standard contact metric structure on the tangent sphere bundle being locally symmetric are those in which the base manifold is flat or 2-dimensional and of constant curvature $+1$. The second case follows from Musso–Tricerri [1988], who prove that (T_1M, g') is Einstein only when $\dim M = 2$, or by Theorem 9.6 below on the conformally flat case. For a direct proof of this result on locally symmetric tangent sphere bundles see the author's paper [1989].

For the more general case of locally ϕ -symmetric contact metric manifolds we note the following theorem of Boeckx and Vanhecke [1997].

Theorem 9.5 *The standard contact metric structure (ϕ, ξ, η, g) on T_1M is strongly locally ϕ -symmetric if and only if the base manifold (M, G) is of constant curvature.*

When the base manifold is either odd-dimensional or of dimension 2, this is also equivalent to the contact metric structure on T_1M satisfying $\nabla_\xi h = ah\phi$ for some function a that is constant on the fibers (Boeckx, Perrone and Vanhecke [1998]).

The result for (T_1M, g) being conformally flat is much stronger than that of being locally symmetric (Koufogiorgos and the author [1994]).

Theorem 9.6 *The standard contact metric structure on T_1M is conformally flat if and only if the base manifold (M, G) is a surface of constant Gaussian curvature 0 or +1.*

We now study the condition $\nabla_\xi h = 0$ and prove a result of Perrone [1994]. For each unit tangent vector $t \in T_mM$, let $[t]^\perp$ be the subspace of T_mM orthogonal to t and define a symmetric linear transformation $L_t : [t]^\perp \longrightarrow [t]^\perp$ by $L_tX = \mathbf{R}_Xt$.

Lemma 9.1 *If the contact metric structure on T_1M satisfies $\nabla_\xi h = 0$, then for any orthonormal pair $\{X, t\}$ on (M, G) , $L_t^2X = L_tX$ and (M, G) is locally symmetric.*

Proof. Since $hU_t = U_t - (\mathbf{R}_{KU}t)^V$ for a vertical vector $U \in T_tT_1M$, we have $h^2U_t = U_t - 2(L_tKU)^V + (L_t^2KU)^V$. Proceeding as in the proof of Theorem 9.4, we also have $lU_t = (L_t^2KU)^V + 2((D_t\mathbf{R})_{KU}t)^H$. On the other hand, since $\nabla_\xi h = 0$, applying ϕ to the first equation of Proposition 7.1 gives $\phi^2 + h^2 + l = 0$. Applying this to U , the vertical part gives $L_t^2 = L_t$ on (M, G) . The horizontal part gives us that (M, G) is locally symmetric by a result of Cartan that a Riemannian manifold (M, G) is locally symmetric if and only if $G((D_X\mathbf{R})_YX, X) = 0$ for all orthonormal pairs $\{X, Y\}$ (English translation: Cartan [1983, pp. 257–258]). ■

Theorem 9.7 *The standard contact metric structure (ϕ, ξ, η, g) on T_1M satisfies $\nabla_\xi h = 0$ if and only if the base manifold (M, G) is of constant curvature 0 or +1.*

Proof. This theorem follows from Lemma 9.1 and the purely Riemannian result (Perrone [1994]) that a Riemannian manifold (M, G) is locally symmetric and satisfies $L_t^2 = L_t$ if and only if (M, G) is of constant curvature 0 or +1. ■

As we have been seeing from time to time, conditions on $\nabla_\xi h$ on a contact metric manifold are of significant geometric interest. Here we mention a result of Boeckx, Perrone and Vanhecke [1998]. Define a tensor field S of type $(1, 1)$ by $SX^H = X^H$ and $SX^T = -X^T$.

Theorem 9.8 *The base manifold (M, G) is locally isometric to a two-point homogeneous space if and only if the contact metric structure*

(ϕ, ξ, η, g) on T_1M satisfies

$$\nabla_\xi h = ah\phi + b\phi S,$$

where a and b are functions that are constant along the fibers of T_1M . Moreover, this is true if and only if the eigenvalues of h are constant along the fibers and l maps vertical vectors to vertical vectors.

In Section 6.4 we saw that a contact metric structure gives rise to a strongly pseudoconvex CR-structure if and only if

$$(\nabla_X \phi)Y = g(X + hX, Y)\xi - \eta(Y)(X + hX).$$

For the tangent sphere bundle we have the following result of Mitric [1991], Tanno [1992].

Theorem 9.9 *For $\dim M \geq 3$ the standard contact metric structure on T_1M gives rise to a strongly pseudoconvex CR-structure if and only if the base manifold (M, G) is of constant curvature.*

Proof. Using $hU_t = U_t - (\mathbf{R}_{KU}t)^V$, evaluate the right-hand side of

$$(\nabla_X \phi)Y = g(X + hX, Y)\xi - \eta(Y)(X + hX)$$

on two vertical vectors U and W and compare with our earlier expression for the left-hand side. The right-hand side becomes $2g(U, W)\xi_t - g((\mathbf{R}_{KU}t)_t^V, W)\xi_t$, while the left-hand side is $2g(U, W)\xi_t + \frac{1}{2}(\mathbf{R}_{KU}tKW)_t^H$. Equality implies that on the base manifold, $\mathbf{R}_{Xt}Y$ is collinear with t for all X, Y orthogonal to t . In particular, $G(\mathbf{R}_{Xt}Y, X) = 0$ for every orthonormal triple $\{t, X, Y\}$, and hence (M, G) is of constant curvature. Conversely, if $\mathbf{R}_{XY}Z = c(G(Y, Z)X - G(X, Z)Y)$, evaluate both sides of $(\nabla_X \phi)Y = g(X + hX, Y)\xi - \eta(Y)(X + hX)$ for the four cases of X, Y being both horizontal, both vertical and each one horizontal with the other vertical, and compare. ■

Tanno actually proves more: In [1992] (and [1991]) he introduces a gauge invariant B of type $(1, 3)$ (on the contact subbundle) whose vanishing implies the CR-condition of Theorem 6.7 and conversely under the CR-condition, B reduces to the Chern–Moser–Tanaka invariant. See Section 10.5 for the notion of a gauge transformation. Tanno proves that on T_1M , $\dim M \geq 3$, B vanishes if and only if (M, G) is of constant curvature -1 .

Th. Koufogiorgos studied in [1997a] the idea of constant ϕ -sectional curvature for non-Sasakian contact metric manifolds, especially for (κ, μ) -manifolds. For the tangent sphere bundle he proved the following theorem.

Theorem 9.10 *If (M, G) is of constant curvature c and dimension ≥ 3 , the standard contact metric structure on T_1M has constant ϕ -sectional curvature (equal to c^2) if and only if $c = 2 \pm \sqrt{5}$.*

For (M, G) a surface of constant curvature $c \neq 1$, (T_1M, g) has constant ϕ -sectional curvature c^2 , as follows readily from Theorem 7.9 and the formula $K(X, \phi X) = -(\kappa + \mu)$ from the same section.

Recently E. Boeckx [2003] [2005] studied the question of the reducibility of the tangent sphere bundle.

Theorem 9.11 *The standard contact metric structure on the tangent sphere bundle of a Riemannian manifold of dimension greater than 2 is locally reducible if and only if the base manifold has a flat factor. If the tangent sphere bundle is a global Riemannian product, then the base manifold is either flat or a global Riemannian product.*

Additional results on the geometry of the tangent sphere bundle can be found in the survey of G. Calvaruso [2005].

9.3 Geometry of vector bundles

The geometric constructions on the tangent bundle described in Section 9.1 can be carried out on a general vector bundle. In this section we describe this construction and the corresponding results without proofs. We follow the treatment given by K. Bang in [1994].

Let (M^n, G) be a Riemannian manifold with Levi-Civita connection D and curvature tensor \mathbf{R} as before. Consider a vector bundle $\pi : \mathbf{E}^{n+k} \longrightarrow M^n$ with fiber metric g^\perp and a metric connection ∇ . If (x^1, \dots, x^n) are local coordinates on M , set $q^i = x^i \circ \pi$; if $\{e_\alpha\}$ is a local orthonormal basis of sections of \mathbf{E} , writing a point $(m, U) \in \mathbf{E}$ as $U = \sum u^\alpha e_\alpha$, (q^1, \dots, q^n) together with the fiber coordinates (u^1, \dots, u^k) form local coordinates on \mathbf{E} .

For a section $\zeta = \sum \zeta^\alpha e_\alpha$ of \mathbf{E} , the connection ∇ is given by

$$\nabla_X \zeta = X^i \left(\frac{\partial \zeta^\alpha}{\partial x^i} + \zeta^\beta \mu_{\beta i}^\alpha \right) e_\alpha,$$

where $\nabla_{\partial_i} e_\beta = \mu_{\beta i}^\alpha e_\alpha$.

If X is a tangent vector field on M and if ζ is a section of \mathbf{E} , the *horizontal lift* X^H of X and the *vertical lift* ζ^V of ζ are defined by

$$X^H = X^i \frac{\partial}{\partial q^i} - X^i u^\beta \mu_{\beta i}^\alpha \frac{\partial}{\partial u^\alpha}$$

and

$$\zeta^V = \zeta^\alpha \frac{\partial}{\partial u^\alpha}.$$

Then $\pi_* X^H = X$ and $\pi_* \zeta^V = 0$. Define a linear map $K : T\mathbf{E} \longrightarrow \mathbf{E}$ by

$$KX^H = 0, \quad K\zeta_{(m,U)}^V = \zeta_m, \quad (m, U) \in \mathbf{E},$$

or by its local expression: If $(\tilde{X}^i, \tilde{X}^{n+\alpha})$ are the components of a vector \tilde{X} tangent to \mathbf{E} at (m, U) with respect to the coordinate basis, then

$$K\tilde{X} = (\tilde{X}^{n+\alpha} + \tilde{X}^i u^\beta \mu_{\beta i}^\alpha) e_\alpha.$$

We define a Riemannian metric \bar{g} on \mathbf{E} , called the *Sasaki metric*, by

$$\bar{g}(\tilde{X}, \tilde{Y}) = G(\pi_* \tilde{X}, \pi_* \tilde{Y}) + g^\perp(K\tilde{X}, K\tilde{Y}),$$

where \tilde{X} and \tilde{Y} are vector fields on \mathbf{E} . When $\mathbf{E} = TM$ and $\nabla = D$, \bar{g} is the Sasaki metric on TM .

The curvature tensor R of ∇ is given by $R_{XY}\zeta = \nabla_X \nabla_Y \zeta - \nabla_Y \nabla_X \zeta - \nabla_{[X,Y]}\zeta$ and ∇ is said to be *flat* if R vanishes for all vector fields X, Y on M and all sections ζ of \mathbf{E} . Since $R_{XY}\zeta$ is also a section of \mathbf{E} , we can compute its inner product $g^\perp(R_{XY}\zeta, \psi)$ with another section ψ . We then define the adjoint $\hat{R}_{\zeta\psi}X$ by

$$G(\hat{R}_{\zeta\psi}X, Y) = g^\perp(R_{XY}\zeta, \psi).$$

With this notation in mind we give the the covariant derivatives of the Levi-Civita connection $\bar{\nabla}$ of the Sasaki metric \bar{g} at a point $(m, U) \in \mathbf{E}$:

$$\begin{aligned} \bar{\nabla}_{X^H} Y^H &= (D_X Y)^H - \frac{1}{2}(R_{XY}U)^V, & \bar{\nabla}_{X^H} \zeta^V &= \frac{1}{2}(\hat{R}_U \zeta X)^H + (\nabla_X \zeta)^V, \\ \bar{\nabla}_{\zeta^V} Y^H &= \frac{1}{2}(\hat{R}_U \zeta Y)^H, & \bar{\nabla}_{\zeta^V} \psi^V &= 0. \end{aligned}$$

The curvature \bar{R} of the Sasaki metric at (m, U) is given as follows: \bar{R} vanishes on three vertical lifts and

$$\begin{aligned}
 \bar{R}_{\zeta^V \psi^V} Z^H &= \left(\hat{R}_{\zeta \psi} Z + \frac{1}{4} \hat{R}_{U \zeta} \hat{R}_{U \psi} Z - \frac{1}{4} \hat{R}_{U \psi} \hat{R}_{U \zeta} Z \right)^H, \\
 \bar{R}_{X^H \zeta^V} \psi^V &= - \left(\frac{1}{2} \hat{R}_{\zeta \psi} X + \frac{1}{4} \hat{R}_{U \zeta} \hat{R}_{U \psi} X \right)^H, \\
 \bar{R}_{X^H \zeta^V} Z^H &= \frac{1}{2} \left((D_X \hat{R})_{U \zeta} Z \right)^H + \left(\frac{1}{2} R_{X Z \zeta} + \frac{1}{4} R_{\hat{R}_{U \zeta} X Z} U \right)^V, \\
 \bar{R}_{X^H Y^H} \zeta^V &= \frac{1}{2} \left((D_X \hat{R})_{U \zeta} Y - (D_Y \hat{R})_{U \zeta} X \right)^H \\
 &\quad + \left(R_{X Y \zeta} + \frac{1}{4} R_{\hat{R}_{U \zeta} Y X} U - \frac{1}{4} R_{\hat{R}_{U \zeta} X Y} U \right)^V, \\
 \bar{R}_{X^H Y^H} Z^H &= \left(R_{X Y Z} + \frac{1}{4} \hat{R}_{U R_{Z Y} U} X + \frac{1}{4} \hat{R}_{U R_{X Z} U} Y + \frac{1}{2} \hat{R}_{U R_{X Y} U} Z \right)^H \\
 &\quad + \frac{1}{2} \left((\nabla_Z R)_{X Y} U \right)^V.
 \end{aligned}$$

Concerning the questions of the vector bundle \mathbf{E} with the Sasaki metric being locally symmetric or conformally flat, K. Bang proved the following theorems.

Theorem 9.12 *Let $\pi : \mathbf{E}^{n+k} \longrightarrow M^n$ be a vector bundle over a Riemannian manifold (M, G) with fiber metric g^\perp and a metric connection ∇ . Then the Sasaki metric on \mathbf{E} is locally symmetric if and only if the connection ∇ is flat and (M, G) is locally symmetric.*

Theorem 9.13 *Let $\pi : \mathbf{E}^{n+k} \longrightarrow M^n$ be a vector bundle over a Riemannian manifold (M, G) with fiber metric g^\perp and a metric connection ∇ . Then the Sasaki metric on \mathbf{E} is conformally flat if and only if either (M, G) is flat with flat connection ∇ or (M, G) has constant curvature with flat connection ∇ and $k = 1$.*

As a corollary K. Bang pointed out that the tangent bundle TM with the Sasaki metric \bar{g} is conformally flat if and only if the base manifold is flat (Theorem 9.2 above).

9.4 Normal bundles

For the case of the normal bundle of a submanifold of a Riemannian manifold, the above construction of the Sasaki metric was developed by Reckziegel [1979] and Borisenko and Yampol'skii [1987b] using the normal connection ∇^\perp . In this section we consider the special cases in which the submanifold is a Lagrangian submanifold of a Kähler manifold or an integral submanifold of a Sasakian manifold. Again the results were obtained by K. Bang in his thesis [1994].

First let L be a Lagrangian submanifold of a symplectic manifold M^{2n} with associated metric \tilde{g} and corresponding almost complex structure \tilde{J} . Recall that if X is tangent to L , then $\tilde{J}X$ is normal. Typically, normal vectors will be denoted by ζ , ν , ψ . Using the ideas of horizontal and vertical lifts in the case of vector bundles from the last section, we define an almost complex structure \bar{J} on the normal bundle $T^\perp L$ of L by

$$\bar{J}X^H = (\tilde{J}X)^V, \quad \bar{J}\zeta^V = (\tilde{J}\zeta)^H.$$

That $\bar{J}^2 = -I$ and that the Sasaki metric \bar{g} is Hermitian with respect to \bar{J} are easily verified. However, the symplectic nature of $(T^\perp L, \bar{J}, \bar{g})$ depends on the ambient manifold $(M^{2n}, \tilde{J}, \tilde{g})$ being Kähler. Let $\bar{\Omega}$ be the fundamental 2-form of the almost Hermitian structure just defined on $T^\perp L$.

Theorem 9.14 *Let L be a Lagrangian submanifold of a Kähler manifold $(M^{2n}, \tilde{J}, \tilde{g})$. Then the normal bundle $(T^\perp L, \bar{\Omega})$ is a symplectic manifold.*

Proof. Since (\bar{J}, \bar{g}) is an almost Hermitian structure, it is immediate that $\bar{\Omega}^n \neq 0$, so we have only to show that $\bar{\Omega}$ is a closed 2-form. First, it is immediate that $\bar{\Omega}(X^H, Y^H) = 0$, $\bar{\Omega}(X^H, \zeta^V) = \tilde{g}(X, J\zeta)$, and $\bar{\Omega}(\zeta^V, \psi^V) = 0$. Now for horizontal lifts of vector fields tangent to L , we have

$$[X^H, Y^H] = [X, Y]^H - (R_X^\perp Y \nu)^V$$

at the point $\nu \in T^\perp L$. Thus using the coboundary formula for $d\bar{\Omega}$, we have

$$\begin{aligned} 3d\bar{\Omega}(X^H, Y^H, Z^H) &= X^H \bar{\Omega}(Y^H, Z^H) + Y^H \bar{\Omega}(Z^H, X^H) + Z^H \bar{\Omega}(X^H, Y^H) \\ &\quad - \bar{\Omega}([X^H, Y^H], Z^H) - \bar{\Omega}([Y^H, Z^H], X^H) - \bar{\Omega}([Z^H, X^H], Y^H) \\ &= \bar{\Omega}((R_X^\perp Y \nu)^V, Z^H) + \bar{\Omega}((R_Y^\perp Z \nu)^V, X^H) + \bar{\Omega}((R_Z^\perp X \nu)^V, Y^H). \end{aligned}$$

In keeping with the notation of this chapter, let G be the induced metric on L , D its Levi-Civita connection and \mathbf{R} is curvature tensor. Using the Gauss–Weingarten equations and the Kähler condition, one readily obtains

$$\tilde{J}\nabla_X^\perp\nu = D_X\tilde{J}\nu.$$

Thus we have

$$\begin{aligned}\bar{\Omega}((R_{X\,Y}^\perp\nu)^V, Z^H) &= \bar{g}((R_{X\,Y}^\perp\nu)^V, \tilde{J}Z^H) = -\bar{g}((\tilde{J}R_{X\,Y}^\perp\nu)^H, Z^H) \\ &= -G(\mathbf{R}_{X\,Y}\tilde{J}\nu, Z) = G(\mathbf{R}_{X\,Y}Z, \tilde{J}\nu).\end{aligned}$$

Substituting this and like expressions in the coboundary formula and using the Bianchi identity, we have $3d\bar{\Omega}(X^H, Y^H, Z^H) = 0$. In much the same way one shows that $d\bar{\Omega}(X^H, Y^H, \zeta^V) = 0$ and $d\bar{\Omega}(X^H, \zeta^V, \psi^V) = 0$, and of course $d\bar{\Omega}$ vanishes on three vertical vectors. ■

We now turn to the question of when the above almost Kähler structure on $T^\perp L$ is itself Kähler.

Theorem 9.15 *Let L be a Lagrangian submanifold of a Kähler manifold $(M^{2n}, \tilde{J}, \tilde{g})$. Then the following are equivalent: (i) $(T^\perp L, \bar{J}, \bar{g})$ is Kähler. (ii) L has flat normal connection. (iii) L is flat.*

Proof. In the previous proof we noted that $\tilde{J}\nabla_X^\perp\nu = D_X\tilde{J}\nu$ and therefore $\mathbf{R}_{X\,Y}\tilde{J}\nu = \tilde{J}R_{X\,Y}^\perp\nu$. Thus L is flat if and only if L has flat normal connection.

To complete the proof we show that the almost complex structure \bar{J} on $T^\perp L$ is integrable if and only if L has flat normal connection. This will be done by computing the Nijenhuis torsion of \bar{J} :

$$\begin{aligned}[\bar{J}, \bar{J}](X^H, \zeta^V) &= -[X^H, \zeta^V] + [(\tilde{J}X)^V, (\tilde{J}\zeta)^H] - \bar{J}[(\tilde{J}X)^V, \zeta^V] \\ &\quad - \bar{J}[X^H, (\tilde{J}\zeta)^H] \\ &= -(\nabla_X^\perp\zeta)^V - (\nabla_{\tilde{J}\zeta}^\perp\tilde{J}X)^V - \bar{J}([X, \tilde{J}\zeta]^H - (R_{X\,\tilde{J}\zeta}^\perp\nu)^V) \\ &= -(\nabla_X^\perp\zeta + \nabla_{\tilde{J}\zeta}^\perp\tilde{J}X + \tilde{J}[X, \tilde{J}\zeta])^V + (\tilde{J}R_{X\,\tilde{J}\zeta}^\perp\nu)^H.\end{aligned}$$

Now using the Kähler condition and the fact that $\tilde{J}[X, \tilde{J}\zeta]$ is normal to L one can show that $\nabla_X^\perp\zeta + \nabla_{\tilde{J}\zeta}^\perp\tilde{J}X + \tilde{J}[X, \tilde{J}\zeta] = 0$. Therefore

$$[\bar{J}, \bar{J}](X^H, \zeta^V) = (\tilde{J}R_{X\,\tilde{J}\zeta}^\perp\nu)^H.$$

Similarly

$$[\bar{J}, \bar{J}](X^H, Y^H) = (R_{X Y}^\perp \nu)^V, \quad [\bar{J}, \bar{J}](\zeta^V, \psi^V) = -(R_{\bar{J}\zeta}^\perp \bar{J}\psi^V)^V,$$

and the result follows. \blacksquare

We remark that in Chapter 1 we proved a theorem of Weinstein that a symplectic manifold is locally the cotangent bundle of any Lagrangian submanifold. Here one may note that since $\tilde{J}\nabla_X^\perp \nu = D_X \tilde{J}\nu$, \tilde{J} provides a connection preserving isomorphism between the tangent bundle and the normal bundle of L .

Turning to the contact case, let M^n be an integral submanifold of a contact metric manifold M^{2n+1} with structure tensors $(\tilde{\phi}, \tilde{\xi}, \tilde{\eta}, \tilde{g})$. On the normal bundle $T^\perp M^n$, we define an almost contact structure $(\bar{\phi}, \bar{\xi}, \bar{\eta})$ by

$$\bar{\phi}X^H = (\tilde{\phi}X)^V, \quad \bar{\phi}\tilde{\xi}^V = 0, \quad \bar{\phi}\zeta^V = (\tilde{\phi}\zeta)^H$$

for all tangent vectors X and normal vectors ζ orthogonal to $\tilde{\xi}$. Also let

$$\bar{\xi} = \tilde{\xi}^V, \quad \bar{\eta}(X) = \bar{g}(X, \bar{\xi})$$

for any vector X . Then $\bar{\eta}(\bar{\xi}) = 1$ and $\bar{\phi}^2 = -I + \bar{\eta} \otimes \bar{\xi}$ follow easily.

Using the coboundary formula for $d\bar{\eta}$ we have at a point $\nu \in T^\perp M^n$,

$$\begin{aligned} 2d\bar{\eta}(X^H, Y^H) &= X^H \bar{\eta}(Y^H) - Y^H \bar{\eta}(X^H) - \bar{\eta}([X^H, Y^H]) \\ &= -\bar{g}([X^H, Y^H], \bar{\xi}) = \tilde{g}(R_{X Y}^\perp \nu, \tilde{\xi}) \\ &= \tilde{g}(\tilde{R}_{X Y} \nu, \tilde{\xi}) + \tilde{g}([A_\nu, A_{\tilde{\xi}}]X, Y) \end{aligned}$$

by the equation of Ricci–Kühne. For a normal vector ζ orthogonal to $\tilde{\xi}$ we have

$$2d\bar{\eta}(X^H, \zeta^V) = -\bar{g}([X^H, \zeta^V], \bar{\xi}) = -\tilde{g}(\nabla_X^\perp \zeta, \tilde{\xi}) = -\tilde{g}(\tilde{\nabla}_X \zeta, \tilde{\xi}) = \tilde{g}(\zeta, \tilde{\nabla}_X \tilde{\xi}).$$

Similarly $d\bar{\eta}(X^H, \bar{\xi}) = 0$, $d\bar{\eta}(\zeta^V, \bar{\xi}) = 0$ and $d\bar{\eta}(\zeta^V, \psi^V) = 0$.

If now M^{2n+1} is Sasakian, by Lemma 8.1, $A_{\tilde{\xi}} = 0$, and by Proposition 7.3, $\tilde{R}_{X Y} \tilde{\xi} = \tilde{\eta}(Y)X - \tilde{\eta}(X)Y$, giving $\tilde{g}(R_{X Y}^\perp \nu, \tilde{\xi}) = \tilde{g}(\tilde{R}_{X Y} \nu, \tilde{\xi}) = 0$. Thus $R_{X Y}^\perp \tilde{\xi} = 0$ for any tangent vectors X and Y , and in turn, $\hat{R}_{\tilde{\xi}\nu} X = 0$ for all $\nu \in T^\perp M^n$. Now setting

$$\phi = \bar{\phi}, \quad \xi = 2\bar{\xi}, \quad \eta = \frac{1}{2}\bar{\eta}, \quad g = \frac{1}{4}\bar{g},$$

we see that $d\eta(X, Y) = g(X, \phi Y)$ for all vector fields on $T^\perp M$, giving $T^\perp M^n$ a contact metric structure (ϕ, ξ, η, g) .

In Section 7.6 we mentioned in passing that there exist many contact metric manifolds satisfying $l = 0$ ($lX = R_X \xi$). This is seen from the following theorem of Bang [1994].

Theorem 9.16 *Let M^n be an integral submanifold of a Sasakian manifold M^{2n+1} with structure tensors $(\tilde{\phi}, \tilde{\xi}, \tilde{\eta}, \tilde{g})$. Then the normal bundle, $T^\perp M^n$, has a contact metric structure (ϕ, ξ, η, g) satisfying $l = 0$.*

Proof. We have just seen that when M^n is an integral submanifold of a Sasakian manifold M^{2n+1} , $T^\perp M^n$ has a contact metric structure (ϕ, ξ, η, g) . So it remains only to show that $l = 0$. From the curvature expressions of Section 9.3 and $\hat{R}_{\tilde{\xi}\nu} X = 0$ we have

$$R_{X^H} \xi = 4R_{X^H \tilde{\xi}^V} \tilde{\xi}^V = -(\hat{R}_\nu \tilde{\xi} \hat{R}_\nu X)^H = 0$$

and $R_{\zeta^V} \xi = 0$. ■

In particular, we see that the contact metric structure (ϕ, ξ, η, g) on $T^\perp M^n$ is never Sasakian. Concerning the question of flat normal connection we have the following result of Yano and Kon [1983, p. 50].

Theorem 9.17 *Let M^n be an integral submanifold of a Sasakian manifold M^{2n+1} . Then M^n has flat normal connection if and only if (M^n, G) has constant curvature 1.*

As an example of Theorem 9.17 recall Example 5.3.1 of S^n as a totally geodesic integral submanifold of the Sasakian manifold S^{2n+1} . In this case the normal bundle $T^\perp S^n$ has flat normal connection, and $T^\perp S^n$ as a contact metric manifold with the structure (ϕ, ξ, η, g) is again the common example $E^{n+1} \times S^n(4)$ (cf. Theorem 7.5).

We also point out that for a normal vector ζ orthogonal to $\tilde{\xi}$, we have on $T^\perp M^n$ that $\nabla_{\zeta^V} \xi = 2\nabla_{\zeta^V} \tilde{\xi}^V = 0$ where ∇ is the Levi-Civita connection of g . Thus from Lemma 6.2, $h\zeta^V = -\zeta^V$, and hence -1 is an eigenvalue of h with multiplicity n . Since $h\phi + \phi h = 0$, $+1$ is also an eigenvalue with multiplicity n and $hX^H = X^H$. From $[X^H, Y^H] = [X, Y]^H - (R_{X^H}^\perp Y^H)^V$ we see that the subbundle $[+1]$ is integrable if and only if M^n has flat normal connection.

We close this section with a continuation of Example 5.3.2. There we gave an embedding of the 2-torus as a flat minimal integral submanifold of S^5 . In keeping with the notation of Section 9.3 we let $\{x^1, x^2\}$ be

the local coordinates on T^2 instead of $\{u, v\}$; $X_1 = \frac{\partial}{\partial x^1}, X_2 = \frac{\partial}{\partial x^2}$ are orthonormal in the metric G , the restriction of \tilde{g} to T^2 . Let $e_1 = \tilde{\phi}X_1$, $e_2 = \tilde{\phi}X_2$ and $e_3 = \tilde{\xi}$. To find $\nabla_{\partial_i}^\perp e_\beta = \mu_{\beta i}^\alpha e_\alpha$ explicitly, compute $\tilde{\nabla}e_\alpha$ using the Sasakian condition. Then one obtains

$$\mu_{31}^1 = -1, \quad \mu_{32}^2 = -1, \quad \mu_{11}^3 = 1, \quad \mu_{22}^3 = 1$$

as the nonzero $\mu_{\beta i}^\alpha$'s. Also $\{u^1, u^2, u^3\}$ denote the fiber coordinates, so that $\{q^i = x^i \circ \pi, u^\alpha\}$ are local coordinates on $T^\perp T^2$. Now computing the Sasaki metric \bar{g} , we have

$$\begin{aligned} \bar{g}\left(\frac{\partial}{\partial q^i}, \frac{\partial}{\partial q^j}\right) &= \delta_{ij} + \sum \mu_{\beta i}^\alpha \mu_{\delta j}^\alpha u^\beta u^\delta, \\ \bar{g}\left(\frac{\partial}{\partial q^i}, \frac{\partial}{\partial u^\alpha}\right) &= \mu_{\beta i}^\alpha u^\beta, \quad \bar{g}\left(\frac{\partial}{\partial u^\alpha}, \frac{\partial}{\partial u^\beta}\right) = \delta_{\alpha\beta}. \end{aligned}$$

Thus for the contact metric metric structure (η, g) on $T^\perp T^2$ we have

$$\eta = \frac{1}{2}(du^3 + u^1 dq^1 + u^2 dq^2)$$

and

$$g = \frac{1}{4} \begin{pmatrix} 1 + (u^1)^2 + (u^3)^2 & u^1 u^2 & -u^3 & 0 & u^1 \\ u^1 u^2 & 1 + (u^2)^2 + (u^3)^2 & 0 & -u^3 & u^2 \\ -u^3 & 0 & 1 & 0 & 0 \\ 0 & -u^3 & 0 & 1 & 0 \\ u^1 & u^2 & 0 & 0 & 1 \end{pmatrix}.$$

Compare this metric with the metric

$$g = \frac{1}{4} \begin{pmatrix} \delta_{ij} + y^i y^j + \delta_{ij} z^2 & \delta_{ij} z & -y^i \\ \delta_{ij} z & \delta_{ij} & 0 \\ -y^j & 0 & 1 \end{pmatrix}$$

associated to the Darboux form $\eta = \frac{1}{2}(dz - \sum y^i dx^i)$ on \mathbb{R}^{2n+1} introduced in Section 7.2 as a formal generalization of the flat associated metric of the Darboux form on \mathbb{R}^3 .

9.5 The geodesic flow on the projectivized tangent bundle

In this section we consider the geodesic flow on the projectivized tangent bundle rather than on its usual home, the unit tangent bundle. We do this in anticipation of our discussion of the complex case, Sections 13.2–13.4, and for illustrative purposes we restrict ourselves to case where the base manifold is the Beltrami model of the hyperbolic plane. This model is the open unit disk in the xy -plane with the Beltrami metric G given by

$$ds^2 = \frac{(1 - y^2)dx^2 + 2xy \, dx \, dy + (1 - x^2)dy^2}{(1 - x^2 - y^2)^2}.$$

Geodesics in the model are chords of the disk.

The projectivized tangent bundle, PTM , is the product of the disk with a real projective line, which one can imagine as vertical line fibers over a horizontal disk. Let $m = \frac{v^2}{v^1}$ be a nonhomogenous coordinate on the fibers, v^1 and v^2 being homogeneous coordinates. We will use (x, y, m) , or (x^1, x^2, m) to accommodate indexing $(1, 2, 3)$, to denote the local coordinates in the projectivized tangent bundle. The metric g on PTM will be the projection of the Sasaki metric on the tangent bundle, and the formalism is the following. Set $P_i = G_{i1}v^1 + G_{i2}v^2$. Instead of tangential lifts we introduce *projective lifts* by

$$\left(\frac{\partial}{\partial x^i}\right)^P = v^1 \left(\frac{\partial}{\partial v^i} - P_i \sum v^j \frac{\partial}{\partial v^j}\right).$$

The projective lifts of the coordinate fields project to PTM as

$$\left(\frac{\partial}{\partial x}\right)^P \longrightarrow -m \frac{\partial}{\partial m}, \quad \left(\frac{\partial}{\partial y}\right)^P \longrightarrow \frac{\partial}{\partial m}.$$

To obtain the metric g , first recall that the unit tangent bundle is defined by $G_{11}(v^1)^2 + 2G_{12}v^1v^2 + G_{22}(v^2)^2 = 1$, and we construct the fiber as a projective line ($m = \frac{v^2}{v^1}$) from this restriction. Note that $v^1 = \frac{1-x^2-y^2}{\sqrt{1+m^2-(mx-y)^2}}$. First,

$$g_{33} = g\left(\frac{\partial}{\partial m}, \frac{\partial}{\partial m}\right) = \bar{g}\left(\left(\frac{\partial}{\partial y}\right)^P, \left(\frac{\partial}{\partial y}\right)^P\right) = \frac{1 - x^2 - y^2}{(1 + m^2 - (mx - y)^2)^2}.$$

Using this, the component g_{13} is readily computed from

$$\begin{aligned} 0 &= \bar{g}\left(\left(\frac{\partial}{\partial x}\right)^H, \left(\frac{\partial}{\partial y}\right)^P\right) = g\left(\frac{\partial}{\partial x} + m\frac{x+my}{1-x^2-y^2}\frac{\partial}{\partial m}, \frac{\partial}{\partial m}\right) \\ &= g_{13} + \frac{m(x+my)}{(1+m^2-(mx-y)^2)^2}. \end{aligned}$$


Similarly, one finds g_{23} , and then using horizontal lifts one finds g_{11} , g_{12} and g_{22} . In summary,

$$\begin{aligned} g_{11} &= \frac{1-y^2}{(1-x^2-y^2)^2} + \frac{m^2(x+my)^2}{(1-x^2-y^2)(1+m^2-(mx-y)^2)^2}, \\ g_{12} &= \frac{xy}{(1-x^2-y^2)^2} - \frac{m(x+my)^2}{(1-x^2-y^2)(1+m^2-(mx-y)^2)^2}, \\ g_{22} &= \frac{1-x^2}{(1-x^2-y^2)^2} + \frac{(x+my)^2}{(1-x^2-y^2)(1+m^2-(mx-y)^2)^2}, \\ g_{13} &= -\frac{m(x+my)}{(1+m^2-(mx-y)^2)^2}, \quad g_{23} = \frac{(x+my)}{(1+m^2-(mx-y)^2)^2}, \\ g_{33} &= \frac{1-x^2-y^2}{(1+m^2-(mx-y)^2)^2}. \end{aligned}$$

The “geodesic flow” is given by the vector field $\frac{\partial}{\partial x} + m\frac{\partial}{\partial y}$, and an integral curve through a point (x, y, m) is a line of slope m lying in a copy of the disk at height m and passing through (x, y) . Note that the point at infinity on a fiber corresponds to a vertical line in the (x, y) -plane. The geodesic flow is hyperbolic (an Anosov flow without the compactness requirement), as can be visualized in the following discussion.

Consider a directed line (chord) in the disk and a point on it. Through the point there are two horocycles; these are ellipses tangent to each other at the point and tangent to the boundary of the disk at the ends of the chord. Asymptotic parallels to the line in the positive direction have greater slope on one side of the line and lesser slope on the other and are orthogonal to the horocycle in the positive direction from the point. The curve in the projectivized tangent bundle lying over (and under) the horocycle whose m -value at each point is the slope of the corresponding asymptotic parallel is the stable submanifold of the geodesic flow through each of its points. Similarly, the asymptotic parallels to the line in the negative direction have lesser slope on the first side and greater on the other and are orthogonal to the horocycle in the negative direction.

The curve in the projectivized tangent bundle lying under (and over) the horocycle whose m -value at each point is the slope of the corresponding asymptotic parallel is the unstable submanifold of the geodesic flow through each of its points.



The vector field $2 \sum v^j (\frac{\partial}{\partial x^j})^H$ projects to

$$\xi = 2 \frac{1 - x^2 - y^2}{\sqrt{1 + m^2 - (mx - y)^2}} \left(\frac{\partial}{\partial x} + m \frac{\partial}{\partial y} \right)$$

and its covariant form is

$$\eta = \frac{(1 + y(mx - y))dx + (m - x(mx - y))dy}{2(1 - x^2 - y^2)\sqrt{1 + m^2 - (mx - y)^2}}.$$

This is a contact form with

$$\eta \wedge d\eta = \frac{-dx \wedge dy \wedge dm}{4(1 - x^2 - y^2)(1 + m^2 - (mx - y)^2)}.$$

The metric $\frac{1}{4}g$ is an associated metric for the contact form η , and ξ is the characteristic vector field giving a contact metric structure on the projectivized tangent bundle.

10

Curvature Functionals on Spaces of Associated Metrics

In this chapter we discuss a number of curvature functionals defined on the spaces of associated metrics for both compact symplectic and compact contact manifolds. Since these spaces are smaller than the space of Riemannian metrics of the same total volume, one expects for the classical curvature functionals weaker but still interesting critical point conditions. Other functionals that depend on the symplectic and contact structures are also considered.

10.1 Introduction to critical metric problems

The study of the integral of the scalar curvature, $A(g) = \int_M \tau \, dV_g$, as a functional on the set \mathcal{M}_1 of all Riemannian metrics of the same total volume on a compact orientable manifold M is now classical, dating back to Hilbert [1915] (see also Nagano [1967]). A Riemannian metric g is a critical point of $A(g)$ if and only if g is an Einstein metric. Since there are so many Riemannian metrics on a manifold, one can regard, philosophically, the finding of critical metrics as an approach to searching for the best metric for the given manifold. Other functions of the curvature have been taken as integrands as well, most notably $B(g) = \int_M \tau^2 \, dV_g$, $C(g) = \int_M |\rho|^2 \, dV_g$, where ρ is the Ricci tensor, and

$D(g) = \int_M |R_{kjih}|^2 dV_g$; the critical point conditions for these have been computed by Berger [1970]. From the critical point conditions it is easy to see that Einstein metrics are critical for $B(g)$ and $C(g)$ but not necessarily conversely. For example, an η -Einstein manifold M^{2n+1} with scalar curvature equal to $2n(2n+1)$ or $2n(2n+3)$ is a non-Einstein critical metric of $C(g)$, Yamaguchi and Chūman [1983]. In the case of $B(g)$, Yamaguchi and Chūman showed that a Sasakian critical point is Einstein.

Another area of interest in the functional $B(g)$ is the study of extremal Kähler metrics initiated by Calabi in [1954], [1982] and [1985]. Let (M, J, g) be a compact Kähler manifold with fundamental 2-form Ω and let $\mathcal{M}_{[\Omega]}$ denote the set of Kähler metrics on M whose fundamental 2-forms belong to the same cohomology class as Ω . Calabi defined an *extremal metric* as a critical point of $B(g)$ restricted to $\mathcal{M}_{[\Omega]}$ and proved that a Kähler metric is extremal if and only if the gradient of the scalar curvature is a (real) holomorphic vector field, i.e., an infinitesimal automorphism of the complex structure (see also Besse [1987, p. 334]). In a developing area, Boyer, Galicki and Simanca [2008], [2009] introduced the notion of an *extremal Sasakian structure* by considering the functional $B(g)$ on the set of Sasakian structures with the same characteristic vector field and complex normal bundle as a given Sasakian structure. Unfortunately, a detailed treatment of these topics would take us too far afield, and we refer the interested reader to the references in this paragraph.

Metrics of constant curvature and Kähler metrics of constant holomorphic curvature are critical for $D(g)$, see Muto [1975]. Also, a Sasakian manifold of dimension m and constant ϕ -sectional curvature $3m-1$ is critical for $D(g)$, see Yamaguchi and Chūman [1983].

To introduce the techniques for our study we will prove the classical result that a Riemannian metric is critical for $A(g)$ if and only if it is Einstein. Let M be a compact orientable manifold and \mathcal{M}_1 the set of all Riemannian metrics normalized by the condition of having the same total volume, usually taken to be 1, but one need not insist on the particular value in a given problem. As in Section 4.3 we often denote by the same letter a tensor field of type $(0, 2)$ and its corresponding types $(1, 1)$ and $(2, 0)$ determined by the metric under consideration, e.g., we may write $\text{tr}TD = T^i_j D^j_i = T^{ij} D_{ji}$.

Lemma 10.1 *Let T be a second order symmetric tensor field on M . Then $\int_M \text{tr} T D dV_g = 0$ for all symmetric tensor fields D satisfying $\int_M \text{tr} D dV_g = 0$ if and only if $T = cg$ for some constant c .*

Proof. If $T = cg$, $\text{tr} T D = c \text{tr} D$ and the sufficiency is immediate. Thus we have only to prove the necessity. Let X, Y be an orthonormal pair of vector fields on a neighborhood \mathcal{U} on M and f a C^∞ function with compact support in \mathcal{U} . Regarding X and Y as part of a local orthonormal basis, define a tensor field D on M by $D(X, X) = f$ and $D(Y, Y) = -f$, with all other components equal to zero and $D \equiv 0$ outside \mathcal{U} . Then $\int_M (T(X, X) - T(Y, Y)) f dV_g = 0$ for any C^∞ function with compact support and hence $T(X, X) = T(Y, Y)$ for every orthonormal pair X, Y . Therefore $T = cg$ for some function c and it remains to show that c is a constant. To see this, let X be any vector field and $D = \mathcal{L}_X g$. Then since the integral of a divergence vanishes,

$$0 = \int_M T^{ij} (\nabla_i X_j + \nabla_j X_i) dV_g = -2 \int_M (\nabla_i T^{ij}) X_j dV_g,$$

but X , is arbitrary so that $\nabla_i T^{ij} = 0$ (Lemma 4.7), from which we see that c must be a constant. ■

Now the approach to these critical point problems is to differentiate the functional in question along a path of metrics. The curvature functionals we study are not generally invariant under homothetic transformations; so when necessary we normalize these problems by restricting them to \mathcal{M}_1 . Let $g(t)$ be a path of metrics in \mathcal{M}_1 and

$$D_{ij} = \left. \frac{\partial g_{ij}}{\partial t} \right|_{t=0}$$

its tangent vector at $g = g(0)$. We define two other tensor fields by

$$\begin{aligned} D_{ji}{}^h &= \frac{1}{2} (\nabla_j D_i{}^h + \nabla_i D_j{}^h - \nabla^h D_{ji}), \\ D_{kji}{}^h &= \nabla_k D_{ji}{}^h - \nabla_j D_{ki}{}^h, \end{aligned}$$

where ∇ denotes the Riemannian connection of $g(0)$, and we note that

$$D_{ji}{}^h = \left. \frac{\partial \Gamma_{ji}{}^h}{\partial t} \right|_{t=0}, \quad D_{kji}{}^h = \left. \frac{\partial R_{kji}{}^h}{\partial t} \right|_{t=0},$$

where $\Gamma_{ji}{}^h$ and $R_{kji}{}^h$ denote the Christoffel symbols and curvature tensor of $g(t)$.

Theorem 10.1 *Let M be a compact orientable C^∞ manifold and \mathcal{M}_1 the set of all Riemannian metrics on M with unit volume. Then $g \in \mathcal{M}_1$ is a critical point of $A(g) = \int_M \tau dV_g$ if and only if g is Einstein.*

Proof. The proof is to compute $\frac{dA}{dt}$ at $t = 0$ for a path $g(t)$ in \mathcal{M}_1 . First note that from $g_{ij}g^{jk} = \delta_i^k$,

$$\left. \frac{\partial g^{ij}}{\partial t} \right|_{t=0} = -D^{ij}.$$

Differentiation of the volume element gives

$$\begin{aligned} \frac{d}{dt} dV_g &= \frac{d}{dt} \sqrt{\det(g(t))} dx^1 \wedge \cdots \wedge dx^n = \frac{1}{2\det(g(t))} \left(\frac{d}{dt} \det(g(t)) \right) dV_g \\ &= \frac{1}{2} g^{ij} \left(\frac{d}{dt} g_{ij} \right) dV_g = \frac{1}{2} D_i^i dV_g. \end{aligned}$$

Now

$$\begin{aligned} \left. \frac{dA}{dt} \right|_{t=0} &= \left. \frac{d}{dt} \int_M R_{kji}{}^k g^{ji} dV_g \right|_{t=0} \\ &= \int_M (D_{kji}{}^k g^{ji} - \rho_{ji} D^{ji} + \frac{1}{2} \tau g^{ji} D_{ji}) dV_g \\ &= \int_M (-\rho^{ji} + \frac{1}{2} \tau g^{ji}) D_{ji} dV_g, \end{aligned}$$

since the integral of a divergence vanishes. On the other hand, differentiation of $\int_M dV_g = 1$ gives $\int_M D_i^i dV_g = 0$. Thus setting $\left. \frac{dA}{dt} \right|_{t=0} = 0$ and applying Lemma 10.1, we have

$$\rho_{ji} - \frac{1}{2} \tau g_{ji} = c g_{ji}$$

for some constant c and hence that g is Einstein. The converse is immediate. ■

In [1974a] Y. Muto computed the second derivative of $A(g)$ at a critical point and proved the following theorem.

Theorem 10.2 *The index of $A(g)$ and the index of $-A(g)$ are both positive at each critical point.*

Y. Muto also considered the second derivative of the functional $D(g)$ from the following point of view. Let $Diff$ denote the diffeomorphism group of M ; if $f \in Diff$, then $D(f^*g) = D(g)$, and hence we have an induced mapping $\tilde{D} : \mathcal{M}_1/Diff \rightarrow \mathbb{R}$. We say that a metric g is a critical point of \tilde{D} if its orbit under $Diff$ is a critical point of \tilde{D} . Recall from the introduction to this section that a Riemannian metric of constant curvature is a critical point of $D(g)$; Y. Muto [1974b] proved the following result.

Theorem 10.3 *If M is diffeomorphic to a sphere and g_0 is a metric of positive constant curvature, then the index of $D(g)$ and the index of \tilde{D} are both zero at g_0 and \tilde{D} has a local minimum at g_0 .*

We now turn to integral functionals defined on the set of metrics associated to a symplectic or contact structure. To begin, we recall from Chapter 4 how the set \mathcal{A} of associated metrics sits in the set \mathcal{N} of all Riemannian metrics with the same volume element. In particular, we saw that a symmetric tensor field D is tangent to a path in \mathcal{A} at g if and only if

$$DJ + JD = 0$$

in the symplectic case and

$$D\xi = 0, \quad D\phi + \phi D = 0$$

in the contact case.

Similar to the role played by Lemma 10.1 we have the following lemma for critical point problems on \mathcal{A} .

Lemma 10.2 *Let T be a second order symmetric tensor field on M . Then $\int_M T^{ij} D_{ij} dV = 0$ for all symmetric tensor fields D satisfying $DJ + JD = 0$ in the symplectic case and $D\xi = 0$, $D\phi + \phi D = 0$ in the contact case if and only if $TJ = JT$ in the symplectic case and $\phi T - T\phi = \eta \otimes \phi T\xi - (\eta \circ T\phi) \otimes \xi$ in the contact case (i.e., ϕ and T commute when restricted to the contact subbundle).*

Proof. We give the proof in the symplectic case, the proof in the contact case being similar (see A. J. Ledger and the author [1986]). Let X_1, \dots, X_{2n} be a local J -basis defined on a neighborhood \mathcal{U} (i.e., X_1, \dots, X_{2n} is an orthonormal basis with respect to g and $X_{2i} = JX_{2i-1}$) and note that the first vector field X_1 may be any unit vector field on \mathcal{U} . Let f be a C^∞ function with compact support in \mathcal{U} and define a path

of metrics $g(t)$ as follows. Make no change in g outside \mathcal{U} and within \mathcal{U} change g only in the planes spanned by X_1 and X_2 by the matrix

$$\begin{pmatrix} 1 + tf + \frac{1}{2}t^2 f^2 & \frac{1}{2}t^2 f^2 \\ \frac{1}{2}t^2 f^2 & 1 - tf + \frac{1}{2}t^2 f^2 \end{pmatrix}.$$

It is easy to check that $g(t) \in \mathcal{A}$, and clearly the only nonzero components of D are $D_{11} = -D_{22} = f$. Then $\int_M T^{ij} D_{ij} dV = 0$ becomes

$$\int_M (T^{11} - T^{22}) f dV = 0.$$

Thus since X_1 was any unit vector field on \mathcal{U} ,

$$T(X, X) = T(JX, JX)$$

for any vector field X . Since T is symmetric, linearization gives $TJ = JT$. Conversely, if T commutes with J and D anticommutes with J , then $\text{tr}TD = \text{tr}TJDJ = \text{tr}JTDJ = -\text{tr}TD$, giving $T^{ij} D_{ij} = 0$. ■

We end this section with our first main result (S. Ianus and the author [1986]), namely we consider the functional $A(g)$ restricted to the set \mathcal{A} and find the critical point condition. Since \mathcal{A} is a smaller set of metrics than \mathcal{M}_1 , we expect a weaker critical point condition than that of being Einstein; we will see that the condition is that the Ricci operator commute with the corresponding almost complex structure, and hence we still have a very natural condition.

Theorem 10.4 *Let M be a compact symplectic manifold and \mathcal{A} the set of metrics associated to the symplectic form. Then $g \in \mathcal{A}$ is a critical point of $A(g) = \int_M \tau dV_g$ restricted to \mathcal{A} if and only if the Ricci operator Q of g commutes with the almost complex structure corresponding to g .*

Proof. The proof is again to compute $\frac{dA}{dt}$ at $t = 0$ for a path $g(t)$ in \mathcal{A} . Since all associated metrics have the same volume element, this is easier than in the Riemannian case. In particular, we have

$$\begin{aligned} \left. \frac{dA}{dt} \right|_{t=0} &= \left. \frac{d}{dt} \int_M R_{kji}{}^k g^{ji} dV_g \right|_{t=0} \\ &= \int_M D_{kji}{}^k g^{ji} - \rho_{ji} D^{ji} dV_g \\ &= - \int_M \rho^{ji} D_{ji} dV_g, \end{aligned}$$

Since $D_{kji}{}^k g^{ji} = (\nabla_k D_{ji}{}^k) g^{ji} = \nabla_k (D_{ji}{}^k g^{ji})$ which is a divergence; note that $\text{tr} D = 0$ and hence $D_{ki}{}^k = 0$. Setting $\frac{dA}{dt}|_{t=0} = 0$, the result follows from Lemma 10.2. ■

The commutativity $QJ = JQ$ is equivalent to $\rho(JX, JY) = \rho(X, Y)$ and is often referred to as the *J-invariance of the Ricci tensor* or as the manifold having *Hermitian Ricci tensor*.

10.2 The $*$ -scalar curvature

In Section 7.2 we defined the $*$ -Ricci tensor and the $*$ -scalar curvature in contact geometry. In almost Hermitian geometry these are defined similarly by

$$\rho_{ij}^* = R_{iklt} J^{kl} J_j{}^t, \quad \tau^* = \rho_i{}^i{}^*.$$

On a Kähler manifold, $\rho_{ij}^* = \rho_{ij}$. The most important property of τ^* on an almost Kähler manifold is the analogue or forerunner of Proposition 7.7, viz.

$$\tau - \tau^* = -\frac{1}{2} |\nabla J|^2.$$

Therefore $\tau - \tau^* \leq 0$, with equality holding if and only if the metric is Kähler. Thus for M compact, Kähler metrics are maxima of the functional

$$K(g) = \int_M \tau - \tau^* dV$$

on \mathcal{A} , and hence it is natural to ask for the critical point condition in general. This was the main question of S. Ianus and the author in [1986]; the critical point condition for $K(g)$ turns out to be the same as for $A(g)$ on \mathcal{A} , viz. $QJ = JQ$.

Theorem 10.5 *Let M be a compact symplectic manifold and \mathcal{A} the set of metrics associated to the symplectic form. Then $g \in \mathcal{A}$ is a critical point of $K(g)$ if and only if $QJ = JQ$.*

At first it may seem surprising that $A(g)$ and $K(g)$ have the same critical point condition, but we will see in the course of our discussion that this is natural. The proof of Theorem 10.5 will then be an easy consequence of Theorem 10.4 and Theorem 10.6 below. It is also natural to ask whether Kähler metrics are the only critical points of $K(g)$; the answer to this is negative and a counterexample was given by Davidov

and Muškarov [1990] on the twistor space of a compact Einstein, self-dual 4-manifold with negative scalar curvature.

In [1969] S. I. Goldberg showed that if $R_{XY}JZ = JR_{XY}Z$ on an almost Kähler manifold, then the metric is Kähler and conjectured that a compact almost Kähler Einstein manifold is Kähler. This conjecture is still open, but under the additional assumption of nonnegative scalar curvature it was proved by Sekigawa [1987]. Without the assumption of compactness, the conjecture is false; nonexistence was shown by Armstrong [1998]; Nurowski and Przanowski [1999] gave an example of a 4-dimensional non-Kähler almost Kähler that is Ricci flat; and Apostolov, Draghici and Moroianu [2001] have given noncompact counterexamples to the conjecture in dimensions ≥ 6 . The scalar curvature in the twistor space example, mentioned above, of a non-Kähler, almost Kähler manifold satisfying $QJ = JQ$ is negative; thus one might ask in light of Sekigawa's result whether a compact almost Kähler manifold with $QJ = JQ$ and nonnegative scalar curvature is Kähler. Draghici [1999] answered this question affirmatively in dimension 4 and negatively in general dimension. As a partial result in general dimension, Draghici proved in [1994] that if M is a compact almost Kähler manifold with Hermitian Ricci tensor and if there exists a constant $\lambda \geq 0$ such that $\lambda \leq Ric(X) \leq 2\lambda$ for any direction X , then M is Kähler. Recently there has been interest in other curvature conditions that imply that an almost Kähler structure is Kähler, especially questions involving the Weyl curvature tensor; see, e.g., Kirchberg [2004] and Draghici and the author [2009].

The original proof of Theorem 10.5 was to proceed as in Theorem 10.4 and differentiate $-\tau^*$. This is very complicated and only after clever use of many identities does one see that differentiation of this term again yields a contribution of $-\rho^{ij}D_{ij}$ to the integrand and hence that one has the same critical point condition (precisely $2Q$ commuting with J). This suggests that if we consider the “total scalar curvature” $I(g) = \int_M \tau + \tau^* dV$, the contributions of each term to the derivative of the integrand would have canceled each other, and hence every metric in \mathcal{A} would have been a critical point; thus, since \mathcal{A} is path connected, $I(g)$ must be constant on \mathcal{A} and hence a symplectic invariant (the author [1991a]). We now prove this using only relatively short computations. The motivation for studying the integral of the sum of τ and τ^* lies in

the author's work with D. Perrone [1992] on critical point problems in the contact case and will be discussed later in this section.

Theorem 10.6 *Let M be a compact symplectic manifold. Then $\int_M \tau + \tau^* dV$ is a symplectic invariant and to within a constant is the cup product*

$$(c_1(M) \cup [\Omega]^{n-1})([M]),$$

where $c_1(M)$ is the first Chern class of M .

Proof. Consider an almost Hermitian manifold with structure tensors (J, g) . The *generalized Chern form* γ is given by

$$8\pi\gamma_{ij} = -4J^k{}_j\rho_{ik}^* - J^{kl}(\nabla_j J^h{}_k)\nabla_i J_{lh}.$$

Now on an almost Kähler manifold,

$$(\nabla_k J_{ip})J_j{}^p = (\nabla_p J_{ij})J_k{}^p;$$

this is the condition for an almost Hermitian structure to be *quasi-Kähler* (see e.g., Gray and Hervella [1980]). Using this we have the following computation:

$$\begin{aligned} 8\pi\gamma_{ij}J^{ji} &= 4\tau^* - J^{kl}(\nabla^j J_k{}^h)(\nabla_i J_{hl})J_j{}^i \\ &= 4\tau^* - J^{kl}(\nabla^j J_k{}^h)(\nabla_j J_{hi})J_l{}^i \\ &= 4\tau^* - |\nabla J|^2, \end{aligned}$$

but $\tau - \tau^* = -\frac{1}{2}|\nabla J|^2$, and hence

$$2(\tau + \tau^*) = 4\tau^* - |\nabla J|^2 = 8\pi\gamma_{ij}J^{ji}.$$

Thus the “total scalar curvature” of an associated metric becomes $I(g) = 4\pi \int_M \gamma_{ij}J^{ji} dV$. Moreover, writing Ω in terms of a J -basis, a simple computation shows that

$$\int_M \gamma_{ij}J^{ji} dV = \frac{1}{2^{n-1}(n-1)!} \int_M \gamma \wedge \Omega^{n-1}.$$

■

Thus one has a relatively easy proof that the “total scalar curvature” is a symplectic invariant, and writing $\tau - \tau^*$ as $2\tau - (\tau + \tau^*)$, we see that $A(g)$ and $K(g)$ have the same critical point condition, proving Theorem 10.5.

On a Kähler manifold the Ricci form is, up to a constant, the first Chern form γ , i.e., $2\pi\gamma_{ij} = -\rho_{ik}J^k_j$. On an almost Kähler manifold the Ricci tensor need not be J -invariant and hence the Ricci form is not in general defined. In [1994] Draghici defined a Ricci form on an almost Kähler manifold by decomposing the Ricci tensor ρ into its J -invariant and J -anti-invariant parts and defining the *Ricci form* ψ by $\psi(X, Y) = \rho^{inv}(X, JY)$. Similarly define the **-Ricci form* by $\psi^*(X, Y) = \rho^*(X, JY)$; since $\rho^*(JX, JY) = \rho^*(Y, X)$, ψ^* is a well-defined 2-form. Draghici then obtained a cohomological version of the Goldberg conjecture and proved that if M is a compact almost Kähler manifold whose Ricci form is cohomologous to the first Chern class, then M is a Kähler manifold.

In the contact case the results corresponding to Theorems 10.4 and 10.5 were obtained by A. J. Ledger and the author in [1986], and the critical point conditions are different. The integral $K(g)$ for a contact manifold M^{2n+1} is taken to be $K(g) = \int_M \tau - \tau^* - 4n^2 dV$ (see Proposition 7.7) even though it is not necessary to include the constant $4n^2$.

Theorem 10.7 *Let M be a compact contact manifold and \mathcal{A} the set of metrics associated to the contact form. Then $g \in \mathcal{A}$ is a critical point of $A(g)$ restricted to \mathcal{A} if and only if Q and ϕ commute when restricted to the contact subbundle.*

Theorem 10.8 *Let M be a compact contact manifold and \mathcal{A} the set of metrics associated to the contact form. Then $g \in \mathcal{A}$ is a critical point of $K(g)$ if and only if $Q - 2nh$ and ϕ commute when restricted to the contact subbundle.*

Moreover, from Proposition 7.7 we see that Sasakian metrics, when they exist, are maxima of $K(g)$.

In approaching Theorem 10.6 we remarked that the study of $\int_M \tau + \tau^* dV$ in symplectic geometry was motivated by the corresponding study in contact geometry (Perrone and the author [1992]). It is interesting to remark that many results in contact and Sasakian geometry were motivated by the corresponding ones in symplectic and Kähler geometry. Here we have an example of a result in contact geometry preceding its symplectic analogue. In contact geometry the functional $I(g) = \int_M \tau + \tau^* dV$ is not an invariant and gives a critical point problem whose critical point condition gives the important class of K-contact metrics, i.e., associated metrics for which the characteristic vector field generates isometries.

Theorem 10.9 *Let M be a compact contact manifold and \mathcal{A} the set of metrics associated to the contact form. Then $g \in \mathcal{A}$ is a critical point of $I(g) = \int_M \tau + \tau^* dV$ if and only if g is K -contact.*

Proof. The terms τ and τ^* are differentiated as in Ledger and the author [1986], and the contributions of $\rho^{ij} D_{ij}$ cancel instead of doubling up as in Theorem 10.8 and the original proof of Theorem 10.5. The result is that

$$\left. \frac{d}{dt} \int_M \tau + \tau^* dV \right|_{t=0} = -4n \int_M h^{jl} D_{jl} dV.$$

Thus from Lemma 10.2 and $h\xi = 0$, the critical point condition becomes $\phi h - h\phi = 0$; but $\phi h + h\phi = 0$ and hence $h = 0$. Therefore ξ is a Killing vector field (see Section 6.2). ■

Turning to the second variation, we have the following result of Perrone and the author [1995].

Theorem 10.10 *The index of $I(g)$ and the index of $-I(g)$ are both positive at each critical point.*

Proof. Referring to Perrone and the author [1995] for details, the second derivative of $I(g)$ evaluated at a critical point is given by the following succinct formula:

$$I''(0) = 2n \int_M \text{tr}(\phi D \mathcal{L}_\xi D) dV.$$

Now as in the proof of Lemma 10.2, let X_1, \dots, X_{2n}, ξ be a local ϕ -basis defined on a neighborhood \mathcal{U} and again note that the first vector field X_1 may be any unit vector field on \mathcal{U} orthogonal to ξ . Let f be a C^∞ function with compact support in \mathcal{U} and define a path of metrics $g(t)$ as follows. Make no change in g outside \mathcal{U} and within \mathcal{U} change g only in the planes spanned by X_1 and X_2 by the matrix

$$\begin{pmatrix} 1 + t^2 f^2 & tf \\ tf & 1 \end{pmatrix}.$$

It is easy to check that $g(t) \in \mathcal{A}$, and clearly the only nonzero components of D are $D_{12} = D_{21} = f$. Denoting the first vector field in the ϕ -basis by X , calculation of $I''(0)$ in the above formula yields

$$I''(0) = -4n \int_M f^2 \eta([[\xi, X], X]) dV,$$

where X may be regarded as any unit vector field on \mathcal{U} belonging to the contact subbundle. Thus the proof reduces to finding unit vector fields belonging to the contact subbundle on \mathcal{U} for which $\eta([\xi, X], X)$ has either sign, and again we refer to Perrone and the author [1995] for details. ■

As an application of Theorem 10.10 we prove the following result (Perrone and the author [1995]).

Theorem 10.11 *The functional $A(g)$ restricted to \mathcal{A} cannot have a local minimum at any Sasakian metric.*

Proof. Suppose that g_0 is a Sasakian metric and a local minimum of $A(g)$ in \mathcal{A} . Then there exists a neighborhood \mathcal{U} of $g_0 \in \mathcal{A}$ on which $A(g_0) \leq A(g)$. Since all associated metrics have the same volume element, $\int_M \tau_0 dV \leq \int_M \tau dV$ for every $g \in \mathcal{U}$. From Proposition 7.7,

$$2\tau - 4n^2 \leq \tau + \tau^*,$$

with equality if and only if the metric is Sasakian. Thus we have

$$I(g_0) = \int_M 2\tau_0 - 4n^2 dV \leq \int_M 2\tau - 4n^2 dV \leq I(g)$$

for every $g \in \mathcal{U}$, that is, g_0 is a local minimum for $I(g)$, contradicting Theorem 10.10. ■

10.3 The integral of $Ric(\xi)$

The integral $L(g) = \int_M Ric(\xi) dV$ was studied in general dimension by the author in [1984] and independently by Chern and Hamilton in [1985] in the 3-dimensional case. Recall (Corollary 7.1) that

$$Ric(\xi) = 2n - \text{tr} h^2;$$

thus K-contact metrics, when they occur, are maxima for $L(g)$ on \mathcal{A} . Moreover, the critical point question for $L(g)$ is the same as that for $\int_M |h|^2 dV$ or $\int_M |T|^2 dV$, where $T(X, Y) = (\mathcal{L}_\xi g)(X, Y) = 2g(X, h\phi Y)$. It is the integral $E(g) = \int_M |T|^2 dV$ that was studied by Chern and Hamilton [1985] for 3-dimensional contact manifolds as a functional on \mathcal{A} regarded as the set of CR-structures on M (there was an error in their calculation of the critical point condition, as was pointed out by Tanno [1989]). The first result concerning $L(g)$ is the following.

Theorem 10.12 *Let M be a compact regular contact manifold and \mathcal{A} the set of metrics associated to the contact form. Then $g \in \mathcal{A}$ is a critical point of $L(g) = \int_M Ric(\xi) dV$ if and only if g is K -contact.*

Proof. As with our other critical point problems, the first step is to compute $\frac{dL}{dt}$ at $t = 0$ for a path $g(t) \in \mathcal{A}$:

$$\left. \frac{dL}{dt} \right|_{t=0} = \int_M (-h^i_m h^{mk} - R^k_{rs}{}^i \xi^r \xi^s + 2h^{ik}) D_{ik} dV.$$

Thus if $g(0)$ is a critical point, Lemma 10.2 gives

$$R_X \xi \xi = -\phi^2 X - h^2 X + 2hX$$

as the critical point condition. Using the first formula of Proposition 7.1, this becomes

$$(\nabla_\xi h)X = -2\phi hX.$$

From this we see that the eigenvalues of h are constant along the integral curves of ξ and that for an eigenvalue $\lambda \neq 0$ and unit eigenvector X , $g(\nabla_\xi X, \phi X) = -1$.

If now M is a regular contact manifold, then M is a principal circle bundle with ξ tangent to the fibers; locally, M is $\mathcal{U} \times S^1$. Since $h\phi + \phi h = 0$, we may choose an orthonormal ϕ -basis of eigenvectors of h at some point of $\mathcal{U} \times S^1$, say X_{2i-1} , $X_{2i} = \phi X_{2i-1}$, ξ . Since the eigenvalues are constant along the fiber, we can continue this basis along the fiber with at worst a change of orientation of some of the eigenspaces when we return to the starting point. Thus if Y is a vector field along the fiber, we may write

$$Y = \sum_i (\alpha_{2i-1} X_{2i-1} + \beta_{2i} X_{2i}) + \gamma \xi,$$

where the coefficients are periodic functions.

Now suppose that the critical point g is not a K -contact metric. Since ϕ and h anticommute, we may assume that all the λ_{2i-1} , $i = 1, \dots, n$, are nonnegative. Also from $(\nabla_\xi h)X = -2\phi hX$ it is easy to see that if some of the λ_{2i-1} vanish, the zero eigenspace of h is parallel along ξ , and hence we may choose the corresponding X_{2i-1} and X_{2i} parallel along a fiber. Again since M is regular, we may choose a vector field Y on $\mathcal{U} \times S^1$ such that at least some $\alpha_{2i-1} \neq 0$ for some $\lambda_{2i-1} \neq 0$ and Y is horizontal and

projectable, that is, $[\xi, Y] = 0$. Writing $Y = \sum_i (\alpha_{2i-1} X_{2i-1} + \beta_{2i} X_{2i})$ along a fiber we have using $\nabla_X \xi = -\phi X - \phi h X$,

$$\begin{aligned} 0 &= [\xi, Y] = \nabla_\xi Y - \nabla_Y \xi \\ &= \sum_i ((\xi \alpha_{2i-1}) X_{2i-1} + \alpha_{2i-1} \nabla_\xi X_{2i-1} + (\xi \beta_{2i}) X_{2i} + \beta_{2i} \nabla_\xi X_{2i} \\ &\quad + \alpha_{2i-1} X_{2i} + \lambda_{2i-1} \alpha_{2i-1} X_{2i} - \beta_{2i} X_{2i-1} + \lambda_{2i-1} \beta_{2i} X_{2i-1}). \end{aligned}$$

Taking components, we have

$$\begin{aligned} 0 &= \xi \alpha_{2j-1} + \sum_i \alpha_{2i-1} g(\nabla_\xi X_{2i}, X_{2j}) + \lambda_{2j-1} \beta_{2j}, \\ 0 &= \xi \beta_{2j} + \sum_i \beta_{2i} g(\nabla_\xi X_{2i}, X_{2j}) + \lambda_{2j-1} \alpha_{2j-1}. \end{aligned}$$

Multiplying the first of these by β_{2j} , the second by α_{2j-1} , and summing on j , we have

$$\xi \left(\sum_j \alpha_{2j-1} \beta_{2j} \right) = - \sum_j \lambda_{2j-1} (\alpha_{2j-1}^2 + \beta_{2j}^2) \leq 0.$$

Thus $\sum_j \alpha_{2j-1} \beta_{2j}$ is a nonincreasing, nonconstant function along the integral curve, contradicting its periodicity. ■

One might conjecture Theorem 10.12 without the regularity. However, we have the following counterexample: The standard contact metric structure on the tangent sphere bundle of a compact manifold of constant curvature -1 is a critical point of L but is not K-contact (the author [1991b]). We give this in the next theorem. Recall also the result of Tashiro, Theorem 9.3, that the standard contact metric structure on the tangent sphere bundle of a Riemannian manifold is K-contact if and only if the base manifold is of constant curvature $+1$.

Theorem 10.13 *Let $T_1 M$ be the tangent sphere bundle of a compact Riemannian manifold (M, G) and \mathcal{A} the set of all Riemannian metrics associated to its standard contact structure. Then the standard associated metric is a critical point of the functional $L(g)$ if and only if (M, G) is of constant curvature $+1$ or -1 .*

Proof. We have seen that the critical point condition of $L(g)$ is

$$R_X \xi \xi = -\phi^2 X - h^2 X + 2hX$$

or $\phi^2 + h^2 + l = 2h$ ($lX = R_X \xi$). Applying this to a vertical vector $U \in T_t T_1 M$ as in the proof of Lemma 9.1, we find that (M, G) is locally symmetric by the Lemma of Cartan [1983, pp.257–258] and that $L_t^2 X = X$ for any orthonormal pair $\{X, t\}$ on (M, G) . Thus the only eigenvalues of $L_t : [t]^\perp \rightarrow [t]^\perp$ are ± 1 . The manifold M is irreducible; for if M had a locally Riemannian product structure, then choosing t tangent to one factor and X tangent to the other, we would have $\mathbf{R}_X t = 0$, contradicting the fact that the only eigenvalues of L_t are ± 1 . Now the sectional curvature of an irreducible locally symmetric space does not change sign. Thus if for some t , L_t had both $+1$ and -1 as eigenvalues, there would be sectional curvatures equal to $+1$ and -1 . Consequently, only one eigenvalue can occur, and hence (M, G) must be a space of constant curvature $+1$ or -1 .

Conversely, if (M, G) has constant curvature c , our expressions for h on horizontal vectors X orthogonal to ξ and vertical vectors U in Section 9.2, namely

$$hX = -X + (\mathbf{R}_{\pi_* X t})^H, \quad hU = U - (\mathbf{R}_{KU t})^V,$$

become $hX = (c - 1)X$ and $hU = (1 - c)U$. Similarly $lU = c^2 U$ and $lX = (4c - 3c^2)X$. Substituting these into the critical point condition, we see that it is satisfied if and only if $c = \pm 1$. ■

In [1991] S. Deng studied the second variation of the functional $L(g)$, or equivalently, of $E(g) = \int_M |T|^2 dV$.

Theorem 10.14 *Let $g \in \mathcal{A}$ be a critical point of $E(g)$. Then g is a minimum.*

Proof. As with Theorem 10.10 we will sketch the proof and refer to Deng [1991] for details. The first step is a lengthy calculation of $E''(0)$ yielding

$$E''(0) = 2 \int_M |\mathcal{L}_\xi D|^2 dV \geq 0.$$

The second step is an auxiliary result that if $\mathcal{L}_\xi D = 0$, then $|T|^2$ is constant along the geodesics $g(t) = ge^{Dt}$ in \mathcal{A} . Now proceed as follows. If $\mathcal{L}_\xi D = 0$ for all D , then $|T|$ is constant and hence $E(g)$ is constant. So suppose that g is not a minimum and that there exist D such that $\mathcal{L}_\xi D \neq 0$. Let V be the subspace of $T_g \mathcal{A}$ consisting of those D with $\mathcal{L}_\xi D = 0$ and V^\perp its orthogonal complement. Let \mathcal{U} be a neighborhood

of g in $\exp_g V^\perp$ (cf. Ebin [1970, p. 37]) on which E exceeds $E(g)$. Let \mathcal{W} be a neighborhood of $\mathcal{U} \subset \mathcal{A}$ formed by geodesic arcs in directions belonging to V . Since g is not a minimum, there exists $\bar{g} \in \mathcal{W}$ such that $E(\bar{g}) < E(g)$. Then there is a geodesic γ from \bar{g} in a direction $\bar{D} \in V$ that meets \mathcal{U} at say \hat{g} . E is constant along γ and therefore $E(\hat{g}) < E(g)$, contradicting $E(\hat{g}) > E(g)$ for $\hat{g} \in \mathcal{U}$. ■

We have noted a number of times a role played by conditions of the form $\nabla_\xi h = ah\phi$ for some function a , and here the critical point condition for the functional L , or equivalently E , is of this type, viz. $\nabla_\xi h = 2h\phi$, equivalently $\nabla_\xi T - 2T \cdot \phi = 0$, where $T \cdot \phi(X, Y) = T(X, \phi Y)$. Barletta and Dragomir [2001] pointed out that this condition has a “universality” property for functionals depending on $|T|^2$ and refer to $\nabla_\xi T - 2T \cdot \phi = 0$ as *Tanno’s equation*. Specifically, let $f : \mathbb{R} \rightarrow \mathbb{R}$ be a real analytic function and set $t = |T|^2$. Then $g \in \mathcal{A}$ is a critical point of the functional $E_f(g) = \int_M f(t) dV$ on \mathcal{A} if and only if

$$f''(t)(\xi t)T + f'(t)(\nabla_\xi T - 2T \cdot \phi) = 0.$$

In [1992b] D. Perrone considered the integral

$$F(g) = \int_M \tau + \tau^* + 2n \operatorname{Ric}(\xi) dV,$$

or setting $\tau_1 = \tau^* + 2n \operatorname{Ric}(\xi)$, $F(g) = \int_M \tau + \tau_1 dV$ on \mathcal{A} . In dimension 3, letting $K(\mathcal{D})$ denote the sectional curvature of the contact subbundle \mathcal{D} , $\tau = 2K(\mathcal{D}) + 2 \operatorname{Ric}(\xi)$, and $\tau^* = 2K(\mathcal{D})$, so $\tau_1 = \tau$. In higher dimensions using Proposition 7.7,

$$\tau_1 = \tau + (2n - 1)(\operatorname{Ric}(\xi) - 2n) + \frac{1}{2}(|\nabla \phi|^2 - 4n),$$

and hence if the manifold is Sasakian, $\tau_1 = \tau$. The critical point condition for $F(g)$ is $\nabla_\xi h = 0$.

Little has been said about the existence of critical metrics, and in general this is a difficult problem. Recall the notion of an almost regular contact manifold as given by C. B. Thomas [1976] (Section 3.4). Rukimbira [1995c] proved that every almost regular contact manifold admits a critical metric for $E(g)$ (equivalently $L(g)$). A number of people, including this author, have raised the question of approaching the existence question as a Ricci flow problem on \mathcal{A} , but to date there are no definitive results.

10.3.1 H -contact manifolds

We have often encountered the condition that ξ be an eigenvector of the Ricci operator Q . One of the more important interpretations of this condition is that of an H -contact manifold as introduced by D. Perrone [2004], and this is a good place to discuss this concept. First, on a given compact m -dimensional Riemannian manifold (M, g) , a unit vector field X is said to be a *harmonic vector field* (C. M. Wood [1997]) if it is a critical point of the energy functional

$$\mathcal{E}(X) = \frac{m}{2} \text{vol}(M, g) + \frac{1}{2} \int_M \|\nabla X\|^2 dV$$

on the space of all unit vector fields. Perrone then defines an H -contact manifold to be a contact metric manifold for which the characteristic vector field ξ is a harmonic vector field and proves the following theorem.

Theorem 10.15 *A contact metric manifold is an H -contact manifold if and only if its characteristic vector field is an eigenvector of the Ricci operator.*

Note that from $\nabla \xi = -\phi - \phi h$, if we consider the energy \mathcal{E} as a functional on \mathcal{A} for a fixed contact form, and hence a fixed unit vector field ξ , then $g \in \mathcal{A}$ is critical for \mathcal{E} if and only if it is critical for the functional L .

In the same paper Perrone also proves the following theorem.

Theorem 10.16 *Let (M^{2n+1}, η, g) be a compact H -contact manifold such that g is critical for L . If $\rho + cg$ is positive definite for some constant $c < 2 - \frac{|\tau|}{\sqrt{2n}}$, the first Betti number of M^{2n+1} vanishes.*

Furthermore, in dimension 3, Perrone showed that a compact H -contact 3-manifold such that g is critical for L is either Sasakian or locally isometric to a non-Sasakian left-invariant contact metric structure on $SL(2, \mathbb{R})$ and conversely. In particular, using the classification of Geiges, Theorem 6.4, Perrone showed that a compact H -contact 3-manifold such that g is critical for L is diffeomorphic to a left quotient of $SU(2)$, the Heisenberg group or $\widetilde{SL}(2, \mathbb{R})$ by a discrete group.

By a result of Han and Yim [1998], a harmonic vector field X on a Riemannian manifold (M, g) defines a harmonic map into the tangent sphere bundle with the metric g' induced from the Sasaki metric on TM (see Section 9.2) if it satisfies the additional condition

$$\text{tr}(Z \longrightarrow -R_{\nabla_Z X, X} Y) = 0$$

for all vector fields Y on M . Perrone [2003] related this idea to the generalized (κ, μ) -manifolds of Koufogiorgos and Tsihlias [2000] by showing that a 3-dimensional contact metric manifold is a (κ, μ) -manifold on an everywhere open dense set if and only if the characteristic vector field defines a harmonic map into the tangent sphere bundle. Following up on these ideas, Koufogiorgos, Markellos and Papantoniou [2008] introduced the notion of a (κ, μ, ν) -manifold as a contact metric manifold whose curvature tensor satisfies

$$\begin{aligned} R_{XY}\xi &= \kappa(\eta(Y)X - \eta(X)Y) + \mu(\eta(Y)hX - \eta(X)hY) \\ &\quad + \nu(\eta(Y)h\phi X - \eta(X)h\phi Y) \end{aligned}$$

for functions κ , μ and ν and showed that for dimensions > 3 such a manifold is a (κ, μ) -manifold. However, in dimension 3 they proved that a (κ, μ, ν) -manifold is an H-contact manifold and conversely, a 3-dimensional H-contact manifold is a (κ, μ, ν) -manifold on an everywhere open dense set. This paper also contains examples of 3-dimensional (κ, μ, ν) -manifolds that are not generalized (κ, μ) -manifolds, and therefore for these examples the characteristic vector fields are harmonic vector fields but do not define harmonic maps.

10.4 The Webster scalar curvature

In Theorem 6.7 we saw that a contact metric manifold is a strongly pseudo-convex CR-manifold if and only if $(\nabla_X \phi)Y = g(X + hX, Y)\xi - \eta(Y)(X + hX)$. On a strongly pseudoconvex CR-manifold Tanaka [1976] introduced a canonical connection. In [1989] Tanno introduced the corresponding connection on a contact metric manifold called the *generalized Tanaka connection*; it agrees with the connection of Tanaka when the contact metric manifold is a strongly pseudoconvex (integrable) CR-manifold. This connection, denoted by ${}^*\nabla$, is defined by

$$\begin{aligned} {}^*\nabla_X Y &= \nabla_X Y + \eta(X)\phi Y - \eta(Y)\nabla_X \xi + (\nabla_X \eta)(Y)\xi \\ &= \nabla_X Y + \eta(X)\phi Y + \eta(Y)(\phi X + \phi hX) + d\eta(X, Y)\xi + d\eta(hX, Y)\xi. \end{aligned}$$

The torsion of this connection, *T , is given by

$$\begin{aligned} {}^*T(X, Y) &= \eta(X)\phi Y - \eta(Y)\phi X - \eta(Y)\nabla_X \xi + \eta(X)\nabla_Y \xi + 2d\eta(X, Y)\xi \\ &= \eta(Y)\phi hX - \eta(X)\phi hY + 2g(X, \phi Y)\xi. \end{aligned}$$

Tanno [1989] then proves the following proposition.

Proposition 10.1 *The generalized Tanaka connection ${}^*\nabla$ on a contact metric manifold is the unique linear connection such that*

$$\begin{aligned} {}^*\nabla\eta &= 0, \quad {}^*\nabla\xi = 0, \quad {}^*\nabla g = 0, \\ ({}^*\nabla_X\phi)Y &= (\nabla_X\phi)Y - g(X + hX, Y)\xi + \eta(Y)(X + hX), \\ {}^*T(\xi, \phi Y) &= -\phi^*T(\xi, Y), \\ {}^*T(X, Y) &= 2d\eta(X, Y)\xi, \quad \text{on } \mathcal{D}. \end{aligned}$$

Tanno also computed the curvature of ${}^*\nabla$ and upon contraction obtains the *generalized Tanaka–Webster scalar curvature*

$$W_1 = \tau - Ric(\xi) + 4n.$$

This is eight times the Webster scalar curvature as defined by Chern and Hamilton [1985] on 3-dimensional contact manifolds and as used by Perrone [1998] in Theorem 7.26.

We now prove a theorem of Chern and Hamilton [1985], an alternate proof of which was given by Perrone in [1990], and a theorem of Tanno [1989]. The proofs will be given simultaneously.

Theorem 10.17 (Chern–Hamilton) *Let M be a compact 3-dimensional contact manifold and \mathcal{A} the set of metrics associated to the contact form. Then $g \in \mathcal{A}$ is a critical point of $E_1(g) = \int_M W_1 dV$ if and only if g is K -contact.*

Theorem 10.18 (Tanno) *Let M be a compact contact manifold and \mathcal{A} the set of metrics associated to the contact form. Then $g \in \mathcal{A}$ is a critical point of $E_1(g) = \int_M W_1 dV$ if and only if*

$$(Q\phi - \phi Q) - (l\phi - \phi l) = 4\phi h - \eta \otimes \phi Q\xi + (\eta \circ Q\phi) \otimes \xi.$$

Proofs. Clearly it is enough to consider $\int_M \tau - Ric(\xi) dV$, and having computed the derivatives of each term separately in our previous theorems, we have

$$\left. \frac{d}{dt} \int_M \tau - Ric(\xi) dV \right|_{t=0} = \int_M (-\rho^{ki} + h^i_m h^{mk} + R^k_{rs} \xi^r \xi^s - 2h^{ik}) D_{ik} dV.$$

Thus by Lemma 10.2 we see that the critical point condition is

$$(Q\phi - \phi Q) - (l\phi - \phi l) = 4\phi h - \eta \otimes \phi Q\xi + (\eta \circ Q\phi) \otimes \xi.$$

Now in dimension 3, the Ricci operator determines the full curvature tensor, i.e.,

$$R_{XY}Z = (g(Y, Z)QX - g(X, Z)QY + g(QY, Z)X - g(QX, Z)Y) - \frac{\tau}{2}(g(Y, Z)X - g(X, Z)Y).$$

Therefore the operator l is given by

$$lX = QX - \eta(X)Q\xi + g(Q\xi, \xi)X - g(QX, \xi)\xi - \frac{\tau}{2}(X - \eta(X)\xi),$$

from which

$$(l\phi - \phi l)X = (Q\phi - \phi Q)X + \eta(X)\phi Q\xi - g(Q\phi X, \xi)\xi.$$

Combining this and the critical point condition, we have $4\phi h = 0$, and hence, since $h\xi = 0$, $h = 0$. ■

As we saw in the last section, in dimension 3, $\tau = 2K(\mathcal{D}) + 2Ric(\xi)$ and $\tau^* = 2K(\mathcal{D})$. Using this, $W_1 = \tau - Ric(\xi) + 4n$ becomes

$$W_1 = \frac{1}{2}(\tau + \tau^* + 8).$$

Thus in dimension 3 the critical point problem for $E_1(g)$ is the same as the critical point problem for $I(g)$ in Theorem 10.9, suggesting that

$$W = \frac{1}{2}(\tau + \tau^* + 4n(n+1))$$

may be the proper generalization of the Webster scalar curvature.

There are, however, other generalizations of the Webster scalar curvature. Again in dimension 3, we may write W_1 as $\tau^* + Ric(\xi) + 4$, and we define a second generalization of the Webster scalar curvature by

$$W_2 = \tau^* + Ric(\xi) + 4n^2.$$

In general dimension the critical point condition of $E_2(g) = \int_M W_2 dV$ is

$$(Q\phi - \phi Q) - (l\phi - \phi l) = -4(2n-1)\phi h - \eta \otimes \phi Q\xi + (\eta \circ Q\phi) \otimes \xi$$

(see Perrone and the author [1992]). The generalization of the Webster scalar curvature $W = \frac{1}{2}(\tau + \tau^* + 4n(n+1))$ is the average of W_1 and W_2 . Th. Koufogiorgos [1997b] has considered the difference of these, and we briefly mention his results.

Proposition 10.2 *Let M^{2n+1} be a contact metric manifold with a strongly pseudoconvex CR-structure. Then*

$$(Q\phi - \phi Q) = (l\phi - \phi l) + 4(n-1)h\phi - \eta \otimes \phi Q\xi + (\eta \circ Q\phi) \otimes \xi.$$

Theorem 10.19 *Let M^{2n+1} be a contact manifold with $n > 1$ and \mathcal{A} the set of associated metrics. If $g \in \mathcal{A}$ gives rise to a strongly pseudoconvex CR-structure, then g is a critical point of $\int_M W_1 - W_2 dV$.*

The idea of the proof of this theorem is to find the critical point condition of $\int_M W_1 - W_2 dV$, which is exactly the formula of Proposition 10.2.

Finally, a word is in order on the constants depending on dimension that occur in the definitions of W_1 , W_2 and W . Again recall the notion of a \mathcal{D} -homothetic deformation, namely a change of structure tensors of the form

$$\bar{\eta} = a\eta, \quad \bar{\xi} = \frac{1}{a}\xi, \quad \bar{\phi} = \phi, \quad \bar{g} = ag + a(a-1)\eta \otimes \eta,$$

where a is a positive constant. By direct computation one shows that τ , $Ric(\xi)$, and τ^* transform in the following manner:

$$\begin{aligned} \bar{\tau} &= \frac{1}{a}\tau + \frac{1-a}{a^2}Ric(\xi) - 2n\left(\frac{a-1}{a}\right)^2, \\ \overline{Ric}(\bar{\xi}) &= \frac{1}{a^2}(Ric(\xi) + 2n(a^2 - 1)), \\ \bar{\tau}^* &= \frac{1}{a}\tau^* + \frac{a-1}{a^2}Ric(\xi) + 2n\left(2n\left(\frac{1-a}{a}\right) + \frac{1-a^2}{a^2}\right). \end{aligned}$$

From these we see that $\bar{W}_1 = \frac{1}{a}W_1$, $\bar{W}_2 = \frac{1}{a}W_2$ and $\bar{W} = \frac{1}{a}W$.

10.5 A gauge invariant

Use has often been made of the notion of a \mathcal{D} -homothetic deformation, as we have seen. The notion of a gauge transformation of a contact metric structure has not received as much attention. No doubt this is due in part to computational complexity; nonetheless, comparing with the notion of a contact structure in the wider sense, the notion of a gauge transformation should be fundamental and deserving of attention.

Let $(M^{2n+1}, \phi, \xi, \eta, g)$ be a contact metric manifold and consider a gauge transformation $\bar{\eta} = \sigma\eta$ of the contact structure, where σ is a

positive function on M^{2n+1} . Let $\zeta = \frac{1}{2\sigma}\phi\nabla\sigma$, $\nabla\sigma$ being the g -gradient of the function σ , and let z be the covariant form of ζ . Now define new structure tensors $\bar{\xi}$, $\bar{\phi}$ and \bar{g} by

$$\begin{aligned}\bar{\xi} &= \frac{1}{\sigma}(\xi + \zeta), & \bar{\phi} &= \phi + \frac{1}{2\sigma}\eta \otimes (\nabla\sigma - (\xi\sigma)\xi), \\ \bar{g} &= \sigma(g - \eta \otimes z - z \otimes \eta) + \sigma(\sigma - 1 + |\zeta|^2)\eta \otimes \eta.\end{aligned}$$

Then $(\bar{\phi}, \bar{\xi}, \bar{\eta}, \bar{g})$ is a contact metric structure, and the change from (ϕ, ξ, η, g) to $(\bar{\phi}, \bar{\xi}, \bar{\eta}, \bar{g})$ is called a *gauge transformation* of the contact metric structure. When σ is a constant this is a \mathcal{D} -homothetic deformation. The notion of a gauge transformation in contact metric geometry is due to Sasaki [1965, p. PTSB-1-3] (see also Tanno [1989]).

Tanno [1989] computed the change of the generalized Tanaka–Webster scalar curvature under a gauge transformation. The result of his computation is

$$\bar{W}_1 = \frac{1}{\sigma} \left(W_1 - \frac{2(n+1)}{\sigma} (\Delta\sigma - \xi\xi\sigma) - \frac{(n+1)(n-2)}{\sigma^2} (|d\sigma|^2 - (\xi\sigma)^2) \right).$$

In the same paper Tanno generalizes an invariant of Jerison and Lee [1984] for strongly pseudoconvex CR-manifolds; specifically, he considers

$$\kappa_{(\eta,g)} = \inf \left\{ \int_M \left(\frac{4(n+1)}{n} (|df|^2 - (\xi f)^2) + W_1 f^2 \right) dV_g \right\},$$

where the infimum is taken over all nonnegative functions f such that $\int_M f^p dV_g = 1$, $p = 2 + \frac{2}{n}$, and proves that $\kappa_{(\eta,g)}$ is a gauge invariant.

Now for a compact contact manifold (M^{2n+1}, η) let $\mathcal{F}(p)$ be the set of all nonnegative functions f such that $\int_M f^p dV_g = 1$ and define a functional $F_\eta : \mathcal{M} \times \mathcal{F}(p) \rightarrow \mathbb{R}$ by

$$F_\eta(g, f) = \int_M \left(\frac{4(n+1)}{n} (|df|^2 - (\xi f)^2) + W_1 f^2 \right) dV_g.$$

Also define $F_{(\eta,g)} : \mathcal{F}(p) \rightarrow \mathbb{R}$ by $F_{(\eta,g)}(f) = F_\eta(g, f)$. In [1989] Tanno studied these functionals in detail; here we mention only the following result.

Theorem 10.20 *Let $(M^{2n+1}, \phi, \xi, \eta, g)$ be a compact contact metric manifold with constant generalized Tanaka–Webster scalar curvature W_1 .*

Then F_η is critical at the pair (g, f) with $f = (\text{vol}(M, g))^{-1/p}$ if and only if

$$(Q\phi - \phi Q) - (l\phi - \phi l) = 4\phi h - \eta \otimes \phi Q\xi + (\eta \circ Q\phi) \otimes \xi.$$

Note that this is also the critical point condition of $E_1(g) = \int_M W_1 dV$ in Theorem 10.18.

10.6 The Abbena metric as a critical point

In Section 1.1 we discussed Thurston's example of a compact symplectic manifold with no Kähler structure and a natural Riemannian metric on this manifold introduced by E. Abbena [1984]. She computed the curvature, and with respect to the basis $\{\mathbf{e}_i\}$ introduced in Section 1.1, the Ricci operator Q is given by the matrix

$$\begin{pmatrix} -\frac{1}{2} & 0 & 0 & 0 \\ 0 & -\frac{1}{2} & 0 & 0 \\ 0 & 0 & \frac{1}{2} & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix}.$$

From the expression for Q it is clear that (M, g) is not Einstein, nor is $QJ = JQ$. Thus this metric is not a critical point for $A(g)$ considered as a functional on \mathcal{M}_1 or on \mathcal{A} , or for $K(g)$ on \mathcal{A} . Also, $\tau = -1/2$ and $\tau^* = +1/2$, giving zero for the “total scalar curvature”.

In [1996] Park and Oh discussed a functional for which the Abbena metric on the Thurston manifold is a critical point; their results are given in the following theorem. Recall that \mathcal{M}_1 denotes the space of Riemannian metrics with unit volume.

Theorem 10.21 *The Abbena metric on the Thurston manifold is a critical point of the functional*

$$\int_M \left(\frac{4}{3} \text{tr} Q^3 - \tau \right) dV_g$$

on \mathcal{M}_1 . The index of this functional and its negative are both positive at the Abbena metric on the Thurston manifold.

We remark that the Abbena metric on the Thurston manifold is a critical point for $K(g)$ in a different context. C. M. Wood [1995] showed

that the Abbena metric is a critical point of $K(g) = -\frac{1}{2} \int_M |\nabla J|^2 dV$ defined with respect to variations through almost complex structures J that preserve g . For this problem the critical point condition is

$$[J, \nabla^* \nabla J] = 0,$$

where $\nabla^* \nabla J$ is the rough Laplacian of the metric in question.

11

Negative ξ -sectional Curvature

In this chapter we introduce some special directions that belong to the contact subbundle of a contact metric manifold with negative sectional curvature for plane sections containing the characteristic vector field. We also discuss in this chapter some questions concerning Anosov and conformally Anosov flows.

11.1 Special directions in the contact subbundle

In [1998] the author introduced some special directions belonging to the contact subbundle of a contact metric manifold with negative sectional curvature for plane sections containing the characteristic vector field ξ or more generally when the operator h admits an eigenvalue greater than 1. For simplicity we will often refer to the sectional curvature of plane sections containing the characteristic vector field ξ as *ξ -sectional curvature*.

We may regard the equation

$$\nabla_X \xi = -\phi X - \phi hX$$

of Lemma 6.2 as indicating how ξ or, by orthogonality, the contact subbundle rotates as one moves around on the manifold. For example,

when $h = 0$, as we move in a direction X orthogonal to ξ , ξ is always “turning” or “falling” toward $-\phi X$. If $hX = \lambda X$, then $\nabla_X \xi = -(1 + \lambda)\phi X$, and again ξ is turning toward $-\phi X$ if $\lambda > -1$ or toward ϕX if $\lambda < -1$. Recall that we noted above that if λ is an eigenvalue of h with eigenvector X , then $-\lambda$ is also an eigenvalue with eigenvector ϕX .

Now one can ask whether there can ever be directions, say Y orthogonal to ξ , along which ξ “falls” forward or backward in the direction of Y itself.

Theorem 11.1 *Let M^{2n+1} be a contact metric manifold. If the tensor field h admits an eigenvalue $\lambda > 1$ at a point p , then there exists a vector Y orthogonal to ξ at p such that $\nabla_Y \xi$ is collinear with Y . In particular, if M^{2n+1} has negative ξ -sectional curvature, such directions Y exist.*

Proof. Let λ denote a positive eigenvalue of h and X a corresponding unit eigenvector. Then

$$\nabla_X \xi = -(1 + \lambda)\phi X, \quad \nabla_{\phi X} \xi = (1 - \lambda)X.$$

Now let $Y = aX + b\phi X$ with $a > 0, b > 0, a^2 + b^2 = 1$ and suppose that $\nabla_Y \xi = \alpha Y$. Then

$$\alpha(aX + b\phi X) = \nabla_Y \xi = -(1 + \lambda)a\phi X + (1 - \lambda)bX,$$

from which $\alpha a = (1 - \lambda)b$, $\alpha b = -(1 + \lambda)a$ and hence

$$a^2 = \frac{\lambda - 1}{2\lambda}, \quad b^2 = \frac{\lambda + 1}{2\lambda}, \quad \alpha = -\sqrt{\lambda^2 - 1}.$$

Thus we see that directions along which $\nabla_Y \xi$ is collinear with Y exist whenever h admits an eigenvalue greater than 1. From the fact that $Ric(\xi) = 2n - \text{tr} h^2$ (Corollary 7.1) we see that if M^{2n+1} has negative ξ -sectional curvature, at least one of the eigenvalues of h must exceed 1. ■

Note that when there exists a direction Y along which $\nabla_Y \xi$ is collinear with Y as above, there is also a second such direction, namely $Z = aX - b\phi X$. For Z we have $\nabla_Z \xi = -\alpha Z$; thus we think of ξ as falling backward as we move in the direction Y and falling forward as we move in the direction Z .

Next note that

$$g(Y, Z) = a^2 - b^2 = -\frac{1}{\lambda}$$

and hence that such directions Y and Z are never orthogonal. Also, if λ has multiplicity $m \geq 1$, then there are m -dimensional subbundles \mathcal{Y} and \mathcal{Z} such that $\nabla_Y \xi = \alpha Y$ for any $Y \in \mathcal{Y}$ and $\nabla_Z \xi = -\alpha Z$ for any $Z \in \mathcal{Z}$. We refer to directions along which the covariant derivative of ξ is collinear with the direction as *special directions*.

11.2 Anosov flows

The most notable example of a contact manifold for which the characteristic vector field is Anosov is the tangent sphere bundle of a negatively curved manifold; here the characteristic vector field is (twice) the geodesic flow, as we saw in Section 9.2. In the case of the tangent sphere bundle of a surface, this is closely related to the structure on $SL(2, \mathbb{R})$ from both the topological and Anosov points of view. If we set $Z_2 = \left\{ \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}, \begin{pmatrix} -1 & 0 \\ 0 & -1 \end{pmatrix} \right\}$, then $PSL(2, \mathbb{R}) = SL(2, \mathbb{R})/Z_2$ is homeomorphic to the tangent sphere bundle of the hyperbolic plane. Moreover, the geodesic flow on a compact surface of constant negative curvature may be realized on $PSL(2, \mathbb{R})/\Gamma$ by $\left\{ \begin{pmatrix} e^t & 0 \\ 0 & e^{-t} \end{pmatrix} \right\}$, where Γ is a discrete subgroup of $SL(2, \mathbb{R})$ for which $SL(2, \mathbb{R})/\Gamma$ is compact (see e.g., Auslander, Green and Hahn [1963, pp. 26–27]). However, from the Riemannian point of view these examples are quite different, as we shall see. In fact, in the case of the tangent sphere bundle of a negatively curved surface, the special directions never agree with the stable and unstable directions of the Anosov flow (Theorem 11.3).

Classically an Anosov flow is defined as follows (Anosov [1967]). Let M be a compact differentiable manifold, ξ a nonvanishing vector field and $\{\psi_t\}$ its 1-parameter group of (C^k) diffeomorphisms. $\{\psi_t\}$ is said to be an *Anosov flow* (or ξ to be *Anosov*) if there exist subbundles E^s and E^u that are invariant along the flow and such that $TM = E^s \oplus E^u \oplus \{\xi\}$, and there exists a Riemannian metric such that

$$\begin{aligned} |\psi_{t*} Y| &\leq a e^{-ct} |Y| \text{ for } t \geq 0 \text{ and } Y \in E_p^s, \\ |\psi_{t*} Y| &\leq a e^{ct} |Y| \text{ for } t \leq 0 \text{ and } Y \in E_p^u, \end{aligned}$$

where a and c are positive constants independent of $p \in M$ and Y in E_p^s or E_p^u . The subbundles E^s and E^u are called the *stable* and *unstable* subbundles or the *contracting* and *expanding* subbundles. The subbundles

E^s and E^u are integrable with C^k integral submanifolds, but in general the subbundles themselves are only continuous.

When M is compact the notion is independent of the Riemannian metric. If M is not compact the notion is metric dependent; in fact, we will give an example of a metric on \mathbb{R}^3 with respect to which a coordinate field is Anosov, even though a coordinate field is clearly not Anosov with respect to the Euclidean metric on \mathbb{R}^3 . Since we are dealing with Riemannian metrics associated to a contact structure, when we speak of the *characteristic vector field being Anosov*, we will mean that it is Anosov with respect to an associated metric of the contact structure.

The following properties of Anosov flows will be of importance here. The subbundles $E^s \oplus \{\xi\}$ and $E^u \oplus \{\xi\}$ are integrable (Anosov [1967, Theorem 8]). Let μ denote the measure induced on M by the Riemannian metric. Recall that a flow is *ergodic* if for every measurable set S , $\psi_t(S) = S$ for all t implies $\mu(S)\mu(M - S) = 0$. If on a compact manifold an Anosov flow admits an integral invariant, i.e., an invariant measure that is equivalent to the measure μ , in particular if it is volume preserving, then it is ergodic (Anosov [1967, Theorem 4]), and in turn, by the ergodic theorem almost all orbits are dense (see e.g., Walters [1975, pp. 29–30]).

As an aside we note that on a compact manifold, an Anosov flow has a countable number of periodic orbits (Anosov [1967, Theorem 2]) and if the flow admits an integral invariant, then the set of periodic orbits is dense in M (Anosov [1967, Theorem 3]). This has an immediate implication for contact geometry. In Section 3.4 we discussed the conjecture of Weinstein that on a simply connected compact contact manifold, ξ must have a closed orbit, so in particular the Weinstein conjecture holds (without the simple connectivity) for a compact contact manifold on which ξ is Anosov.

Let us now turn our attention to the case of a 3-dimensional contact metric manifold and suppose that the ξ -sectional curvature is negative and that ξ is Anosov with respect to the associated metric. We then have both the special directions of Section 11.1 and the *Anosov directions*, i.e., the 1-dimensional stable and unstable bundles. One can then ask what happens if the special directions and the Anosov directions agree.

Theorem 11.2 *Let (M^3, η, g) be a 3-dimensional contact metric manifold with negative ξ -sectional curvature. If the characteristic vector field ξ generates an Anosov flow with respect to g and the special directions agree with the Anosov directions, then the contact metric structure*

satisfies $\nabla_\xi h = 0$. Moreover, if M^3 is compact, then it is a compact quotient of $\widetilde{SL}(2, \mathbb{R})$.

Proof. Suppose that Y is a local unit vector field such that $\nabla_Y \xi = \alpha Y$, $\alpha = -\sqrt{\lambda^2 - 1}$. Since ξ is Anosov and \mathcal{Y} agrees with the stable Anosov subbundle, the subbundle $\mathcal{Y} \oplus \{\xi\}$ is integrable. Thus from $[\xi, Y] = \nabla_\xi Y - \alpha Y$, $\nabla_\xi Y$ belongs to $\mathcal{Y} \oplus \{\xi\}$; but $g(\nabla_\xi Y, \xi) = 0$ and Y is unit, so $g(\nabla_\xi Y, Y) = 0$. Thus $\nabla_\xi Y = 0$. Similarly $\nabla_\xi Z = 0$. Recall the operator l defined by $lX = R_{X\xi}\xi$ for any X ; clearly l is a symmetric operator. Computing $R_{Y\xi}\xi$ and $R_{Z\xi}\xi$ we have

$$lY = -(\xi\alpha + \alpha^2)Y, \quad lZ = (\xi\alpha - \alpha^2)Z;$$

but Y and Z are not orthogonal, so $\xi\alpha = 0$ and $l|_{\mathcal{D}} = -\alpha^2 I|_{\mathcal{D}}$. Now compute $\nabla_\xi h$ acting on each vector of the h -eigenvector basis, $\{X, \phi X, \xi\}$, using the first equation of Proposition 7.1; this gives

$$(\nabla_\xi h)X = \phi(X - h^2 X - lX) = \phi(X - \lambda^2 X + \alpha^2 X) = 0$$

and similarly $(\nabla_\xi h)\phi X = 0$; $(\nabla_\xi h)\xi = 0$ is immediate. Therefore $\nabla_\xi h = 0$.

Finally, recall a result of E. Ghys [1987] that if ξ is Anosov on a compact 3-dimensional contact manifold M and the Anosov directions are smooth, then M is a compact quotient of $\widetilde{SL}(2, \mathbb{R})$. This may also be proved directly using consequences of $\nabla_\xi h = 0$; see the author's paper [1998]. ■

We now exhibit a family of contact metric structures on the Lie group $SL(2, \mathbb{R})$, show that the characteristic vector field is Anosov, and show that the special directions agree with the Anosov directions.

On a 3-dimensional unimodular Lie group we have a Lie algebra structure of the form

$$[e_2, e_3] = c_1 e_1, \quad [e_3, e_1] = c_2 e_2, \quad [e_1, e_2] = c_3 e_3.$$

In [1976] J. Milnor gave a complete classification of 3-dimensional Lie groups and their left-invariant metrics. If one c_i is nonzero, the dual 1-form ω_i is a contact form and e_i is the characteristic vector field. However, for the Riemannian metric defined by $g(e_i, e_j) = \delta_{ij}$ at the identity and extended by left translation to be an associated metric for ω_i , we must have $c_i = 2$. For $SL(2, \mathbb{R})$ two of the c_i 's are positive and one

negative in the Milnor classification, so taking ω_1 as the contact form, we write the Lie algebra as

$$[e_2, e_3] = 2e_1, \quad [e_3, e_1] = (1 - \lambda)e_2, \quad [e_1, e_2] = (1 + \lambda)e_3, \quad (*)$$

where $\lambda > 1$. Further, by way of notation, set

$$SL(2, \mathbb{R}) = \left\{ \begin{pmatrix} x & y \\ u & v \end{pmatrix} \mid xv - yu = 1 \right\}.$$

Now consider the matrices

$$\begin{pmatrix} \frac{1}{2}\sqrt{\lambda^2 - 1} & 0 \\ 0 & -\frac{1}{2}\sqrt{\lambda^2 - 1} \end{pmatrix}, \quad \begin{pmatrix} 0 & -\sqrt{\frac{\lambda+1}{2}} \\ \sqrt{\frac{\lambda+1}{2}} & 0 \end{pmatrix}, \quad \begin{pmatrix} 0 & -\sqrt{\frac{\lambda-1}{2}} \\ -\sqrt{\frac{\lambda-1}{2}} & 0 \end{pmatrix}$$

in the Lie algebra $\mathfrak{sl}(2, \mathbb{R})$, which we regard as the tangent space of $SL(2, \mathbb{R})$ at the identity. Applying the differential of left translation by $\begin{pmatrix} x & y \\ u & v \end{pmatrix}$ to these matrices gives the vector fields

$$\zeta_1 = \frac{1}{2}\sqrt{\lambda^2 - 1} \left(x \frac{\partial}{\partial x} - y \frac{\partial}{\partial y} + u \frac{\partial}{\partial u} - v \frac{\partial}{\partial v} \right),$$

$$\zeta_2 = \sqrt{\frac{\lambda+1}{2}} \left(y \frac{\partial}{\partial x} - x \frac{\partial}{\partial y} + v \frac{\partial}{\partial u} - u \frac{\partial}{\partial v} \right),$$

$$\zeta_3 = -\sqrt{\frac{\lambda-1}{2}} \left(y \frac{\partial}{\partial x} + x \frac{\partial}{\partial y} + v \frac{\partial}{\partial u} + u \frac{\partial}{\partial v} \right),$$

whose Lie brackets satisfy (*). Using these matrices again, define a left-invariant metric g ; then $\{\zeta_1, \zeta_2, \zeta_3\}$ is an orthonormal basis. The contact form ω_1 is given by

$$\omega_1 = \frac{2}{\sqrt{\lambda^2 - 1}}(v \, dx - y \, du).$$

The characteristic vector field ξ is ζ_1 . The metric g is an associated metric and ϕ as a skew-symmetric operator is given by $\phi\xi = 0$ and $\phi\zeta_2 = \zeta_3$. The symmetric operator h is given by $h\xi = 0$, $h\zeta_2 = \lambda\zeta_2$, $h\zeta_3 = -\lambda\zeta_3$. The special directions are

$$Y = \sqrt{\frac{\lambda-1}{2\lambda}}\zeta_2 + \sqrt{\frac{\lambda+1}{2\lambda}}\zeta_3 = -\frac{\sqrt{\lambda^2-1}}{\sqrt{\lambda}} \left(x \frac{\partial}{\partial y} + u \frac{\partial}{\partial v} \right)$$

and

$$Z = \sqrt{\frac{\lambda-1}{2\lambda}}\zeta_2 - \sqrt{\frac{\lambda+1}{2\lambda}}\zeta_3 = \frac{\sqrt{\lambda^2-1}}{\sqrt{\lambda}} \left(y \frac{\partial}{\partial x} + v \frac{\partial}{\partial u} \right).$$

The 1-parameter group of $\{\psi_t\}$ of ξ is given by

$$\psi_t \begin{pmatrix} x & y \\ u & v \end{pmatrix} = \begin{pmatrix} xe^{\frac{1}{2}\sqrt{\lambda^2-1}t} & ye^{-\frac{1}{2}\sqrt{\lambda^2-1}t} \\ ue^{\frac{1}{2}\sqrt{\lambda^2-1}t} & ve^{-\frac{1}{2}\sqrt{\lambda^2-1}t} \end{pmatrix}.$$

Then applying ψ_{t*} to Y at a point p yields

$$\begin{aligned} \psi_{t*}Y_p &= e^{-\frac{1}{2}\sqrt{\lambda^2-1}t}Y_p = e^{-\sqrt{\lambda^2-1}t}Y_{\psi_t(p)}, \\ \psi_{t*}Z_p &= e^{\frac{1}{2}\sqrt{\lambda^2-1}t}Z_p = e^{\sqrt{\lambda^2-1}t}Z_{\psi_t(p)}. \end{aligned}$$

Thus the corresponding subbundles \mathcal{Y} and \mathcal{Z} are invariant under the flow. Finally, since $\{\zeta_1, \zeta_2, \zeta_3\}$ is orthonormal, $|Y|^2 = \frac{\lambda-1}{2\lambda} + \frac{\lambda+1}{2\lambda} = 1$ and hence

$$|\psi_{t*}Y_p| = e^{-\frac{1}{2}\sqrt{\lambda^2-1}t}|Y_p|;$$

similarly

$$|\psi_{t*}Z_p| = e^{\frac{1}{2}\sqrt{\lambda^2-1}t}|Z_p|.$$

Thus ξ is an Anosov vector field and the special directions Y and Z agree with the Anosov directions.

In contrast to this, consider the contact metric structure on the tangent sphere bundle. We mentioned at the beginning of this section that the tangent sphere bundle of a surface is closely related to the structure on $SL(2, \mathbb{R})$ from both the topological and Anosov points of view. Comparing the above with the following theorem, we see that from the Riemannian point of view these are quite different.

Theorem 11.3 *With respect to the standard contact metric structure on the tangent sphere bundle of a negatively curved surface, the characteristic vector field is Anosov, but the special directions never agree with the stable and unstable directions.*

Proof. We noted in Section 9.2 that for the standard contact metric structure on the tangent sphere bundle of a Riemannian manifold, the characteristic vector field ξ is (twice) the geodesic flow, which is an Anosov vector field when the base manifold is negatively curved (see, e.g.,

Anosov [1967]). By Theorem 11.2, if the special directions of the contact metric structure agree with the Anosov directions, then $\nabla_\xi h = 0$. Now by Theorem 9.7 the standard contact metric structure of the tangent sphere bundle of any Riemannian manifold satisfies $\nabla_\xi h = 0$ if and only if the base manifold is of constant curvature 0 or +1. ■

Perrone [2000], utilizing the special directions Y and Z belonging to the contact subbundle on a contact metric manifold of negative ξ -sectional curvature, introduced another notion. Let \mathcal{Y} and \mathcal{Z} denote the subbundles generated by Y and Z . The special directions are said to be *Anosov-like* if the subbundles $\mathcal{Y} \oplus \{\xi\}$ and $\mathcal{Z} \oplus \{\xi\}$ are integrable. Perrone then proved that a contact metric 3-manifold admits Anosov-like special directions if and only if it satisfies $\nabla_\xi h = 0$ and has negative Ricci curvature in the direction ξ .

Three-dimensional contact metric manifolds satisfying $\nabla_\xi h = 0$ were called *3- τ -manifolds* by F. Gouli-Andreou and Ph. Xenos [1998]. The name comes from the equivalent condition $\nabla_\xi \tau = 0$, where here $\tau = \mathcal{L}_\xi g$; in particular, τ and h are related by $\tau(X, Y) = 2g(h\phi X, Y)$. A 3-dimensional contact metric manifold on which the Ricci operator Q and ϕ commute satisfies $\nabla_\xi h = 0$ but not conversely.

We close this section with an example (the author [1996]) of a contact metric manifold satisfying $\nabla_\xi h = 0$ but with $Q\phi \neq \phi Q$. We include this example here because it is also an example of a metric on \mathbb{R}^3 with respect to which the coordinate field $\frac{\partial}{\partial z}$ is Anosov.

Consider the standard Darboux contact form $\eta = \frac{1}{2}(dz - y dx)$ and characteristic vector field $\xi = 2\frac{\partial}{\partial z}$. Let f be a smooth function of x and y bounded below by a positive constant c . Then the metric given by

$$g = \frac{1}{4} \begin{pmatrix} \frac{e^{zf} + (1+f^2)e^{-zf} - 2}{f^2} + y^2 & \frac{e^{zf} - 1}{f} & -y \\ \frac{e^{zf} - 1}{f} & e^{zf} & 0 \\ -y & 0 & 1 \end{pmatrix}$$

is an associated metric. The tensor fields ϕ and h are given by

$$\phi = \begin{pmatrix} \frac{e^{zf} - 1}{f} & e^{zf} & 0 \\ -(\frac{e^{zf} + (1+f^2)e^{-zf} - 2}{f^2}) & -(\frac{e^{zf} - 1}{f}) & 0 \\ y(\frac{e^{zf} - 1}{f}) & ye^{zf} & 0 \end{pmatrix},$$

$$h = \begin{pmatrix} e^{zf} & fe^{zf} & 0 \\ -(\frac{fe^{zf} + (1+f^2)(-f)e^{-zf}}{f^2}) & -e^{zf} & 0 \\ ye^{zf} & yfe^{zf} & 0 \end{pmatrix}.$$

By direct computation, $\nabla_\xi h = 0$. Also $2\lambda^2 = \text{tr}h^2 = 2(1 + f^2)$, and hence the positive eigenfunction of h is $\lambda = \sqrt{1 + f^2} > 1$. Now on a 3-dimensional contact metric manifold satisfying $Q\phi = \phi Q$, the eigenfunction λ is a constant (Koufogiorgos, Sharma, and the author [1990]). Thus if f is not constant this structure on \mathbb{R}^3 satisfies $\nabla_\xi h = 0$ but not $Q\phi = \phi Q$.

For this structure the special directions discussed in Section 11.1 are given by

$$Y = f \frac{\partial}{\partial x} - \frac{\partial}{\partial y} + yf \frac{\partial}{\partial z}, \quad Z = \frac{\partial}{\partial y}.$$

To check that $\xi = 2\frac{\partial}{\partial z}$ is Anosov with respect to g , consider for simplicity just $\frac{\partial}{\partial z}$; its flow ψ_t maps a point $p_0(x, y, z)$ to the point $p(x, y, z + t)$. Now recalling that the function f was chosen to be bounded below by a positive constant c , we have for $t \leq 0$,

$$\left| \psi_{t*} \frac{\partial}{\partial y}(p_0) \right| = \left| \frac{\partial}{\partial y}(p) \right| = \frac{1}{2} e^{\frac{(z+t)f}{2}} = e^{\frac{tf}{2}} \left| \frac{\partial}{\partial y}(p_0) \right| \leq e^{\frac{ct}{2}} \left| \frac{\partial}{\partial y}(p_0) \right|.$$

Similarly for $t \geq 0$,

$$\left| \psi_{t*} \left(f \frac{\partial}{\partial x} - \frac{\partial}{\partial y} + yf \frac{\partial}{\partial z} \right)(p_0) \right| \leq e^{\frac{-ct}{2}} \left| \left(f \frac{\partial}{\partial x} - \frac{\partial}{\partial y} + yf \frac{\partial}{\partial z} \right)(p_0) \right|.$$

Thus $\frac{\partial}{\partial z}$, equivalently ξ , is Anosov with respect to this metric; Y determines the stable subbundle and Z the unstable subbundle.

11.3 Conformally Anosov flows

Another interesting notion, more general than an Anosov flow, is that of a conformally Anosov flow. We will show that if a 3-dimensional compact contact metric manifold has negative ξ -sectional curvature, then the characteristic vector field is conformally Anosov.

Mitsumatsu [1995] and Eliashberg and Thurston [1998] introduced a generalization of Anosov flows as follows. A flow ψ_t and its corresponding vector field are said to be *conformally Anosov*, Eliashberg and Thurston

[1998] (*projectively Anosov* Mitsumatsu [1995]), if there is a continuous Riemannian metric and a continuous, invariant splitting $TM = E^s \oplus E^u \oplus \{\xi\}$ as in the Anosov case such that for $Z \in E^u$ and $Y \in E^s$,

$$\frac{|\psi_{t*}Z|}{|\psi_{t*}Y|} \geq e^{ct} \frac{|Z|}{|Y|}$$

for some constant $c > 0$ and all $t \geq 0$.

Now a contact structure η on a 3-dimensional contact manifold M^3 determines an orientation on M^3 . This is true in dimension 3 even for a contact structure in the wider sense, since the sign of $\eta \wedge d\eta$ is independent of the choice of local contact form η .

The main result for our purpose from Mitsumatsu [1995, p. 1418] and Eliashberg and Thurston [1998, pp. 26–27] is the following.

Theorem 11.4 *If two contact structures (in the wider sense) on a compact 3-dimensional contact manifold M^3 induce opposite orientations, then the vector field directing the intersection of the two contact subbundles is a conformally Anosov flow. Conversely, given a conformally Anosov flow on M^3 , there exist two contact structures giving opposite orientations on M^3 whose contact subbundles intersect tangent to the flow.*

We now show that certain curvature hypotheses on a compact contact metric 3-manifold imply that the characteristic vector field ξ is conformally Anosov; in particular, negative ξ -sectional curvature is such a hypothesis (the author [2000]). A variation of this result appears in the author's paper with D. Perrone [1998].

Theorem 11.5 *Let $(M^3, \phi, \xi, \eta, g)$ be a compact 3-dimensional contact metric manifold with nowhere vanishing h and $\{e_1, e_2 (= \phi e_1), \xi\}$ an orthonormal eigenvector basis of h with $he_1 = \lambda e_1$ and λ the positive eigenvalue. If $K(\xi, e_1) < (1+\lambda)^2$ and $K(\xi, e_2) < (1-\lambda)^2$, then ξ is conformally Anosov. In particular, if the ξ -sectional curvature is negative, ξ is conformally Anosov.*

Proof. Let ω^1, ω^2 be the dual 1-forms of e_1 and e_2 . Since the three eigenvalues of h are everywhere distinct, the corresponding line fields are global. By the orientability, one may choose local bases directing the line fields that either agree or have two directions reversed in the overlap of coordinate neighborhoods. Thus in computing $\omega^1 \wedge d\omega^1$ and $\omega^2 \wedge d\omega^2$ on

such a basis we may regard the computations as global. By straightforward computation we have

$$\begin{aligned}\omega^1 \wedge d\omega^1(e_1, e_2, \xi) &= \frac{\lambda - 1}{6} - \frac{1}{6}g(\nabla_\xi e_1, e_2), \\ \omega^2 \wedge d\omega^2(e_1, e_2, \xi) &= -\frac{\lambda + 1}{6} - \frac{1}{6}g(\nabla_\xi e_1, e_2).\end{aligned}$$

On the other hand, applying the first equation of Proposition 7.1 to e_1 and e_2 , we obtain

$$\begin{aligned}K(\xi, e_1) &= 1 - \lambda^2 - 2\lambda g(\nabla_\xi e_1, e_2), \\ K(\xi, e_2) &= 1 - \lambda^2 + 2\lambda g(\nabla_\xi e_1, e_2).\end{aligned}$$

Combining these equations, we have

$$\omega^1 \wedge d\omega^1(e_1, e_2, \xi) = \frac{1}{12\lambda}((\lambda - 1)^2 - K(\xi, e_2))$$

and

$$\omega^2 \wedge d\omega^2(e_1, e_2, \xi) = \frac{1}{12\lambda}(-(\lambda + 1)^2 + K(\xi, e_1)).$$

The hypotheses now imply that $\omega^1 \wedge d\omega^1(e_1, e_2, \xi) > 0$ and $\omega^2 \wedge d\omega^2(e_1, e_2, \xi) < 0$. Therefore ξ is conformally Anosov by the result of Mitsumatsu and Eliashberg–Thurston. ■

We remark that the curvature of the standard contact metric structure on the tangent sphere bundle of a surface of negative curvature satisfies the curvature hypotheses in the first statement of the theorem. In particular, the standard contact metric structure on the tangent sphere bundle of a surface of constant curvature -1 has sectional curvature -7 for horizontal plane sections and sectional curvature $+1$ for plane sections spanned by ξ and the vertical direction; the positive eigenvalue λ is $+2$ with vertical vectors as eigenvectors (see Section 9.2).

We also remark that our argument uses two contact structures to study a third, an interesting idea in view of the result of Gonzalo [1987] (Section 3.2) that a 3-dimensional compact orientable manifold admits three independent contact structures.

It is immediate that the characteristic vector field ξ of a contact structure can never be Anosov or conformally Anosov with respect to a Sasakian metric. This is a consequence of the fact that on a Sasakian

manifold ξ is a Killing vector field; thus its flow is metric preserving and therefore cannot satisfy the exponential growth behavior in the definition of a classical or conformal Anosov flow. It is possible, however, for a vector field to belong to the contact subbundle of a Sasakian structure and be conformally Anosov with respect to this metric. We present an example of this which has the additional feature that the characteristic vector field is invariant along the flow (again see the author and D. Perrone [1998]). We begin with the following lemma.

Lemma 11.1 *On a 3-dimensional contact manifold, in terms of local Darboux coordinates (x, y, z) ($\eta = \frac{1}{2}(dz - y dx)$), any associated metric is of the form*

$$g = \frac{1}{4} \begin{pmatrix} a & b & -y \\ b & c & 0 \\ -y & 0 & 1 \end{pmatrix}$$

with $ac - b^2 - cy^2 = 1$; the metric is Sasakian if and only if the functions a , b and c are independent of z .

Proof. The form of the last row and column follows from the requirement that $\eta(X) = g(X, \xi)$. In dimension 3 the remaining requirements reduce to the determinant of the matrix (without the $\frac{1}{4}$) being 1. Also in dimension 3 the Sasakian condition is equivalent to the contact metric structure being K-contact. Thus evaluating the Lie derivative, $\mathcal{L}_\xi g$, on the coordinate vector fields, we see that $\xi = 2\frac{\partial}{\partial z}$ is Killing if and only if the functions a , b and c are independent of z . ■

To construct the example, consider $\mathbf{R}_+^3 = \{(x, y, z) | y > 0\}$ with the standard Darboux contact form $\eta = \frac{1}{2}(dz - y dx)$. The characteristic vector field is $2\frac{\partial}{\partial z}$ and the Riemannian metric given by the following matrix is Sasakian by the lemma:

$$g = \frac{1}{4} \begin{pmatrix} e^{2y} & \sqrt{e^{2y} - y^2 - 1} & -y \\ \sqrt{e^{2y} - y^2 - 1} & 1 & 0 \\ -y & 0 & 1 \end{pmatrix}.$$

The vector field $\frac{\partial}{\partial y}$ is conformally Anosov with respect to this metric. To see this we observe that the subbundles determined by $\frac{\partial}{\partial z} = \frac{1}{2}\xi$ and $\frac{\partial}{\partial x}$ correspond to E^s and E^u respectively. The flow simply maps a point

$p_0(x, y, z)$ to the point $p(x, y + t, z)$, and we have easily that

$$\frac{|\frac{\partial}{\partial x}(p)|}{|\frac{\partial}{\partial z}(p)|} = e^t \frac{|\frac{\partial}{\partial x}(p_0)|}{|\frac{\partial}{\partial z}(p_0)|}.$$

Clearly this flow satisfies $\psi_{t*}\xi = \xi$.

For a discussion of conformally Anosov flows on 3-dimensional homogeneous contact metric manifolds including an Anosov flow belonging to the contact subbundle on the Lie group $E(1, 1)$, see Perrone and the author [1998].

12

Complex Contact Manifolds

While the study of complex contact manifolds is almost as old as the modern theory of real contact manifolds, the subject has received much less attention, and since many examples are now appearing in the literature, we devote this and the next chapter to the subject.

12.1 Complex contact manifolds and associated metrics

The notion of a complex contact manifold stems from the late 1950s and early 1960s with the papers of Kobayashi [1959] and Boothby [1961], [1962]; this is just shortly after the Boothby–Wang fibration in real contact geometry. Then in [1965], J. A. Wolf studied homogeneous complex contact manifolds and their relation to quaternionic symmetric spaces. An example of more recent work is the result of Moroianu and Semmelmann [1994] that on a compact spin Kähler manifold M of positive scalar curvature and complex dimension $4l + 3$, the following are equivalent: (i) M is a Kähler–Einstein manifold admitting a complex contact structure, (ii) M is the twistor space of a quaternionic Kähler manifold of positive scalar curvature, (iii) M admits Kählerian Killing spinors. LeBrun [1995] proves that a complex contact manifold of positive first Chern class, i.e., a

Fano contact manifold, is a twistor space if and only if it admits a Kähler Einstein metric and conjectures that every Fano contact manifold is a twistor space.

In the 1970s and early 1980s there was a development, of the Riemannian theory of complex contact manifolds by Ishihara and Konishi [1979], [1980], [1982]. In this development, however, the notion of normality seems too strong, since it precludes the complex Heisenberg group as one of the canonical examples, although it does include complex projective spaces of odd complex dimension as one would expect (see Section 12.4). In the real case both the Heisenberg group and the odd-dimensional spheres have natural Sasakian (normal contact metric) structures. As a subject, the Riemannian geometry of complex contact manifolds is still in its infancy.

A *complex contact manifold* is a complex manifold of odd complex dimension $2n + 1$ together with an open covering $\{\mathcal{O}_\alpha\}$ by coordinate neighborhoods such that:

1. On each \mathcal{O}_α there is a holomorphic 1-form θ_α such that

$$\theta_\alpha \wedge (d\theta_\alpha)^n \neq 0.$$

2. On $\mathcal{O}_\alpha \cap \mathcal{O}_\beta \neq \emptyset$ there is a nonvanishing holomorphic function $f_{\alpha\beta}$ such that $\theta_\alpha = f_{\alpha\beta}\theta_\beta$.

The subspaces $\{X \in T_m \mathcal{O}_\alpha : \theta_\alpha(X) = 0\}$ define a nonintegrable holomorphic subbundle \mathcal{H} of complex dimension $2n$ called the *complex contact subbundle* or *horizontal subbundle*. The quotient $L = TM/\mathcal{H}$ is a complex line bundle over M . Kobayashi [1959] proved that $c_1(M) = (n + 1)c_1(L)$ and hence for a compact complex contact manifold, a complex contact structure is given by a global 1-form if and only if its first Chern class vanishes (see also Boothby [1961], [1962]). It is for this reason that our definition of complex contact structure is analogous to that of a contact structure in the wider sense. Even for the most canonical example of a complex contact manifold, $\mathbb{C}P^{2n+1}$, the structure is not given by a global form. Since a holomorphic p -form on a compact Kähler manifold is closed (see, e.g., Goldberg [1962, p. 177]), no compact Kähler manifold has a complex contact structure given by a global contact form. Moreover, Ye [1994] showed that a compact Kähler manifold with vanishing first Chern class has no complex contact structure. There are, however,

interesting examples of complex contact manifolds with global complex contact forms, as we shall see; these are called *strict complex contact manifolds* (Foreman [2000a], [2000b]).

We will not need a complex Darboux theorem in our development here, but a complex version of the real Darboux theorem is possible, and such a result is discussed briefly by LeBrun [1995, p. 423].

If $(M, \{\theta_\alpha\})$ is a complex contact manifold, the transition functions $f_{\alpha\beta}$ define a holomorphic line bundle over M , viz. L^{-1} . Using local sections of this bundle we define complex-valued 1-forms $\{\pi_\alpha\}$ such that each π_α is a nonvanishing, complex-valued function multiple of θ_α and on $\mathcal{O}_\alpha \cap \mathcal{O}_\beta \neq \emptyset$,

$$\pi_\alpha = h_{\alpha\beta} \pi_\beta, \quad h_{\alpha\beta} : \mathcal{O}_\alpha \cap \mathcal{O}_\beta \longrightarrow S^1.$$

The $h_{\alpha\beta}$ are then the transition functions of a circle bundle P over M (see Ishihara and Konishi [1982]) and on \mathcal{O}_α ,

$$\pi_\alpha \wedge (d\pi_\alpha)^n \wedge \bar{\pi}_\alpha \wedge (d\bar{\pi}_\alpha)^n \neq 0.$$

Writing $\pi_\alpha = u_\alpha - iv_\alpha$, we have $v_\alpha = u_\alpha \circ J$, since θ_α is holomorphic. Moreover, u_α and v_α transform naturally with respect to S^1 ; namely, if $h_{\alpha\beta} = a + ib$, then

$$u_\beta = au_\alpha - bv_\alpha, \quad v_\beta = bu_\alpha + av_\alpha, \quad a^2 + b^2 = 1.$$

The set $\{\pi_\alpha\}$ is called a *normalized complex contact structure* with respect to $\{\theta_\alpha\}$; Foreman [1996].

For simplicity we will often omit the subscripts on the local tensor fields. Define a local section U of TM , i.e., a section of $T\mathcal{O}$, by $du(U, X) = 0$ for every $X \in \mathcal{H}$, $u(U) = 1$ and $v(U) = 0$. Such local sections then define a global subbundle \mathcal{V} by $\mathcal{V}|_{\mathcal{O}} = \text{Span}\{U, JU\}$. We now have $TM = \mathcal{H} \oplus \mathcal{V}$, and we denote the projection map to \mathcal{H} by

$$p : TM \longrightarrow \mathcal{H}.$$

The subbundle \mathcal{V} is called the *vertical subbundle* or *characteristic subbundle*. It is generally assumed that \mathcal{V} is integrable, but in Subsection 13.4.1 we will give an example of a complex contact structure for which \mathcal{V} is not integrable. Except for this example the integrability will be assumed throughout.

On the other hand, if M is a complex manifold with almost complex structure J , Hermitian metric g and open covering by coordinate neighborhoods $\{\mathcal{O}_\alpha\}$, M is called a *complex almost contact metric manifold* if it satisfies the following two conditions:

(1) On each \mathcal{O}_α there exist 1-forms u_α and $v_\alpha = u_\alpha \circ J$ with orthogonal dual vector fields U_α and $V_\alpha = -JU_\alpha$ and $(1, 1)$ tensor fields G_α and $H_\alpha = G_\alpha J$ such that

$$\begin{aligned} G_\alpha^2 &= H_\alpha^2 = -I + u_\alpha \otimes U_\alpha + v_\alpha \otimes V_\alpha, \\ G_\alpha J &= -JG_\alpha, \quad G_\alpha U = 0, \quad g(X, G_\alpha Y) = -g(G_\alpha X, Y). \end{aligned}$$

(2) On $\mathcal{O}_\alpha \cap \mathcal{O}_\beta \neq \emptyset$,

$$\begin{aligned} u_\beta &= au_\alpha - bv_\alpha, \quad v_\beta = bu_\alpha + av_\alpha, \\ G_\beta &= aG_\alpha - bH_\alpha, \quad H_\beta = bG_\alpha + aH_\alpha, \end{aligned}$$

where a and b are functions with $a^2 + b^2 = 1$.

It is clear that H_α also anticommutes with J and is skew-symmetric with respect to g and that G_α and H_α annihilate both U and V .

Returning to the local forms $\pi_\alpha = u_\alpha - iv_\alpha$ on a complex contact manifold, a Hermitian metric g is called an *associated metric* if there are local fields of endomorphisms G_α such that the tensor fields G_α , $H_\alpha = G_\alpha J$, U_α , $V_\alpha = -JU_\alpha$, u_α , $v_\alpha = u_\alpha \circ J$, g form a complex almost contact metric structure satisfying

$$g(X, G_\alpha Y) = du_\alpha(X, Y), \quad g(X, H_\alpha Y) = dv_\alpha(X, Y), \quad X, Y \in \mathcal{H}.$$

As a consequence we also have $U_\beta = aU_\alpha - bV_\alpha$, $V_\beta = bU_\alpha + aV_\alpha$ and hence $U_\alpha + iV_\alpha = h_{\alpha\beta}^{-1}(U_\beta + iV_\beta)$. In particular, $\{h_{\alpha\beta}^{-1}\}$ are the transition functions of the bundle \mathcal{V} . Another way to look at this subbundle is to note that on $\mathcal{O}_\alpha \cap \mathcal{O}_\beta$, $U_\alpha \wedge V_\alpha = U_\beta \wedge V_\beta$ and hence that each $U_\alpha \wedge V_\alpha$ defines part of a global object.

Kobayashi [1959] also showed that the structural group of a complex contact manifold is reducible to $(Sp(n) \cdot U(1)) \times U(1)$. Such a reduction is equivalent to a complex almost contact metric structure; cf. Shibuya [1978]. Shibuya's approach is the complex analogue of that of Hatakeyama [1962], which we utilized in Section 4.2. A formulation in terms of global tensor fields similar to the fundamental 4-form of a quaternionic Kähler manifold (see, e.g., Ishihara [1974]) was given by Ishihara, Ludden and the author [1978].

Ishihara and Konishi [1982] (see also Foreman [1996]) prove that a complex contact manifold admits a complex almost contact metric structure for which the local contact form θ is of the form $u - iv$ to within a non-vanishing complex-valued function multiple and the local tensor fields G and H are related to du and dv by

$$\begin{aligned} du(X, Y) &= g(X, GY) + (\sigma \wedge v)(X, Y), \\ dv(X, Y) &= g(X, HY) - (\sigma \wedge u)(X, Y) \end{aligned}$$

for some 1-form σ . When \mathcal{V} is integrable, σ takes the form $\sigma(X) = g(\nabla_X U, V)$. We refer to a complex contact manifold with a complex almost contact metric structure satisfying these conditions as a *complex contact metric manifold*. For a given normalized contact structure we denote by \mathcal{A} , as in the real case, the space of associated metrics. As a matter of notation we define local 2-forms \hat{G} and \hat{H} by $\hat{G}(X, Y) = g(X, GY)$ and $\hat{H}(X, Y) = g(X, HY)$.

In the case of a strict complex contact structure, u and v may be taken globally such that $\theta = u - iv$ and $\sigma = 0$.

Bearing in mind that the local contact form θ is $u - iv$ to within a nonvanishing complex-valued function multiple and since in the overlap $\mathcal{O}_\alpha \cap \mathcal{O}_\beta$, $u_\beta = au_\alpha - bv_\alpha$, $v_\beta = bu_\alpha + av_\alpha$, we can define an *integral submanifold* as a submanifold whose tangent spaces belong to the complex contact subbundle, i.e., $u(X) = v(X) = 0$ or equivalently $\theta(X) = 0$. If the submanifold is itself a complex submanifold, we call it a *holomorphic integral submanifold*. When the holomorphic integral submanifold has complex dimension 1, it is called a *holomorphic Legendre curve*. Recall that in real contact geometry the maximum dimension of an integral submanifold of a contact manifold of dimension $2n + 1$ is only n . Similarly, since U and V are normal, from the above equations we see that for any integral submanifold, GX and HX are normal for any tangent vector X . Thus an integral submanifold of a complex contact manifold of complex dimension $2n + 1$ has real dimension at most $2n$.

We have seen in the earlier chapters that for a contact metric structure (ϕ, ξ, η, g) , the tensor field h defined by $h = \frac{1}{2}\mathcal{L}_\xi\phi$ plays a fundamental role. Now for a complex contact metric structure we define local tensor fields by

$$h_U = \frac{1}{2}\text{sym}(\mathcal{L}_U G) \circ p, \quad h_V = \frac{1}{2}\text{sym}(\mathcal{L}_V H) \circ p,$$

where sym denotes the symmetric part; h_U anticommutes with G , h_V anticommutes with H , and

$$\nabla_X U = -GX - Gh_U X + \sigma(X)V, \quad \nabla_X V = -HX - Hh_V X - \sigma(X)U.$$

From these equations one readily sees that the integral surfaces of \mathcal{V} are totally geodesic submanifolds. Just as the meaning of the vanishing of h in the real case is that the metric was invariant under the action of ξ , i.e., ξ is Killing, an associated metric g is projectable with respect to the foliation induced by the integrable subbundle \mathcal{V} if and only if h_U and h_V vanish.

12.2 Examples of complex contact manifolds

12.2.1 Complex Heisenberg group

We have seen that \mathbb{R}^3 with the Darboux form $\eta = \frac{1}{2}(dz - y dx)$ as its contact structure and Sasakian metric $g = \frac{1}{4}(dx^2 + dy^2) + \eta \otimes \eta$ is a standard example of a contact metric manifold. Identifying \mathbb{R}^3 with the Heisenberg group

$$H_{\mathbb{R}} = \left\{ \left(\begin{pmatrix} 1 & y & z \\ 0 & 1 & x \\ 0 & 0 & 1 \end{pmatrix} \middle| x, y, z \in \mathbb{R} \right\} \simeq \mathbb{R}^3,$$

left translation preserves η , and g is a left-invariant metric on $H_{\mathbb{R}}$ (see Example 4.5.1).

The complex Heisenberg group is the closed subgroup $H_{\mathbb{C}}$ of $GL(3, \mathbb{C})$ given by

$$H_{\mathbb{C}} = \left\{ \left(\begin{pmatrix} 1 & z_2 & z_3 \\ 0 & 1 & z_1 \\ 0 & 0 & 1 \end{pmatrix} \middle| z_1, z_2, z_3 \in \mathbb{C} \right\} \simeq \mathbb{C}^3.$$

If L_B denotes left translation by $B \in H_{\mathbb{C}}$, then $L_B^* dz_1 = dz_1$, $L_B^* dz_2 = dz_2$, $L_B^*(dz_3 - z_2 dz_1) = dz_3 - z_2 dz_1$. The vector fields $\frac{\partial}{\partial z_1} + z_2 \frac{\partial}{\partial z_3}$, $\frac{\partial}{\partial z_2}$, $\frac{\partial}{\partial z_3}$ are dual to the 1-forms dz_1 , dz_2 , $dz_3 - z_2 dz_1$ and are left-invariant vector fields. Moreover, relative to the coordinates $(z_1, z_2, z_3, \bar{z}_1, \bar{z}_2, \bar{z}_3)$

the Hermitian metric (Jayne [1992, p. 234])

$$g = \frac{1}{8} \left(\begin{array}{ccc|ccc} & & & 1 + |z_2|^2 & 0 & -z_2 \\ & & O & 0 & 1 & 0 \\ & & & -\bar{z}_2 & 0 & 1 \\ \hline 1 + |z_2|^2 & 0 & -\bar{z}_2 & & & \\ 0 & 1 & 0 & & O & \\ -z_2 & 0 & 1 & & & \end{array} \right)$$

is a left-invariant metric on $H_{\mathbb{C}}$, but it is not a Kähler metric. The form

$$\theta = \frac{1}{2}(dz_3 - z_2 dz_1)$$

is a complex contact structure on $H_{\mathbb{C}}$, and in our view, $(H_{\mathbb{C}}, \theta, g)$ plays the role in the geometry of complex contact manifolds that \mathbb{R}^3 with its standard Sasakian structure does in the geometry of real contact manifolds.

As we have seen, a complex contact manifold admits a complex almost contact metric structure. Here $H_{\mathbb{C}} \simeq \mathbb{C}^3$ and θ is global, so the structure tensors may be taken globally. With J denoting the standard almost complex structure on \mathbb{C}^3 , $J \frac{\partial}{\partial x_i} = \frac{\partial}{\partial y_i}$, we may give a complex almost contact metric structure to $H_{\mathbb{C}}$ as follows. Since θ is holomorphic, setting $\theta = u - iv$, we have $v = u \circ J$. Also set $4 \frac{\partial}{\partial z_3} = U + iV$; then $u(X) = g(U, X)$ and $v(X) = g(V, X)$. Finally, in complex coordinates G and H were given by Jayne [1992, p. 235]:

$$G = \left(\begin{array}{ccc|ccc} & & & 0 & 1 & 0 \\ & & O & -1 & 0 & 0 \\ & & & 0 & z_2 & 0 \\ \hline 0 & 1 & 0 & & & \\ -1 & 0 & 0 & & O & \\ 0 & \bar{z}_2 & 0 & & & \end{array} \right),$$

$$H = \left(\begin{array}{ccc|ccc} & & & 0 & -i & 0 \\ & & & i & 0 & 0 \\ & & & 0 & -iz_2 & 0 \\ \hline 0 & i & 0 & & & \\ -i & 0 & 0 & & & \\ 0 & i\bar{z}_2 & 0 & & & \end{array} \right).$$

Note that the matrix gG is that of the real part $d\theta$ and the matrix gH is that of the imaginary part of $d\theta$. For this structure the 1-form σ vanishes.

Now let

$$\Gamma = \left\{ \left(\begin{array}{ccc} 1 & \gamma_2 & \gamma_3 \\ 0 & 1 & \gamma_1 \\ 0 & 0 & 1 \end{array} \right) \mid \gamma_k = m_k + in_k, m_k, n_k \in \mathbb{Z} \right\};$$

Γ is a subgroup of $H_{\mathbb{C}} \simeq \mathbb{C}^3$. The 1-form $dz_3 - z_2 dz_1$ is invariant under the action of Γ and hence the quotient $H_{\mathbb{C}}/\Gamma$ is a compact complex contact manifold with a global complex contact form. $H_{\mathbb{C}}/\Gamma$ is known as the *Iwasawa manifold*. The Iwasawa manifold has no Kähler structure, but it does have an indefinite Kähler structure and it has symplectic forms; see Fernández and Gray [1986].

12.2.2 Odd-dimensional complex projective space

We will need a local expression for the complex contact structure on $\mathbb{C}P^{2n+1}$ and will use homogeneous coordinates $(z_1, \dots, z_{n+1}, w_1, \dots, w_{n+1})$. Then the complex contact structure is given by the holomorphic 1-form

$$\psi = \sum_{k=1}^{n+1} (z_k dw_k - w_k dz_k).$$

To give a little more detail we remark that the complex contact structure on $\mathbb{C}P^{2n+1}$ is closely related to the Sasakian 3-structure on the sphere S^{4n+3} (see Chapter 14) and to the quaternionic Kähler structure on quaternionic projective space $\mathbb{H}P^n$. The space $\mathbb{C}^{2n+2} \simeq \mathbb{H}^{n+1}$ has three almost complex structures I, J, K , which act on the position

vector \mathbf{x} as

$$\begin{aligned} I\mathbf{x} &= i\mathbf{x} = (iz_1, \dots, iz_{n+1}, iw_1, \dots, iw_{n+1}), \\ J\mathbf{x} &= (i\bar{w}_1, \dots, i\bar{w}_{n+1}, -i\bar{z}_1, \dots, -i\bar{z}_{n+1}), \\ K\mathbf{x} &= (\bar{w}_1, \dots, \bar{w}_{n+1}, -\bar{z}_1, \dots, -\bar{z}_{n+1}). \end{aligned}$$

The vector fields on S^{4n+3} given by $\xi_1 = -I\mathbf{x}$, $\xi_2 = -J\mathbf{x}$, $\xi_3 = -K\mathbf{x}$ are the characteristic vector fields of the three contact structures η_1, η_2, η_3 on S^{4n+3} . In terms of the complex coordinates, η_1, η_2, η_3 on S^{4n+3} are the restrictions of the following forms on \mathbb{C}^{2n+2} , which we denote by the same letters:

$$\begin{aligned} \eta_1 &= -\frac{i}{2} \sum_{k=1}^{n+1} (z_k d\bar{z}_k - \bar{z}_k dz_k + w_k d\bar{w}_k - \bar{w}_k dw_k), \\ \psi &= \eta_3 + i\eta_2 = \sum_{k=1}^{n+1} (z_k dw_k - w_k dz_k). \end{aligned}$$

Ishihara and Konishi [1979] proved that if one of the contact structures of a manifold \tilde{M}^{4n+3} with a Sasakian 3-structure (see Theorem 14.10) is regular, the base manifold M of the induced fibration is a complex contact manifold. The structure is constructed as follows. If $(\phi_1, \xi_1, \eta_1, g)$ is the regular Sasakian structure, then ϕ_1 and g are projectable. Let $\tilde{\pi}$ denote the horizontal lift with respect to the principal S^1 bundle connection defined by η_1 . Then J defined by $JX = \pi_* \phi_1 \tilde{\pi} X$ and the projected metric form a Kähler structure on M (cf. Example 6.7.2). For a coordinate neighborhood \mathcal{U} on M and a local cross section s of \tilde{M}^{4n+3} over \mathcal{U} , the 1-forms u and v and a tensor field G defined on \mathcal{U} by

$$\begin{aligned} u(X) \circ \pi &= \eta_2(s_* X), \quad v(X) \circ \pi = \eta_3(s_* X), \\ GX &= \pi_*(\phi_2 s_* X - \eta_1(s_* X)\xi_3 + \eta_3(s_* X)\xi_1), \end{aligned}$$

define the complex contact and complex almost contact structures on M . In the case of the Hopf fibration this is the standard Kähler structure on $\mathbb{C}P^{2n+1}$ with the Fubini–Study metric. With the Hopf fibration induced by ξ_1 , $\psi = \eta_3 + i\eta_2$ is a local expression for the complex contact structure on $\mathbb{C}P^{2n+1}$.

12.2.3 Twistor spaces

Generalizing the previous example, we discuss twistor spaces over quaternionic Kähler manifolds, the twistor space of quaternionic projective space being odd-dimensional complex projective space. Consider a Riemannian manifold (M^{4n}, g) whose holonomy group is contained in $Sp(n) \cdot Sp(1) = Sp(n) \times Sp(1)/\{\pm I\}$. This means that there exists a subbundle $E \subset \text{End}(TM)$ with 3-dimensional fibers such that locally there exists a basis of E consisting of almost complex structures $\{\mathcal{I}, \mathcal{J}, \mathcal{K}\}$ satisfying, $\mathcal{I}\mathcal{J} = -\mathcal{J}\mathcal{I} = \mathcal{K}$ and such that $\nabla_X \mathcal{I}, \nabla_X \mathcal{J}, \nabla_X \mathcal{K}$ belong to the span of $\{\mathcal{I}, \mathcal{J}, \mathcal{K}\}$ for any vector field X on M . In dimension 4 ($n = 1$), $Sp(1) \cdot Sp(1) = SO(4)$, so the holonomy group condition is not a restriction. For $n > 1$, (M^{4n}, g) is called a *quaternionic Kähler manifold*. A well-known result of Alekseevskii [1968] (see also Berger [1966], Ishihara [1974]) is that in this case, the metric g is Einstein.

Since $Sp(1) \cdot Sp(1) = SO(4)$, a stronger definition of quaternionic Kähler manifold is needed in dimension 4. A 4-dimensional manifold is said to be a *quaternionic Kähler manifold* if it is Einstein and self-dual with nonzero scalar curvature (see, e.g., LeBrun [1991]).

Returning to the subbundle $\bar{\pi} : E \longrightarrow M$, regardless of dimension, we define the *horizontal* subspace $\bar{\mathcal{H}}_p$ at $p \in E$ as follows. Let α denote a curve in M such that $\alpha(0) = \bar{\pi}(p)$ and let s denote a section of E such that $s(\alpha(0)) = p$. Then set

$$\bar{\mathcal{H}}_p = \{(s \circ \alpha)_*(0) \in T_p E \mid \nabla_{\dot{\alpha}(0)} s = 0\}.$$

Now induce a bundle metric on E by making each quaternionic Kähler frame $\{\mathcal{I}, \mathcal{J}, \mathcal{K}\}$ an orthonormal basis. Let Z be the space of all unit elements of E . Locally

$$Z = \{x\mathcal{I} + y\mathcal{J} + z\mathcal{K} \in E \mid x^2 + y^2 + z^2 = 1\};$$

$\pi : Z \longrightarrow M$, is called the *twistor space* of M and each element $j \in Z$ is an almost complex structure on the tangent space of M at $\pi(j)$.

Let $\mathcal{V} = \ker \pi_*$. Since each fiber of Z is a unit sphere, then at $j = x\mathcal{I} + y\mathcal{J} + z\mathcal{K}$, $x^2 + y^2 + z^2 = 1$, we may make the identification

$$\mathcal{V}_j = \{X \in E_{\pi(j)} \mid X \perp j\} = \{a\mathcal{I} + b\mathcal{J} + c\mathcal{K} \mid ax + by + cz = 0\}.$$

Setting $\mathcal{H} = \bar{\mathcal{H}}|_Z$, we have the splitting $TZ \cong \mathcal{V} \oplus \mathcal{H}$. Define an almost complex structure J on Z as follows. For a vertical vector $V \in \mathcal{V}_j$, set

$JV = j \times V$, where \times denotes the usual vector product in Euclidean 3-space; in particular, this is the usual almost complex structure on S^2 . To define the action of J on $X \in \mathcal{H}_j$, first let $\tilde{}$ denote the horizontal lift with respect to the connection on E determined by $\bar{\mathcal{H}}$. Then since j is an almost complex structure on the tangent space of M at $\pi(j)$, set $JX = \widetilde{j\pi_*X}$. Now local \mathcal{V} -valued forms θ defining \mathcal{H} give Z a complex contact structure; see Salamon [1982] or for verification of the integrability, Besse [1987, pp. 413–415] and for $\theta \wedge (d\theta)^n \neq 0$, Besse [1987, p. 416].

Now let $\{\mathcal{U}_\alpha\}$ be an open covering of the quaternionic Kähler manifold (M^{4n}, g) , and corresponding to a neighborhood \mathcal{U}_α set

$$\mathcal{O}_\alpha = \{x\mathcal{I} + y\mathcal{J} + z\mathcal{K} \in Z \mid z \neq 1\}, \quad \mathcal{O}'_\alpha = \{x\mathcal{I} + y\mathcal{J} + z\mathcal{K} \in Z \mid z \neq -1\}.$$

The collection of pairs $\{\mathcal{O}_\alpha, \mathcal{O}'_\alpha\}$ is then an atlas on Z . On \mathcal{O}_α define vector fields \hat{U} and \hat{V} by

$$\hat{U} = \frac{1 - z - x^2}{1 - z}\mathcal{I} - \frac{xy}{1 - z}\mathcal{J} + x\mathcal{K}$$

and $\hat{V} = -J\hat{U}$. Then $\{\hat{U}, \hat{V}\}$ forms a basis of \mathcal{V} on \mathcal{O}_α . As elements of E , \hat{U} and \hat{V} are unit and there exist real 1-forms \hat{u} and \hat{v} such that $\hat{u}(\hat{U}) = \hat{v}(\hat{V}) = 1$, $\hat{u}(\hat{V}) = \hat{v}(\hat{U}) = 0$. On \mathcal{O}'_α define vector fields \hat{U}' and \hat{V}' by

$$\hat{U}' = \frac{1 + z - x^2}{1 + z}\mathcal{I} - \frac{xy}{1 + z}\mathcal{J} + x\mathcal{K}$$

and $\hat{V}' = -J\hat{U}'$. Again we have the corresponding 1-forms \hat{u}' , \hat{v}' and

$$\hat{U} + i\hat{V} = h(\hat{U}' + i\hat{V}'), \quad \hat{u} - i\hat{v} = h(\hat{u}' - i\hat{v}'),$$

where $h : \mathcal{O}_\alpha \cap \mathcal{O}'_\alpha \longrightarrow S^1$. Thus $\{\hat{u} - i\hat{v}\}$ is a normalized contact structure on Z and $\hat{u} \otimes \hat{u} + \hat{v} \otimes \hat{v}$ is a global tensor field on Z .

Let $c > 0$ and define a Riemannian metric g_c on Z by

$$g_c = c^2(\hat{u} \otimes \hat{u} + \hat{v} \otimes \hat{v}) + \pi^*g,$$

$g_c|_{\mathcal{V}} = c^2\langle, \rangle$, where \langle, \rangle is the Euclidean metric on fibers of E restricted to S^2 . The metric g_c is called the *Salamon, Bérard-Bergery metric with vertical coefficient c* . Setting $u_c = c\hat{u}$, $v_c = c\hat{v}$, $\{u_c - iv_c\}$ is a normalized contact structure on Z and

$$g_c = u_c \otimes u_c + v_c \otimes v_c + \pi^*g.$$

In [1996] Foreman proved the following theorem.

Theorem 12.1 *Let (M^{4n}, g) be a quaternionic Kähler manifold and τ the scalar curvature of g . Consider the twistor space Z with the Salamon, Bérard-Bergery metric g_c , $c > 0$, and normalized contact structure $\{u_c - iv_c\}$. Then we have the following:*

1. g_c is Hermitian with respect to the complex structure J on Z .
2. g_c is an associated metric if and only if $c|\tau| = 8n(n+2)$.
3. g_c is Kähler if and only if $c^2\tau = 4n(n+2)$.
4. g_c is associated and Kähler if and only if $c = \frac{1}{2}$ and $\tau = 16n(n+2)$.

Recall that in Section 1 we noted conditions for an associated metric g to be projectable with respect to the foliation induced by the integrable subbundle \mathcal{V} , namely that h_U and h_V vanish. Foreman [1996] also proves the following result.

Theorem 12.2 *Let Z be the twistor space over a quaternionic Kähler manifold (M^{4n}, g) of constant scalar curvature τ and let c be such that $c|\tau| = 8n(n+2)$. Then in the space \mathcal{A} of all associated metrics with respect to the normalized contact structure $\{u_c - iv_c\}$, the Salamon, Bérard-Bergery metric g_c is the only projectable associated metric in \mathcal{A} .*

Again it seems worthwhile to mention the result of LeBrun [1995].

Theorem 12.3 *A complex contact manifold of positive first Chern class is a twistor space if and only if it admits a Kähler Einstein metric.*

Further discussion of the geometry of twistor space can be found in Foreman [2000b], [2002b]. In particular he gives curvature conditions for a complex contact metric manifold to be the twistor space of a quaternionic Kähler manifold. His main result is the following.

Theorem 12.4 *A complex contact metric manifold is isometric to the twistor space of a quaternionic Kähler manifold with positive scalar curvature if and only if*

$$\begin{aligned} R_{XY}(aU + bV) &= (au + bv)(Y)X - (au + bv)(X)Y \\ &\quad + (au + bv)(JY)JX - (au + bv)(JX)JY \\ &\quad + 2g(X, JY)(aU + bV) \end{aligned}$$

12.2.4 The Complex Boothby–Wang fibration

In [2000a] and again in [2000b] Foreman constructed complex contact manifolds with global complex contact forms and fibrations with vertical

fibers $S^1 \times S^1$, and hence these examples are quite different from the twistor space examples. Here we discuss the complex Boothby–Wang fibration of Foreman [2000a]. Let (M, Ω) be a *complex symplectic manifold* of complex dimension $2n$ and complex structure J_0 , i.e., M is a complex manifold together with a closed holomorphic 2-form Ω such that $\Omega^n \neq 0$. Writing $\Omega = \Omega_1 + i\Omega_2$, we see that Ω_1 and Ω_2 are closed 2-forms. On the other hand, Ω may be written in terms of a local basis of holomorphic 1-forms as $\theta_1 \wedge \theta_{n+1} + \cdots + \theta_n \wedge \theta_{2n}$. Then taking real and imaginary parts of $\theta_k = \alpha_k + i\beta_k$, we have

$$\begin{aligned}\Omega_1 &= \alpha_1 \wedge \alpha_{n+1} - \beta_1 \wedge \beta_{n+1} + \cdots + \alpha_n \wedge \alpha_{2n} - \beta_n \wedge \beta_{2n}, \\ \Omega_2 &= \alpha_1 \wedge \beta_{n+1} + \beta_1 \wedge \alpha_{n+1} + \cdots + \alpha_n \wedge \beta_{2n} - \beta_n \wedge \alpha_{2n},\end{aligned}$$

from which we see that $\Omega_1^{2n} \neq 0$ and $\Omega_2^{2n} \neq 0$. Thus we have two distinct symplectic structures on M . If each of these is of integral class, we have two principal circle bundles P_1 and P_2 with contact (connection) forms η_1 and η_2 as in Sections 3.3, 4.5.4 and 6.7.2, and each $d\eta_k = \Omega_k$. Finally, let ξ_1 and ξ_2 be the characteristic vector fields of these contact structures.

Define a principal $S^1 \times S^1$ -bundle P over M by $P = P_1 \oplus P_2$ and let π denote the projection map. For $z \in P$, set $\mathcal{V}_z P = \ker \pi_*|_z$. This defines a vector bundle \mathcal{V} over P , and by extending each ξ_k to be trivial on the other factor, we may regard ξ_1 and ξ_2 as vector fields on P . Moreover, the pair (η_1, η_2) defines a connection on P (see Foreman [2000b] for details). We denote the horizontal subspace determined by the connection at z by $\mathcal{H}_z P$ and the horizontal lift by $\tilde{\pi}$ or \tilde{X} .

The bundle space P carries an almost complex structure J defined as follows. On \mathcal{V} define J by $J\xi_1 = \xi_2$ and $J\xi_2 = -\xi_1$, and for $X \in \mathcal{H}_z P$, set $JX = \tilde{\pi} J_0 \pi_* X$. Since the Lie algebra $\mathfrak{s}^1 \oplus \mathfrak{s}^1$ is abelian, $[\xi_1, \xi_2] = 0$ and hence $[J, J](\xi_1, \xi_2) = 0$. Also it is easy to check that $[\xi_k, \tilde{X}] = 0$ and hence that $[J, J](\xi_k, \tilde{X}) = 0$. Finally, utilizing the integrability of J_0 on M one can readily show that $\pi_*[J, J](\tilde{X}, \tilde{Y}) = 0$. Therefore P is a complex manifold.

We can now define a complex contact structure on the complex manifold P . First note that $\eta_2 = -\eta_1 \circ J$, where again by extending each η_k to be trivial on the other factor we regard η_1 and η_2 as 1-forms on P . Set $\theta = \eta_1 + i\eta_2$; then θ is of type $(1, 0)$. Since Ω is holomorphic and $d\theta = \pi^*\Omega$, we see that θ is a holomorphic 1-form. Again since $d\theta = \pi^*\Omega$, a straightforward computation shows that $\theta \wedge (d\theta)^n \neq 0$. Thus P becomes a complex contact manifold with a global complex contact form.

We summarize the above construction in the following theorem.

Theorem 12.5 *Let M be a complex symplectic manifold with a complex symplectic form $\Omega = \Omega_1 + i\Omega_2$ such that both Ω_1 and Ω_2 determine integral classes. Then the $S^1 \times S^1$ -bundle defined by $([\Omega_1], [\Omega_2]) \in H^2(M, \mathbb{Z}) \oplus H^2(M, \mathbb{Z})$ has a complex contact structure given by a holomorphic connection form whose curvature form is Ω .*

For example, $\mathbb{C}P^n \times \mathbb{C}P^n$ is a complex symplectic manifold, and hence the Calabi–Eckmann manifold $S^{2n+1} \times S^{2n+1}$ carries a complex contact structure as described in the theorem. In [2000a] Foreman also gave examples for which the base complex symplectic manifold is a complex torus of even complex dimension.

Foreman next proved a converse to this theorem as a complex Boothby–Wang fibration; we state the result here and refer to Foreman [2000a] for the proof. For a global complex contact form θ we write $\theta = u - iv$, where u and v are real forms with $v = u \circ J$; the vertical bundle \mathcal{V} is then spanned by global vector fields U and $V = -JU$, where

$$\begin{aligned} u(U) &= 1, & v(U) &= 0, & \iota(U)du &= 0, \\ u(V) &= 0, & v(V) &= 1, & \iota(V)dv &= 0. \end{aligned}$$

Theorem 12.6 *Let P be a $(2n + 1)$ -dimensional compact complex contact manifold with a global form $\theta = u - iv$ such that the corresponding vertical vector fields U and V are regular. Then θ generates a free $S^1 \times S^1$ -action on P and $p : P \rightarrow M$ is a principal $S^1 \times S^1$ -bundle over a complex symplectic manifold M such that θ is a connection form for this fibration and the complex symplectic form Ω on M is given by $p^*\Omega = d\theta$.*

12.2.5 3-dimensional homogeneous examples

In the previous example we saw that B. Foreman had given a complex Boothby–Wang fibration for (regular) complex contact manifolds with a global contact form. In [1999] Foreman studied 3-dimensional complex homogeneous complex contact manifolds with a global complex contact form and obtained the following classification.

Theorem 12.7 *If M is a 3-dimensional complex homogeneous complex contact manifold with global complex contact form, then M is of the*

form $M = G/\Gamma$, where G is a simply connected 3-dimensional complex Lie group and $\Gamma \subset G$ is a discrete subgroup.

1. Suppose G is unimodular. Then G is one of the following:

- (a) $SL(2, \mathbb{C})$ if $\text{rk}(\text{ad}(\mathcal{V})) = 2$;
- (b) The universal cover of the group of rigid motions of the complex Euclidean plane if $\text{rk}(\text{ad}(\mathcal{V})) = 1$;
- (c) $H_{\mathbb{C}}$ if $\text{rk}(\text{ad}(\mathcal{V})) = 0$.

2. Suppose G is not unimodular. Then G is solvable, $\text{rk}(\text{ad}(\mathcal{V})) = 1$, and G is one of the following complex Lie groups:

- (a) The semidirect product $G_{\alpha} = \mathbb{C} \times_{\tau_{\alpha}} \mathbb{C}^2$, for any $\alpha \in \mathbb{C}^* \setminus 1$, where τ_{α} is the representation of \mathbb{C} in $GL(2, \mathbb{C})$ given by $\tau_{\alpha}(t) = \begin{pmatrix} e^{-t} & 0 \\ 0 & e^{-\alpha t} \end{pmatrix}$;
- (b) $G = \left\{ \begin{pmatrix} e^t & te^t & u \\ 0 & e^t & v \\ 0 & 0 & 1 \end{pmatrix} \mid t, u, v \in \mathbb{C} \right\}$.

The question of the regularity of the vertical foliation on compact quotients of $SL(2, \mathbb{C})$ by a discrete group was discussed by Foreman in [2010]. In Section 13.5 we will discuss complex contact structures on the Lie group $SL(2, \mathbb{C})$ in more detail.

12.2.6 Complex contact Lie groups

A *complex contact Lie group* is a $(2n+1)$ -dimensional complex Lie group G with a left-invariant holomorphic 1-form θ such that $\theta \wedge (d\theta)^n \neq 0$ on the complex manifold G . Define ξ in the Lie algebra \mathfrak{g} by $d\theta(\xi, \cdot) = 0$ and $\theta(\xi) = 1$. Then $\mathfrak{g} = \mathcal{V} \oplus \mathcal{H}$ with $\mathcal{V} = \langle \xi \rangle_{\mathbb{C}}$; ξ is called the *Reeb vector field* of θ .

Complex contact Lie groups in dimensions greater than 3 and for which the adjoint representation of ξ is diagonalizable were studied by Foreman in [2006]. His result is the following.

Theorem 12.8 *Let (G, θ) be a $(2n+1)$ -dimensional complex contact Lie group G such that $\text{ad}(\xi) : \mathfrak{g} \rightarrow \mathfrak{g}$ is diagonalizable. If $n > 1$, then the universal cover group of G is the semidirect product $\mathbb{C}^{2n} \times_{\Omega} \mathbb{C}$, where Ω is the standard symplectic form on \mathbb{C}^{2n} .*

12.2.7 $\mathbb{C}^{n+1} \times \mathbb{C}P^n(16)$

In [1998] B. Korkmaz gave a complex analogue of the real contact metric manifold $E^{n+1} \times S^n(4)$ together with a curvature characterization analogous to Theorem 7.5. We describe the example and state the theorem.

Let (t_0, \dots, t_n) be homogeneous coordinates on $\mathbb{C}P^n$ and \mathcal{U}_i the neighborhood defined by $t_i \neq 0$. On \mathcal{U}_i introduce nonhomogeneous coordinates by $w^j = \frac{t_j}{t_i}$, $j = 0, \dots, n$, $j \neq i$. Let $\mathcal{O}_i = \mathbb{C}^{n+1} \times \mathcal{U}_i$ and let (z^0, \dots, z^n) be coordinates on \mathbb{C}^{n+1} . Define a holomorphic 1-form θ_i on \mathcal{O}_i by $\theta_i = \frac{1}{t_i} \sum_{k=0}^n t_k dz^k$. Then $\theta_i \wedge (d\theta_i)^n \neq 0$ on \mathcal{O}_i and $\theta_j = \frac{t_i}{t_j} \theta_i$ on $\mathcal{O}_i \cap \mathcal{O}_j$. Thus $\{\theta_i\}_{i=0}^n$ is a complex contact structure on $\mathbb{C}^{n+1} \times \mathbb{C}P^n$.

For convenience we continue our work on \mathcal{O}_0 with $\theta_0 = dz^0 + \sum_{k=1}^n w^k dz^k$. The product metric on $\mathbb{C}^{n+1} \times \mathbb{C}P^n(16)$ is given by the matrix

$$g = \frac{1}{8} \left(\begin{array}{cc|cc} & & I_{n+1} & 0 \\ & O & 0 & \mathbf{g} \\ \hline I_{n+1} & 0 & & \\ 0 & \mathbf{g}^T & & O \end{array} \right),$$

where $\frac{1}{8}\mathbf{g}$ is the metric on $\mathbb{C}P^n(16)$, i.e.,

$$\mathbf{g}_{ij} = \frac{(1 + \sum_{k=1}^n |w^k|^2) \delta_{ij} - \bar{w}^i w^j}{(1 + \sum_{k=1}^n |w^k|^2)^2}.$$

Let $f_0 = 1 + \sum_{k=1}^n |w^k|^2$ and define real 1-forms u_0 and v_0 by

$$u_0 = \frac{1}{4\sqrt{f_0}} \left(dz^0 + d\bar{z}^0 + \sum_{k=1}^n (w^k dz^k + \bar{w}^k d\bar{z}^k) \right),$$

$$v_0 = \frac{i}{4\sqrt{f_0}} \left(dz^0 - d\bar{z}^0 + \sum_{k=1}^n (w^k dz^k - \bar{w}^k d\bar{z}^k) \right).$$

Similarly define vector fields U_0 and V_0 by

$$U_0 = \frac{2}{\sqrt{f_0}} \left(\frac{\partial}{\partial z^0} + \frac{\partial}{\partial \bar{z}^0} + \sum_{k=1}^n \left(\bar{w}^k \frac{\partial}{\partial z^k} + w^k \frac{\partial}{\partial \bar{z}^k} \right) \right),$$

$$V_0 = \frac{-2i}{\sqrt{f_0}} \left(\frac{\partial}{\partial z^0} - \frac{\partial}{\partial \bar{z}^0} + \sum_{k=1}^n \left(\bar{w}^k \frac{\partial}{\partial z^k} - w^k \frac{\partial}{\partial \bar{z}^k} \right) \right).$$

Then $\theta_0 = 2\sqrt{f_0}(u_0 - iv_0)$, $du_0(U_0, X) = 0$ for all $X \in \mathcal{H}$, $u_0(U_0) = 1$, $v_0(U_0) = 0$, and $g(U_0, X) = u_0(X)$ for all X .

Let

$$G_0 = f_0^{-\frac{3}{2}} \left(\begin{array}{cc|cc} & & 0 & A \\ & O & B & 0 \\ \hline 0 & \bar{A} & & \\ \bar{B} & 0 & & O \end{array} \right),$$

where

$$A = \begin{pmatrix} w^1 & w^2 & \cdots & w^n \\ |w^1|^2 - f_0 & \bar{w}^1 w^2 & \cdots & \bar{w}^1 w^n \\ w^1 \bar{w}^2 & |w^2|^2 - f_0 & \cdots & \bar{w}^2 w^n \\ \vdots & \vdots & \ddots & \vdots \\ w^1 \bar{w}^n & w^2 \bar{w}^n & \cdots & |w^n|^2 - f_0 \end{pmatrix},$$

$$B = f_0^{\frac{1}{2}} \begin{pmatrix} -w^1 \\ -w^2 \\ \vdots \\ -w^n \end{pmatrix} I_n.$$

Then $G_0^2 = -I + u_0 \otimes U_0 + v_0 \otimes V_0$, $G_0 J + J G_0 = 0$, etc. On \mathcal{O}_1 set $f_1 = 1 + |w^0|^2 + \sum_{k=2}^n |w^k|^2$. Then $\frac{f_0}{f_1} = \frac{|t_1|^2}{|t_0|^2}$. Setting $a - bi = \frac{t_0}{t_1} \sqrt{\frac{f_0}{f_1}}$ on $\mathcal{O}_0 \cap \mathcal{O}_1$, we have $a^2 + b^2 = 1$ and

$$u_1 = au_0 - bv_0, \quad v_1 = bu_0 + av_0, \quad G_1 = aG_0 - bH_0, \quad H_1 = bG_0 + aH_0,$$

where u_1, v_1, G_1, H_1 are the structure tensors on \mathcal{O}_1 . Therefore

$$\{(u_k, v_k, U_k, V_k, G_k, H_k, g)\} \text{ on } \{\mathcal{O}_k\}_{k=0}^n$$

is a complex contact metric structure on $\mathbb{C}^{n+1} \times \mathbb{C}P^n(16)$.

It can be shown that for this structure, $h_U = h_V$; see Korkmaz [1998]. We now state a characterization of this example due to Korkmaz [1998] analogous to Theorem 7.5 that a contact metric manifold on which $R_{XY}\xi = 0$ is locally isometric to $E^{n+1} \times S^n(4)$.

Theorem 12.9 *Let M be a complex contact metric manifold with $h_U = h_V$. If $R_{XY}\xi = 0$, then M is locally isometric to $\mathbb{C}^{n+1} \times \mathbb{C}P^n(16)$.*

$$12.2.8 \quad \cos z^3 dz^1 + \sin z^3 dz^2$$

In Example 3.2.6, Section 6.2, and Section 7.2 we studied the contact form

$$\frac{1}{2}(\cos x^3 dx^1 + \sin x^3 dx^2)$$

on \mathbb{R}^3 and T^3 with $\frac{1}{4}\delta_{ij}$ as a flat associated metric. We can similarly consider the complex contact form

$$\theta = \frac{1}{2}(\cos z^3 dz^1 + \sin z^3 dz^2)$$

on \mathbb{C}^3 , $z^k = x^k + iy^k$, $k = 1, 2, 3$. For this structure,

$$U + iV = 4\left(\cos z^3 \frac{\partial}{\partial z^1} + \sin z^3 \frac{\partial}{\partial z^2}\right),$$

and an associated metric is given by

$$(g_{\lambda\bar{\mu}}) = \frac{1}{8} \begin{pmatrix} \operatorname{sech} 2y^3 & 0 & 0 \\ 0 & \operatorname{sech} 2y^3 & 0 \\ 0 & 0 & 1 \end{pmatrix}.$$

The other structure tensors can be readily computed (cf. Jayne [1992, pp. 222–223]). One can also compute the curvature of this metric quite readily and one finds that the metric is not flat.

In the real case we have noted that aside from the flat 3-dimensional case, there are no flat associated metrics in any dimension. In the complex case we conjecture that there are no flat complex contact metric structures at all. In view of Olszak's result that the only contact metric manifolds of constant curvature and dimension ≥ 5 are of constant curvature $+1$ (Theorem 7.3), we also conjecture that aside from odd-dimensional complex projective space, there are no complex contact metric structures of constant holomorphic curvature.

12.3 Normality of complex contact manifolds

As we have seen, in real contact geometry the product $M \times \mathbb{R}$ of an almost contact manifold and the real line carries a natural almost complex structure J , and if J is integrable, the structure is said to be normal. Recall also that a Sasakian manifold is a normal contact metric manifold.

Ishihara and Konishi [1979], [1980] introduced a notion of normality for complex contact structures. Their notion is the vanishing of the two tensor fields S and T given by

$$\begin{aligned} S(X, Y) &= [G, G](X, Y) + 2\hat{G}(X, Y)U - 2\hat{H}(X, Y)V \\ &\quad + 2(v(Y)HX - v(X)HY) + \sigma(GY)HX - \sigma(GX)HY \\ &\quad + \sigma(X)GHY - \sigma(Y)GHX, \\ T(X, Y) &= [H, H](X, Y) - 2\hat{G}(X, Y)U + 2\hat{H}(X, Y)V \\ &\quad + 2(u(Y)GX - u(X)GY) + \sigma(HX)GY - \sigma(HY)GX \\ &\quad + \sigma(X)GHY - \sigma(Y)GHX. \end{aligned}$$

However, this notion seems to be too strong; among its implications is that the underlying Hermitian manifold (M, g) is Kähler. Thus while indeed one of the canonical examples of a complex contact manifold, the odd-dimensional complex projective space, is normal in this sense, the complex Heisenberg group is not. B. Korkmaz [2000] generalized the notion of normality, and we adopt her definition in this book. A complex contact metric structure is *normal* if

$$\begin{aligned} S(X, Y) = T(X, Y) &= 0, \text{ for every } X, Y \in \mathcal{H}, \\ S(U, X) = T(V, X) &= 0, \text{ for every } X. \end{aligned}$$

Even though the definition appears to depend on the special nature of U and V , it respects the change in overlaps $\mathcal{O}_\alpha \cap \mathcal{O}_\beta$ and is a global notion (Korkmaz [2000]). With this notion of normality both odd-dimensional complex projective space and the complex Heisenberg group with their standard complex contact metric structures are normal. We remark that for Ishihara and Konishi's notion of normality, the holonomy group is the full unitary group $U(2m + 1)$ Houh [1976]. Also, Ishihara and Konishi's notion of normality is equivalent to the curvature condition in Theorem 12.4, Foreman [2000b].

One consequence of normality is that $h_U = 0$ for every $U \in \mathcal{V}$; this is analogous to the fact that every Sasakian structure is K-contact. Another is that the sectional curvature of a plane section spanned by a vector in \mathcal{V} and a vector in \mathcal{H} is equal to +1 (cf. Korkmaz [2000], Foreman [2000b]).

We now give expressions for the covariant derivatives of G and H on a normal complex contact metric manifold; for proofs see Korkmaz [2000]. A complex contact metric manifold is normal if and only if the covariant

derivatives of G and H have the following forms:

$$\begin{aligned}
g((\nabla_X G)Y, Z) &= \sigma(X)g(HY, Z) + v(X)d\sigma(GZ, GY) \\
&\quad - 2v(X)g(HGY, Z) - u(Y)g(X, Z) - v(Y)g(JX, Z) \\
&\quad + u(Z)g(X, Y) + v(Z)g(JX, Y). \\
g((\nabla_X H)Y, Z) &= -\sigma(X)g(GY, Z) - u(X)d\sigma(HZ, HY) \\
&\quad - 2u(X)g(GHY, Z) + u(Y)g(JX, Z) - v(Y)g(X, Z) \\
&\quad - u(Z)g(JX, Y) + v(Z)g(X, Y).
\end{aligned}$$

In these formulas the first two terms on the right vanish for the complex Heisenberg group (Example 12.2.1), and the second and third terms cancel on $P\mathbb{C}^{2n+1}$ (Example 12.2.2). Also one has

$$\nabla_X U = -GX, \quad \nabla_X V = -HX$$

on $H_{\mathbb{C}}$ and

$$\nabla_X U = -GX + \sigma(X)V, \quad \nabla_X V = -HX - \sigma(X)U$$

on $\mathbb{C}P^{2n+1}$. Finally, on a normal complex contact manifold we have

$$\begin{aligned}
g((\nabla_X J)Y, Z) \\
= u(X)(d\sigma(Z, GY) - 2g(HY, Z)) + v(X)(d\sigma(Z, HY) + 2g(GY, Z)).
\end{aligned}$$

When the complex contact structure is strict, i.e., given by a global complex contact form, the situation is more restrictive. In particular, $\sigma = 0$ and therefore some of the above formulas simplify. Foreman [2000b] defines a *complex Sasakian manifold* to be a normal complex contact metric manifold whose complex contact structure is given by a global complex contact form. His paper gives a number of examples and basic properties including local projectivity to a hyper-kähler manifold.

12.4 GH -sectional curvature

Corresponding to the ideas of holomorphic curvature in complex geometry and ϕ -sectional curvature in real contact geometry, B. Korkmaz [2000] defined the notion of GH -sectional curvature for a complex contact metric manifold. For a unit vector $X \in \mathcal{H}_m$, the plane in $T_m M$ spanned by X

and $Y = aGX + bHX$, $a, b \in \mathbb{R}$, $a^2 + b^2 = 1$, is called a GH -plane section, and its sectional curvature, $K(X, Y)$, the GH -sectional curvature of the plane section. For a given vector X , $K(X, Y)$ is independent of the vector Y in the plane of GX and HX if and only if $K(X, GX) = K(X, HX)$ and $g(R_{XGX}HX, X) = 0$. Let M be a normal complex contact metric manifold; if the GH -sectional curvature is independent of the choice of GH -section at each point, it is constant on the manifold, and we say that M is a *complex contact space form*. The curvature tensor and the following theorems were obtained by Korkmaz [2000]; explicitly, the curvature tensor is

$$\begin{aligned}
 R_{XY}Z = & \frac{c+3}{4} \left(g(Y, Z)X - g(X, Z)Y \right. \\
 & + g(Z, JY)JX - g(Z, JX)JY + 2g(X, JY)JZ \Big) \\
 & + \frac{c-1}{4} \left(- (u(Y)u(Z) + v(Y)v(Z))X + (u(X)u(Z) + v(X)v(Z))Y \right. \\
 & + 2u \wedge v(Z, Y)JX - 2u \wedge v(Z, X)JY + 4u \wedge v(X, Y)JZ \\
 & + g(Z, GY)GX - g(Z, GX)GY + 2g(X, GY)GZ \\
 & + g(Z, HY)HX - g(Z, HX)HY + 2g(X, HY)HZ \\
 & + (-u(X)g(Y, Z) + u(Y)g(X, Z) \\
 & + v(X)g(JY, Z) - v(Y)g(JX, Z) + 2v(Z)g(X, JY))U \\
 & + (-v(X)g(Y, Z) + v(Y)g(X, Z) \\
 & - u(X)g(JY, Z) + u(Y)g(JX, Z) - 2u(Z)g(X, JY))V \Big) \\
 & - \frac{4}{3} (d\sigma(U, V) + c + 1) \left((v(X)u \wedge v(Z, Y) - v(Y)u \wedge v(Z, X) \right. \\
 & + 2v(Z)u \wedge v(X, Y))U - (u(X)u \wedge v(Z, Y) \\
 & - u(Y)u \wedge v(Z, X) + 2u(Z)u \wedge v(X, Y))V \Big).
 \end{aligned}$$

Odd-dimensional complex projective space with the Fubini–Study metric of constant holomorphic curvature 4 is of constant GH -sectional curvature 1. The complex Heisenberg group has holomorphic curvature 0 for horizontal and vertical holomorphic sections and constant GH -sectional curvature -3 .

Theorem 12.10 *Let M be a normal complex contact metric manifold. Then M has constant GH -sectional curvature c if and only if for X*

horizontal, the holomorphic sectional curvature of the plane spanned by X and JX is $c + 3$.

Theorem 12.11 *Let M be a normal complex contact metric manifold of constant GH -sectional curvature $+1$ and satisfying $d\sigma(V, U) = 2$. Then M has constant holomorphic curvature 4 . If, in addition, M is complete and simply connected, then M is isometric to $\mathbb{C}P^{2n+1}$ with the Fubini–Study metric of constant holomorphic curvature 4 .*

Korkmaz [2000] then introduced the idea of an \mathcal{H} -homothetic deformation of a complex contact metric structure. Let α be a positive constant and consider the local structure tensors (G, H, U, V, u, v, g) . Then define new structure tensors by

$$\begin{aligned}\tilde{u} &= \alpha u, & \tilde{v} &= \alpha v, & \tilde{U} &= \frac{1}{\alpha}U, & \tilde{V} &= \frac{1}{\alpha}V, & \tilde{G} &= G, & \tilde{H} &= H, \\ \tilde{g} &= \alpha g + \alpha(\alpha - 1)(u \otimes u + v \otimes v).\end{aligned}$$

This change of structure is called an \mathcal{H} -homothetic deformation. The new structure then respects the transitions on the overlaps of coordinate neighborhoods and hence gives a new complex contact metric structure on M . Moreover, $\tilde{S}(X, Y) = S(X, Y)$ on \mathcal{H} , $\tilde{S}(X, \tilde{U}) = \frac{1}{\alpha}S(X, U)$, etc., so if the given structure is normal, so is the new structure. Korkmaz computed the curvature and showed that if on a normal complex contact metric manifold the original structure has constant GH -sectional curvature c , then the new structure has constant GH -sectional curvature $\tilde{c} = \frac{c+3}{\alpha} - 3$; in particular, she proved the following results.

Theorem 12.12 *Complex projective space $\mathbb{C}P^{2n+1}$ carries a normal complex contact metric structure with constant GH -section curvature $\frac{4}{\alpha} - 3$ for every $\alpha > 0$.*

Theorem 12.13 *A normal complex contact metric manifold with metric \tilde{g} of constant GH -sectional curvature $\tilde{c} > -3$ is \mathcal{H} -homothetic to a normal complex contact metric manifold with metric g of constant GH -section curvature $c = 1$. Moreover, if $d\sigma(\tilde{V}, \tilde{U}) = (\tilde{c} + 3)^2/8$, the metric g is Kähler and has constant holomorphic curvature 4 .*

In [2003] B. Korkmaz continued her study of the curvature of complex contact metric manifolds and of \mathcal{H} -homothetic deformations; in particular, she developed a theory of complex (κ, μ) -spaces analogous to that described in Section 7.3 for the real contact geometry.

12.5 The set of associated metrics and integral functionals

In [1996] B. Foreman began the study of the set of all associated metrics on a complex contact manifold. In Chapter 4 we studied how the set of associated metrics \mathcal{A} for a symplectic or contact form sits in the set of Riemannian metrics with the same volume element, and in particular we studied the tangent space to \mathcal{A} at an associated metric g . As before, we will typically use the same letter for a symmetric tensor field as a tangent vector to a curve of metrics and for the corresponding tensor field of type $(1, 1)$ determined by the metric.

As already mentioned, and in keeping with the notation in the real case, we will denote by \mathcal{A} the set of all associated metrics for a complex contact structure as defined in Section 12.1. As in the real case, \mathcal{A} is infinite dimensional; all associated metrics to a given complex contact structure have the same volume element; \mathcal{A} is totally geodesic in \mathcal{N} in the sense that if $D \in T_g\mathcal{A}$, the geodesic $g_t = ge^{tD}$ is a path in \mathcal{A} ; and two metrics in \mathcal{A} may be joined by a geodesic. We now give a characterization of $T_g\mathcal{A}$; for details of this and the preceding statements see Foreman [1996].

Lemma 12.1 *Let g be a metric in \mathcal{A} . Then a symmetric tensor field D of type $(0, 2)$ is in $T_g\mathcal{A}$ if and only if D , as a tensor field of type $(1, 1)$, satisfies*

$$DJ = JD, \quad D|_{\mathcal{V}} = 0, \quad DG = -GD$$

for any local tensor field G as in the preceding sections.

As before, to study integral functionals defined on \mathcal{A} we will be differentiating such functionals along paths of metrics, and for this we need the following fundamental lemma (Foreman [1996]). For an endomorphism T of a complex vector space, set

$$T^s = \frac{1}{2}(T - JTJ), \quad T^d = \frac{1}{2}(T + JTJ),$$

i.e., T^s is the part of T that commutes with J , and T^d the part of T that anticommutes with J .

Lemma 12.2 *For $g \in \mathcal{A}$ and T a symmetric $(1, 1)$ tensor field,*

$$\int_M \operatorname{tr} TD \, dV = 0$$

for every $D \in T_g \mathcal{A}$ if and only if

$$p(TJ + JT)p = HTG - GTH,$$

or equivalently

$$pT^s p = -GT^s G.$$

In Section 10.3 we studied the integral of the Ricci curvature in the direction of the characteristic vector field ξ , i.e., $L(g) = \int_M Ric(\xi) dV$, as a functional on the set of associated metrics. In [1996] Foreman gave two analogues of $Ric(\xi)$ for complex contact manifolds and proved the following results. For a unit vertical vector field U , $Ric(U) + Ric(JU)$ is global, i.e., it respects transition on $\mathcal{O}_\alpha \cap \mathcal{O}_\beta$, and we denote it by $Ric(\mathcal{V})$. For the second analogue, Foreman utilizes the $*$ -Ricci tensor, and it is easy to check that for a unit vertical vector field U , $Ric^*(U) = R_{ij}^* U^i U^j$ is independent of the unit vertical vector field U and globally defined; we denote it by $Ric^*(\mathcal{V})$. Define functionals L and L^* on \mathcal{A} by

$$L(g) = \int_M Ric(\mathcal{V}) dV, \quad L^*(g) = \int_M Ric^*(\mathcal{V}) dV.$$

Theorem 12.14 *Let M be a compact complex contact manifold and \mathcal{A} the set of associated metrics. Then $g \in \mathcal{A}$ is a critical point of $L(g)$ if and only if*

$$(\nabla_U h_U)^s + (\nabla_V h_V)^s = \sigma(U)h_V^s - \sigma(V)h_U^s - 4Gh_U^d$$

for any local unit vertical vector field U .

Clearly, any projectable associated metric is a critical point of $L(g)$. In fact, Foreman showed that $Ric(\mathcal{V}) = 8n - 2K(\mathcal{V}) - \text{tr}h_U^2 - \text{tr}h_V^2$, where $K(\mathcal{V})$ is the sectional curvature of a vertical plane section, and moreover that $\int_M K(\mathcal{V}) dV$ is independent of the associated metric. Thus projectable metrics are maxima of $L(g)$. If g is Kähler, $h_U^s = 0$ for any vertical vector field U , and moreover, $(\nabla_W h_U^d)^s = 0$ for any vertical vector field W . Thus we also have the following corollary.

Corollary 12.1 *If $g \in \mathcal{A}$ is Kähler and a critical point of $L(g)$, then it is projectable.*

Theorem 12.15 *Let M be a compact complex contact manifold and \mathcal{A} the set of associated metrics. Then $g \in \mathcal{A}$ is a critical point of $L^*(g)$ if and only if*

$$h_U^d(\nabla_U J) = 0$$

for any local unit vertical vector field U .

In particular, if g is projectable or Kähler, it is critical. For the twistor spaces as described in Example 12.2.3 this theorem may be improved as follows. If Z is the twistor space of a compact quaternionic Kähler manifold M^{4n} , $4n \geq 8$, with normalized complex contact structure $\{u_c - iv_c\}$ and if the Salamon, Bérard-Bergery metric g_c is an associated metric, then an associated metric g is critical for L^* if and only if $h_U^d = 0$ for every vertical vector U .

B. Foreman also obtained some important results on the constancy of L^* , i.e., when L^* is independent of the associated metric.

Theorem 12.16 *Let M be a complex contact manifold with corresponding almost complex structure J . Then L^* is constant on \mathcal{A} if and only if J is projectable.*

Theorem 12.17 *Suppose M is a compact complex contact manifold with a global complex contact structure and \mathcal{A} the set of associated metrics. Then L^* is constant on \mathcal{A} .*

Turning now to the scalar curvatures, in real contact geometry the “total scalar curvature” defined by $\int_M \tau + \tau^* dV$ is a functional on \mathcal{A} whose critical points are precisely the K-contact metrics, as we have seen (Theorem 10.9). B. Foreman (unpublished) has also defined a ***scalar curvature* by first contracting the curvature with the local tensor fields G and H as in the **-scalar curvature*, giving **-scalar curvatures* τ_G^* and τ_H^* ; $\tau^{**} = \tau_G^* + \tau_H^*$ is then globally defined, and one defines the “total scalar curvature” $I(g)$ by $I(g) = \int_M \tau + \tau^* + \tau^{**} dV$. Recall that an almost Hermitian structure (J, g) is *semi-Kähler* if the fundamental 2-form Ω is coclosed. Foreman computed the critical point condition for this functional and though complicated, it yields the following result.

Theorem 12.18 *A projectable semi-Kähler metric is a critical point of the functional $I(g)$.*

12.6 Holomorphic Legendre curves

Let \tilde{M} be a Hermitian manifold of complex dimension n with complex structure J and corresponding Riemannian metric g . Following Chern, Cowen and Vitter [1974] and S. Dolbeault [1977], we describe holomorphic curves and Frenet frames. A holomorphic curve in \tilde{M} is a nonconstant holomorphic map $\iota : M \rightarrow \tilde{M}$, where M is a Riemann surface.

If $z_i = z_i(w)$ is a local representation of M in a neighborhood of $w = 0$, then its holomorphic tangent vector at $\iota(0)$ is given by $\sum z'_i(0) \frac{\partial}{\partial z_i} = w^p V$, where p is a nonnegative integer and V a nonzero vector. The isolated points where $p > 0$ are called *stationary points of order 0*. A unitary frame $\{f_1, \dots, f_n\}$ is called a *Frenet frame* if $f_1 = \frac{V}{|V|}$ and

$$\tilde{\nabla}_X f_i = \omega_{ii-1}(X) f_{i-1} + \omega_{ii}(X) f_i + \omega_{ii+1}(X) f_{i+1}$$

for $i = 1$ to $n - 1$, where ω_{ii+1} is a holomorphic 1-form and $\omega_{i+1i}(X) = -\overline{\omega_{ii+1}(X)}$. Points where ω_{ii+1} vanish are called *stationary points of order i* . In general, a unitary frame is not holomorphic, but we will be interested in the case in which the f_i 's are holomorphic vector fields, and we then speak of a *holomorphic Frenet curve* and a *holomorphic Frenet frame*.

Not every holomorphic curve in a Hermitian manifold \tilde{M} has a Frenet frame. Chern, Cowen and Vitter [1974] (or see Dolbeault [1977]) give curvature conditions on \tilde{M} under which every holomorphic curve has a Frenet frame; in particular, a Kähler manifold of complex dimension ≥ 3 has a Frenet frame along every holomorphic curve if and only if it has constant holomorphic curvature.

M being a holomorphic Frenet curve has implications on M as a submanifold. In the following lemma we give three such implications, the first two of which are immediate in any Kähler manifold. For our purpose we give the lemma only for Hermitian manifolds of complex dimension 3 (see Baikoussis, Gouli-Andreou and the author [1998] for the proof). Recall that the span of the second fundamental form α is called the *first normal space* and will be denoted by ν_1 . Let 'proj' denote projection to the orthogonal complement of $TM \oplus \nu_1$. Define $\beta(X, Y, Z)$ by

$$\beta(X, Y, Z) = \text{proj} \tilde{\nabla}_X \tilde{\nabla}_Y Z (= \text{proj} \nabla_X^\perp \alpha(Y, Z)).$$

The span of β is called the *second normal space* and will be denoted by ν_2 . Successively the higher normal spaces may be defined in this manner taking higher order derivatives.

Lemma 12.3 *Let M be a holomorphic Frenet curve in a 3-dimensional Hermitian manifold (\tilde{M}, J, g) with second fundamental form α . Let \tilde{R} denote the curvature tensor of \tilde{M} . Then*

- (1) $\alpha(X, JY) = J\alpha(X, Y)$ for any tangent vectors X, Y , and hence the real dimension of the first normal space is 2;

$$(2) \quad \tilde{\nabla}_X J|_{TM \oplus T^\perp M} = 0;$$

(3) for any tangent vectors X, Y, Z , we have that $\tilde{R}(X, Y)Z$ is orthogonal to the second normal space.

Conversely, if M is a holomorphic curve satisfying (1), (2), and (3), then it has a holomorphic Frenet frame.

Related to the idea of a Frenet frame are the curvatures themselves. These Frenet curvatures for a holomorphic curve date back to Calabi [1953a], who defined $(n - 1)$ real-valued curvature functions (see also Lawson [1970]). When the ambient space is a complex space form, these curvatures are actually intrinsic (Lawson [1970]). We follow the development as presented by Lawson.

Let M be a holomorphic curve in a 3-dimensional Hermitian manifold \tilde{M} for which the properties of the lemma hold. From $\alpha(X, JY) = J\alpha(X, Y)$ we have easily that for all unit tangent vectors X, Y (Y not necessarily distinct from X), $|\alpha(X, Y)|^2$ is a function of position alone, say $\kappa_1(p)$, $p \in M$; κ_1 is called the *curvature* or *first curvature* of M . Now with $\beta(X, Y, Z)$ defined as above, from property (2) of the lemma we have $\beta(X, Y, JZ) = J\beta(X, Y, Z)$. From the second expression for β in the definition we see that β is symmetric in the second and third variables, giving $\beta(X, JY, Z) = J\beta(X, Y, Z)$. Let ν be a vector in the second normal space. Then by property (3), $g(\tilde{\nabla}_X \tilde{\nabla}_Y Z, \nu) = g(\tilde{\nabla}_Y \tilde{\nabla}_X Z, \nu)$, giving that β is symmetric in the first and second variables and therefore $\beta(JX, Y, Z) = J\beta(X, Y, Z)$. From these properties of β we have that

$$\kappa_2(p) = \frac{|\beta(X, Y, Z)|^2}{\kappa_1(p)}$$

is well defined, X, Y, Z being any unit tangent vectors. We call κ_2 the *torsion* or *second curvature* and also denote it by τ .

It is interesting to compare the curvature and torsion with the derivatives of the Frenet frames and the holomorphic connection forms ω_{ii+1} . In particular, writing f_1 as $\frac{e_1 - iJe_1}{\sqrt{2}}$, we have $\kappa = |\omega_{12}(e_1)|^2$ and $\tau = |\omega_{23}(e_1)|^2$.

We begin with the following proposition from Baikoussis, Gouli-Andreou and the author [1998].

Proposition 12.1 *Let M be a real surface in $(H_{\mathbb{C}}, \theta, g)$ such that $\theta(X) = 0$ for any tangent vector X . Then M is a holomorphic Legendre curve as well as a holomorphic Frenet curve with torsion $\tau \equiv 1$.*

We now state two results, due to Baikoussis, Gouli-Andreou and the author [1998].

Theorem 12.19 *If the torsion of a holomorphic Frenet curve in the complex Heisenberg group is not identically zero and at one point the complex contact form annihilates the tangent space, then the curve is a holomorphic Legendre curve.*

Theorem 12.20 *Let M be a holomorphic Legendre curve in $(H_{\mathbb{C}}, \theta, g)$ and N its projection to $\mathbb{C}^2 = \{(z_1, z_2)\}$ with its standard complex structure and Kähler (Euclidean) metric. Then the Gaussian curvature of M is 8 times that of N .*

Turning to the complex contact manifold \mathbb{CP}^3 , we give a complete characterization of holomorphic Legendre curves due to Bryant [1982].

Theorem 12.21 *Let M be a connected Riemann surface and let f and g be meromorphic functions on M with g nonconstant. In terms of homogeneous coordinates define $\iota : M \rightarrow \mathbb{CP}^3$ by*

$$\zeta \longrightarrow \left(1, g, f - g \frac{f'}{2g'}, \frac{f'}{2g'}\right).$$

Then $\iota : M \rightarrow \mathbb{CP}^3$ is a holomorphic Legendre curve. Conversely, any holomorphic Legendre curve is either of this form or has its image in some $\mathbb{CP}^1 \subset \mathbb{CP}^3$.

Specifically, if (z_1, z_2, w_1, w_2) are homogeneous coordinates on \mathbb{CP}^3 and $\psi = z_1 dw_1 - w_1 dz_1 + z_2 dw_2 - w_2 dz_2$ as in Example 12.2.2, then it is easy to check that $\psi(\iota_* \frac{\partial}{\partial \zeta}) = 0$.

As an application of his result, Bryant uses the fibration of \mathbb{CP}^3 over S^4 to show that given a compact Riemann surface, there exists a conformal superminimal generically one-to-one immersion into S^4 whose image in S^4 is an algebraic surface.

12.7 The Calabi (Veronese) embeddings as integral submanifolds of \mathbb{CP}^{2n+1}

In [1953b] Calabi showed that up to holomorphic congruence there is a unique holomorphic embedding of $\mathbb{CP}^n(\frac{4}{\nu})$ into $\mathbb{CP}^N(4)$, $N = \binom{n+\nu}{\nu} - 1$

that does not lie in any totally geodesic complex projective space of lower dimension. Nakagawa and Ogiue [1976] showed that the only full isometric immersions of positively curved complex space forms into positively curved complex space forms are local versions of these embeddings. These embeddings are known as the *Calabi embeddings* or as the *Veronese embeddings*, especially in the case $\mathbb{C}P^2(2) \longrightarrow \mathbb{C}P^5(4)$. For $n = 1$ these embeddings are called *Calabi curves*. Classically the Calabi embeddings are given as follows. Let $\zeta_1, \dots, \zeta_{n+1}$ be homogeneous coordinates for $\mathbb{C}P^n(\frac{4}{\nu})$. The Calabi embedding of $\mathbb{C}P^n(\frac{4}{\nu})$ into $\mathbb{C}P^N(4)$, in terms of homogeneous coordinates for $\mathbb{C}P^N(4)$, is given by

$$(\zeta_1, \dots, \zeta_{n+1}) \longrightarrow \left(\zeta_1^\nu, \sqrt{\nu} \zeta_1^{\nu-1} \zeta_2, \dots, \sqrt{\frac{\nu!}{a_1! \dots a_{n+1}!}} \zeta_1^{a_1} \dots \zeta_{n+1}^{a_{n+1}}, \dots, \zeta_{n+1}^\nu \right),$$

where $\sum_{i=1}^{n+1} a_i = \nu$, the a_i 's being nonnegative integers. The meaning of the integer ν is that there are $\nu - 1$ normal spaces for these embeddings.

The question to be addressed in this section is the following. For which of these embeddings is there a holomorphic congruence of $\mathbb{C}P^{2n+1}$ that positions the submanifold as a holomorphic integral submanifold of the complex contact structure?

For the Calabi curves we have the following positive answer due to Dillen, Verstraelen, Vrancken and the author [1996].

Theorem 12.22 *There exists a holomorphic congruence of $\mathbb{C}P^{2n+1}(4)$ that positions the Calabi curve $\mathbb{C}P^1(\frac{4}{2n+1})$ as a holomorphic Legendre curve in $\mathbb{C}P^{2n+1}(4)$.*

The embedding is given explicitly as follows. For simplicity set $A_k = \sqrt{\binom{2n+1}{k-1}}$; now position $\mathbb{C}P^1(\frac{4}{2n+1})$ in $\mathbb{C}P^{2n+1}(4)$ by

$$(\zeta_1, \zeta_2) \longrightarrow \left(\zeta_1^{2n+1}, \dots, A_k \zeta_1^{2n+2-k} \zeta_2^{k-1}, \dots, A_{n+1} \zeta_1^{n+1} \zeta_2^n, \right. \\ \left. \zeta_2^{2n+1}, \dots, (-1)^{k-1} A_k \zeta_2^{2n+2-k} \zeta_1^{k-1}, \dots, (-1)^n A_{n+1} \zeta_2^{n+1} \zeta_1^n \right).$$

Then

$$\begin{aligned}\frac{\partial}{\partial \zeta_1} &= \sum_{k=1}^{n+1} \left(A_k(2n+2-k) \zeta_1^{2n+1-k} \zeta_2^{k-1} \frac{\partial}{\partial z_k} \right. \\ &\quad \left. + (-1)^{k-1} A_k(k-1) \zeta_2^{2n+2-k} \zeta_1^{k-2} \frac{\partial}{\partial w_k} \right), \\ \frac{\partial}{\partial \zeta_2} &= \sum_{k=1}^{n+1} \left(A_k(k-1) \zeta_1^{2n+2-k} \zeta_2^{k-2} \frac{\partial}{\partial z_k} \right. \\ &\quad \left. + (-1)^{k-1} A_k(2n+2-k) \zeta_2^{2n+1-k} \zeta_1^{k-1} \frac{\partial}{\partial w_k} \right)\end{aligned}$$

represent the tangent space, and the proof is to show that $\psi(\frac{\partial}{\partial \zeta_1}) = \psi(\frac{\partial}{\partial \zeta_2}) = 0$, where $\psi = \sum_{k=1}^{n+1} (z_k dw_k - w_k dz_k)$ as in Example 12.2.2.

On the other hand, there is no holomorphic congruence of $\mathbb{CP}^5(4)$ that brings the Veronese surface $\mathbb{CP}^2(2)$ into position as a Legendre submanifold of the complex contact structure on $\mathbb{CP}^5(4)$ even though the codimension is large enough. This has to do with the fact that the integer ν is equal to 2.

Theorem 12.23 *Assume that $N = \binom{n+2}{2} - 1$ is odd. There is no holomorphic congruence of $\mathbb{CP}^N(4)$ that brings the Calabi embedding of $\mathbb{CP}^n(2)$ into position as an integral submanifold of the complex contact structure on $\mathbb{CP}^N(4)$.*

Proof. Recall that the meaning of the condition $\nu = 2$ is that the first normal space is the whole normal space. Suppose now that $\mathbb{CP}^n(2) \longrightarrow \mathbb{CP}^N(4)$ is an integral submanifold of the complex contact structure with second fundamental form α . We have already noted that the vector fields U and V are normal and that for any tangent vector X , GX is normal. Using $\tilde{\nabla}_X U = -GX + \sigma(X)V$, we have

$$0 = Xg(Y, U) = g(\tilde{\nabla}_X Y, U) - g(Y, GX) = g(\alpha(X, Y), U),$$

and similarly $g(\alpha(X, Y), V) = 0$. Thus U and V are orthogonal to both the tangent space and the first normal space, but the first normal space is the whole normal space, giving a contradiction. ■

For $\nu = 3$ we first give a nonexistence result for $n = 2$ and then a positive result for n odd; these results are due to Korkmaz, Vrancken and the author [2000].

Theorem 12.24 *There is no holomorphic congruence of $\mathbb{CP}^9(4)$ that brings the Calabi embedding of $\mathbb{CP}^2(\frac{4}{3})$ into position as an integral submanifold of the complex contact structure on $\mathbb{CP}^9(4)$.*

Theorem 12.25 *When $\nu = 3$ and n is odd, $N = \binom{n+3}{3} - 1$ is odd and there exists a holomorphic congruence of $\mathbb{CP}^N(4)$ that brings the Calabi embedding of $\mathbb{CP}^n(\frac{4}{3})$ into position as an integral submanifold of the complex contact structure on $\mathbb{CP}^N(4)$.*

Finally, we give an example with $\nu = 5$. In terms of the homogeneous coordinates $(z_1, \dots, z_{28}, w_1, \dots, w_{28})$, $\mathbb{CP}^3(\frac{4}{5})$ may be realized as an integral submanifold of the complex contact structure on $\mathbb{CP}^{55}(4)$ in the following way:

$$\begin{array}{ll}
 z_1 = \zeta_1^5 & w_1 = -\zeta_3^5 \\
 z_2 = \zeta_2^5 & w_2 = -\zeta_4^5 \\
 z_3 = \sqrt{5}\zeta_1^4\zeta_2 & w_3 = -\sqrt{5}\zeta_3^4\zeta_4 \\
 z_4 = \sqrt{5}\zeta_1^4\zeta_3 & w_4 = \sqrt{5}\zeta_3^4\zeta_1 \\
 z_5 = \sqrt{5}\zeta_1^4\zeta_4 & w_5 = \sqrt{5}\zeta_3^4\zeta_2 \\
 z_6 = \sqrt{5}\zeta_2^4\zeta_1 & w_6 = -\sqrt{5}\zeta_4^4\zeta_3 \\
 z_7 = \sqrt{5}\zeta_2^4\zeta_3 & w_7 = \sqrt{5}\zeta_4^4\zeta_1 \\
 z_8 = \sqrt{5}\zeta_2^4\zeta_4 & w_8 = \sqrt{5}\zeta_4^4\zeta_2 \\
 z_9 = \sqrt{10}\zeta_1^3\zeta_2^2 & w_9 = -\sqrt{10}\zeta_3^3\zeta_4^2 \\
 z_{10} = \sqrt{10}\zeta_1^3\zeta_3^2 & w_{10} = -\sqrt{10}\zeta_3^3\zeta_1^2 \\
 z_{11} = \sqrt{10}\zeta_1^3\zeta_4^2 & w_{11} = -\sqrt{10}\zeta_3^3\zeta_2^2 \\
 z_{12} = \sqrt{10}\zeta_2^3\zeta_1^2 & w_{12} = -\sqrt{10}\zeta_4^3\zeta_3^2 \\
 z_{13} = \sqrt{10}\zeta_2^3\zeta_3^2 & w_{13} = -\sqrt{10}\zeta_4^3\zeta_1^2 \\
 z_{14} = \sqrt{10}\zeta_2^3\zeta_4^2 & w_{14} = -\sqrt{10}\zeta_4^3\zeta_2^2 \\
 z_{15} = \sqrt{20}\zeta_1^3\zeta_2\zeta_3 & w_{15} = \sqrt{20}\zeta_3^3\zeta_1\zeta_4 \\
 z_{16} = \sqrt{20}\zeta_1^3\zeta_2\zeta_4 & w_{16} = \sqrt{20}\zeta_3^3\zeta_2\zeta_4 \\
 z_{17} = \sqrt{20}\zeta_1^3\zeta_3\zeta_4 & w_{17} = -\sqrt{20}\zeta_3^3\zeta_1\zeta_2 \\
 z_{18} = \sqrt{20}\zeta_2^3\zeta_1\zeta_3 & w_{18} = \sqrt{20}\zeta_4^3\zeta_1\zeta_3 \\
 z_{19} = \sqrt{20}\zeta_2^3\zeta_1\zeta_4 & w_{19} = \sqrt{20}\zeta_4^3\zeta_2\zeta_3
 \end{array}$$

$$z_{20} = \sqrt{20}\zeta_2^3\zeta_3\zeta_4$$

$$z_{21} = \sqrt{30}\zeta_1^2\zeta_2^2\zeta_3$$

$$z_{22} = \sqrt{30}\zeta_1^2\zeta_2^2\zeta_4$$

$$z_{23} = \sqrt{30}\zeta_1^2\zeta_3^2\zeta_2$$

$$z_{24} = \sqrt{30}\zeta_1^2\zeta_4^2\zeta_2$$

$$z_{25} = \sqrt{30}\zeta_1^2\zeta_4^2\zeta_3$$

$$z_{26} = \sqrt{30}\zeta_2^2\zeta_4^2\zeta_1$$

$$z_{27} = \sqrt{60}\zeta_1^2\zeta_2\zeta_3\zeta_4$$

$$z_{28} = \sqrt{60}\zeta_2^2\zeta_1\zeta_3\zeta_4$$

$$w_{20} = -\sqrt{20}\zeta_4^3\zeta_1\zeta_2$$

$$w_{21} = \sqrt{30}\zeta_3^2\zeta_4^2\zeta_1$$

$$w_{22} = \sqrt{30}\zeta_3^2\zeta_4^2\zeta_2$$

$$w_{23} = -\sqrt{30}\zeta_3^2\zeta_1^2\zeta_4$$

$$w_{24} = -\sqrt{30}\zeta_3^2\zeta_2^2\zeta_4$$

$$w_{25} = \sqrt{30}\zeta_3^2\zeta_2^2\zeta_1$$

$$w_{26} = -\sqrt{30}\zeta_2^2\zeta_4^2\zeta_3$$

$$w_{27} = -\sqrt{60}\zeta_3^2\zeta_1\zeta_2\zeta_4$$

$$w_{28} = -\sqrt{60}\zeta_4^2\zeta_1\zeta_2\zeta_3$$

13

Additional Topics in Complex Geometry

Before turning to our main topics we first discuss partially hyperbolic diffeomorphisms and holomorphic Anosov flows as introduced by Étienne Ghys [1995]. In Section 13.2 we discuss the geometry of the projectivized holomorphic tangent and cotangent bundles. The study of the projectivized holomorphic tangent bundle naturally raises the question of a complex geodesic flow, which we discuss in Section 13.3. In Section 13.4 we return to the projectivized holomorphic tangent bundle and develop its complex almost contact metric structure. In Section 13.5 we first discuss special directions on complex contact manifolds analogous to our treatment in the real case in Chapter 11 and then discuss complex contact structures on the Lie group $SL(2, \mathbb{C})$ in detail.

13.1 Partial and holomorphic hyperbolicity

The purpose of this section is to present, as preliminaries, some ideas related to real and complex hyperbolicity. A diffeomorphism f of a (usually compact) Riemannian manifold M is said to be *partially hyperbolic in the narrow sense* (see, e.g., Pesin [2004, pp. 13–14]) if there exist numbers $C > 0$ and

$$0 < \lambda_1 \leq \mu_1 < \lambda_2 \leq \mu_2 < \lambda_3 \leq \mu_3, \quad \mu_1 < 1, \quad \lambda_3 > 1,$$

independent of $p \in M$, and an invariant splitting

$$TM = E^s \oplus E^c \oplus E^u$$

such that for $n > 0$,

$$C^{-1}\lambda_1^n\|v\| \leq \|f_*^n v\| \leq C\mu_1^n\|v\|, \quad v \in E_p^s,$$

$$C^{-1}\lambda_2^n\|v\| \leq \|f_*^n v\| \leq C\mu_2^n\|v\|, \quad v \in E_p^c,$$

$$C^{-1}\lambda_3^n\|v\| \leq \|f_*^n v\| \leq C\mu_3^n\|v\|, \quad v \in E_p^u.$$

As mentioned in Section 11.2 for Anosov flows, when M is compact the notion is independent of the choice of metric, but when M is noncompact, the notion is in general metric dependent. The subbundles E^s , E^c and E^u are called the *stable*, *central*, and *unstable subbundles* respectively.

For example, let ψ_t be an Anosov flow corresponding to a vector field ξ . For fixed t the diffeomorphism ψ_t is partially hyperbolic with 1-dimensional central direction generated by ξ . In Section 13.5 we will encounter a real vector field whose corresponding 1-parameter group is a group of partially hyperbolic diffeomorphisms for which the central subbundle has dimension 2.

In [1995] É. Ghys defines the notion of a holomorphic Anosov flow as a particular \mathbb{C}^* -action on a complex manifold that gives rise to an invariant splitting of the real tangent bundle, together with natural growth conditions on the subbundles. Lemma 2.1 of Ghys [1995] shows that the resulting stable and unstable subbundles extend to complex subbundles in the complexified tangent bundle. Also, in remarks on page 600 of his paper, Ghys discusses the possible consideration of starting with a holomorphic vector field and the flow it generates. This is described in terms of a splitting of the complexified tangent bundle, but one could equally well begin with a splitting of the holomorphic tangent bundle, and we take this point of view here.

Let ξ be a holomorphic vector field on a Hermitian manifold M ; strictly speaking, ξ does not determine a flow due to the lack of a natural ordering of the complex numbers. However, for a holomorphic vector field, the theory of complex differential equations goes through as in the real case. Let w_1, \dots, w_n be local complex coordinates on M and $f^j(w_1, \dots, w_n)$ the holomorphic component functions of ξ with respect to the basis $\{\frac{\partial}{\partial w_1}, \dots, \frac{\partial}{\partial w_n}\}$. Then the complex autonomous system

$$\frac{dw_\alpha}{dz} = f^\alpha(w_1, \dots, w_n), \quad \alpha = 1, \dots, n,$$

satisfies existence and uniqueness theorems similar to those in the real case (see, e.g., Hille [1976, Theorem 2.2.2]). Thus for a given point $w_0 \in M$ one can construct a unique local holomorphic curve through the point by the solution $w(z) = (w_1(z), \dots, w_n(z))$, $w(0) = w_0$. Therefore given ξ , one can define a “flow” ψ_z mapping a point w_0 to the point $w(z)$. We say that ξ (and the flow) is a *holomorphic Anosov flow* if there exists an invariant splitting of the holomorphic tangent bundle τ as a direct sum of type- $(1, 0)$ subbundles E^s , E^u and the 2-dimensional bundle tangent to the orbits of ψ_z together with numbers $C > 0$, $\lambda > 0$ such that

$$\begin{aligned}\|\psi_{z*}v\| &\leq Ce^{-\lambda\Re(z)}\|v\|, & v \in E_p^s, \\ \|\psi_{z*}v\| &\geq Ce^{\lambda\Re(z)}\|v\|, & v \in E_p^u,\end{aligned}$$

where \Re denotes the real part.

Before leaving this section we discuss briefly the question of integrability for a complex autonomous system of the form

$$\frac{\partial w^\alpha}{\partial z} = f^\alpha(w^1, \dots, w^n), \quad \alpha = 1, \dots, n,$$

where the functions on the right are complex-valued but not necessarily holomorphic. This is motivated by the consideration of a vector field, $\xi = \sum f^\alpha \frac{\partial}{\partial w^\alpha}$, of type $(1, 0)$ but not necessarily holomorphic. The question is when is there a local foliation by holomorphic curves whose tangent spaces are determined by the vector field. Being of type $(1, 0)$, the vector field is of the form $X - iJX$ for some real vector field X , and hence it determines a subbundle spanned by X and JX . Integrability of the subbundle requires that the Lie bracket $[X, JX]$ be a linear combination of X and JX , say $[X, JX] = aX + bJX$ for some real-valued functions a and b . Since $[\xi, \bar{\xi}] = 2i[X, JX]$, the integrability condition takes the form $[\xi, \bar{\xi}] = i(\alpha\xi + \bar{\alpha}\bar{\xi})$, $\alpha = a + ib$.

Proposition 13.1 *Consider the complex autonomous system*

$$\frac{\partial w^\alpha}{\partial z} = f^\alpha(w^1, \dots, w^n), \quad \alpha = 1, \dots, n,$$

where the complex-valued functions f^α are smooth and nonvanishing on a simply connected domain. Then the system is integrable if and only if the f^j ’s are of the form $f^\alpha = gF^\alpha$, where each F^α is holomorphic and g is complex-valued.

Proof. For the sufficiency suppose $f^\alpha = gF^\alpha$, where each F^α is holomorphic. Then

$$\begin{aligned} [\xi, \bar{\xi}] &= \left[\sum g F^\alpha \frac{\partial}{\partial w^\alpha}, \sum \bar{g} \bar{F}^\beta \frac{\partial}{\partial \bar{w}^\beta} \right] \\ &= \sum g F^\alpha \frac{\partial \bar{g}}{\partial w^\alpha} \bar{F}^\beta \frac{\partial}{\partial \bar{w}^\beta} - \sum \bar{g} \bar{F}^\alpha \frac{\partial g}{\partial \bar{w}^\alpha} F^\beta \frac{\partial}{\partial w^\beta} \\ &= -\left(\frac{\bar{g}}{g} \sum \bar{F}^\alpha \frac{\partial g}{\partial \bar{w}^\alpha} \right) \xi + \left(\frac{g}{\bar{g}} \sum F^\alpha \frac{\partial \bar{g}}{\partial w^\alpha} \right) \bar{\xi}, \end{aligned}$$

giving the integrability.

Conversely, given the integrability of the subbundle, we can view it as a foliation of a domain in \mathbb{C}^n by J -invariant submanifolds, but such submanifolds are complex submanifolds and therefore are given by holomorphic functions $w^j(z)$. The corresponding holomorphic tangent vector field, say $\sum F^\alpha \frac{\partial}{\partial w^\alpha}$, must then be proportional to the given vector field $\sum f^\alpha \frac{\partial}{\partial w^\alpha}$. ■

We can easily generalize this proposition slightly. Suppose that the hypotheses on the f^α 's hold for $\alpha = 1, \dots, p$ and that $f^\alpha \equiv 0$ for $\alpha > p$. Then for any solution, $w^\alpha = \text{const.}$ for $\alpha > p$, and the remaining p equations behave as in the proposition. For example, in Example 12.2.7 we have the nonholomorphic vector field

$$U_0 + iV_0 = \frac{4}{\sqrt{f_0}} \left(\frac{\partial}{\partial z^0} + \sum_{\alpha=1}^n \bar{w}^\alpha \frac{\partial}{\partial z^\alpha} \right),$$

where we caution that the coordinates in this example are $\{z^0, \dots, z^n, w^1, \dots, w^n\}$. For the corresponding autonomous system, the w^α 's become constants and the $n+1$ functions on the right-hand sides of the remaining equations are of the form $g = \frac{4}{\sqrt{f_0}}$ times the constant functions $F^\alpha = \bar{w}^\alpha$. The solutions represent the integral submanifolds of the vertical subbundle \mathcal{V} .

13.2 Projectivized holomorphic bundles

In Section 9.5 we gave an example of a contact structure on the projectivized tangent bundle of the hyperbolic plane, and in Example 3.2.4 we briefly mentioned the projectivized cotangent bundle. In the complex setting we first discuss the projectivized holomorphic cotangent bundle and

then study the projectivized holomorphic tangent bundle as a complex analogue of the tangent sphere bundle.

Let M be an $n + 1$ complex dimensional complex manifold with local coordinates (w^0, \dots, w^n) and τ^* its holomorphic cotangent bundle with fiber coordinates $(\hat{p}^0, \dots, \hat{p}^n)$. Each fiber of τ^* is a copy of \mathbb{C}^{n+1} , and one can consider the corresponding projective space \mathbb{CP}^n giving the projectivized holomorphic cotangent bundle $P\tau^*$.

If $\pi : \tau^* \longrightarrow M$ is the projection map, then w^α and \hat{p}^α are local coordinates on τ^* , where we have identified w^α with $w^\alpha \circ \pi$. Consider the Liouville form β , which is given locally by $\beta = \sum_{\alpha=0}^n \hat{p}^\alpha dw^\alpha$. Now on $P\tau^*$ the \hat{p}^α 's are homogeneous coordinates for the fibers, and we introduce nonhomogeneous coordinates (p^1, \dots, p^n) on the neighborhood defined by $\hat{p}^0 \neq 0$ by $p^\alpha = \frac{\hat{p}^\alpha}{\hat{p}^0}$. Then

$$\theta = dw^0 + \sum_{i=1}^n p^\alpha dq^\alpha$$

is a local complex contact form, and taking charts on $P\tau^*$ defined by $\hat{p}^\alpha \neq 0$, we obtain a complex contact structure.

Turning to the Hermitian setting, we first note that one case of the projectivized holomorphic tangent bundle has already been discussed, namely $\mathbb{C}^{n+1} \times \mathbb{CP}^n$ (16) in Example 12.2.7. More generally, let M be an $n + 1$ complex dimensional Hermitian manifold with local coordinates $\{w^0, \dots, w^n\}$ and $\hat{\pi} : \tau \longrightarrow M$ its holomorphic tangent bundle with fiber coordinates $\{\hat{\zeta}^0, \dots, \hat{\zeta}^n\}$. Each fiber is \mathbb{C}^{n+1} , and we consider the corresponding complex projective space \mathbb{CP}^n . We call the bundle constructed in this way the *projectivized holomorphic tangent bundle* $P\tau$. The $\hat{\zeta}^\alpha$'s are homogeneous coordinates for the fibers of $P\tau$, and we introduce nonhomogeneous coordinates $\{\zeta^1, \dots, \zeta^n\}$ on the neighborhood defined by $\hat{\zeta}^0 \neq 0$ by $\zeta^\alpha = \hat{\zeta}^\alpha / \hat{\zeta}^0$. Let $\pi : P\tau \longrightarrow M$ denote the projection, and we denote by $\{w^0, \dots, w^n, \zeta^1, \dots, \zeta^n\}$ the local coordinates on $P\tau$, where we have identified w^α with $w^\alpha \circ \pi$.

The vertical coordinate fields on τ project to $P\tau$ in the following manner:

$$\frac{\partial}{\partial \hat{\zeta}^0} \longrightarrow -\frac{1}{\hat{\zeta}^0} \sum_{\gamma=1}^n \zeta^\gamma \frac{\partial}{\partial \zeta^\gamma}, \quad \frac{\partial}{\partial \hat{\zeta}^\alpha} \longrightarrow \frac{1}{\hat{\zeta}^0} \frac{\partial}{\partial \zeta^\alpha} \quad \alpha = 1, \dots, n.$$

Let $\mathbf{G}_{\alpha\bar{\beta}}$ denote the Hermitian metric on M and $\mu_{\beta\gamma}^\alpha$ the connection coefficients. The metric gives rise to horizontal lifts of vector fields to τ

by

$$\left(\frac{\partial}{\partial w^\beta}\right)^H = \frac{\partial}{\partial w^\beta} - \hat{\zeta}^\gamma \mu_{\beta\gamma}^\alpha \frac{\partial}{\partial \hat{\zeta}^\alpha}.$$

The vertical lift of $\frac{\partial}{\partial w^\beta}$ is $(\frac{\partial}{\partial w^\beta})^V = \frac{\partial}{\partial \hat{\zeta}^\beta}$.

In the fibers of τ consider the spheres $\sum_{\alpha,\gamma=0}^n \mathbf{G}_{\alpha\bar{\gamma}} \hat{\zeta}^\alpha \bar{\hat{\zeta}}^\gamma = 1$ and set $P_\alpha = \sum_{\gamma=0}^n \mathbf{G}_{\alpha\bar{\gamma}} \bar{\hat{\zeta}}^\gamma$. Now instead of studying the vertical lift of a vector field to τ , we define the *projective lift* to τ by

$$\left(\frac{\partial}{\partial w^\alpha}\right)^P = \hat{\zeta}^0 \left(\frac{\partial}{\partial \hat{\zeta}^\alpha} - P_\alpha \sum_{\gamma=0}^n \hat{\zeta}^\gamma \frac{\partial}{\partial \hat{\zeta}^\gamma} \right).$$

Projecting to $P\tau$, we have

$$\left(\frac{\partial}{\partial w^0}\right)^P \longrightarrow - \sum_{\gamma=1}^n \zeta^\gamma \frac{\partial}{\partial \zeta^\gamma}, \quad \left(\frac{\partial}{\partial w^\alpha}\right)^P \longrightarrow \frac{\partial}{\partial \zeta^\alpha}, \quad a = 1, \dots, n.$$

We also remark that dividing $\sum_{\alpha,\gamma=0}^n \mathbf{G}_{\alpha\bar{\gamma}} \hat{\zeta}^\alpha \bar{\hat{\zeta}}^\gamma = 1$ by $|\hat{\zeta}^0|^2$ gives

$$\mathbf{G}_{0\bar{0}} + \sum_{\gamma=1}^n \mathbf{G}_{0\bar{\gamma}} \bar{\zeta}^\gamma + \sum_{\alpha=1}^n \mathbf{G}_{\alpha\bar{0}} \zeta^\alpha + \sum_{\alpha,\gamma=1}^n \mathbf{G}_{\alpha\bar{\gamma}} \zeta^\alpha \bar{\zeta}^\gamma = \frac{1}{|\hat{\zeta}^0|^2}.$$

The left-hand side is now a local function on $P\tau$, and hence $\frac{1}{|\hat{\zeta}^0|^2}$, or just $|\hat{\zeta}^0|$, becomes an important local function on $P\tau$.

The holomorphic tangent bundle carries a Hermitian metric, similar to the Sasaki metric of Section 9.1 on the tangent bundle of a Riemannian manifold (see, e.g., Munteanu [2004, p. 50]). Denoting this metric by \hat{g} , it is given in terms of vectors X, Y of type $(1, 0)$ on M by

$$\hat{g}(X^H, \bar{Y}^H) = \hat{g}(X^V, \bar{Y}^V) = \mathbf{G}(X, \bar{Y}), \quad \hat{g}(X^H, \bar{Y}^V) = 0.$$

We now define a Hermitian metric g on $P\tau$ by

$$\begin{aligned} g(X^H, \bar{Y}^H) &= \hat{g}(X^H, \bar{Y}^H) = \mathbf{G}(X, \bar{Y}), \\ g(X^H, \bar{Y}^P) &= 0, \quad g(X^P, \bar{Y}^P) = \hat{g}(X^P, \bar{Y}^P). \end{aligned}$$

13.3 The complex geodesic flow

Let us first recall the characterization of complex space forms of Yano and Mogi (Theorem 8.4) that a Kähler manifold M^{2n} , $n \geq 2$, is a complex space form if and only if for every point and every holomorphic section at the point, there exists a unique totally geodesic holomorphic curve through the point and tangent to the given holomorphic section at the point. Thus for complex space forms one has a natural complex analogue of a geodesic.

Now consider the vector field on the holomorphic tangent bundle τ defined by

$$\sum_{\alpha=0}^n \hat{\zeta}^\alpha \left(\frac{\partial}{\partial w^\alpha} \right)^H.$$

For our discussion of the projectivized holomorphic tangent bundle $P\tau$, first divide the above vector field by $\hat{\zeta}^0$ and consider the vector field

$$\Xi = \left(\frac{\partial}{\partial w^0} \right)^H + \sum_{\alpha=1}^n \zeta^\alpha \left(\frac{\partial}{\partial w^\alpha} \right)^H.$$

We will do this explicitly for complex hyperbolic space $\mathbb{C}H^{n+1}$ and complex projective space $\mathbb{C}P^{n+1}$. The case of \mathbb{C}^{n+1} is trivial.

For complex hyperbolic space, the Bergman metric of constant holomorphic curvature -1 is given by

$$ds^2 = 4 \frac{(1 - \sum |w^\varepsilon|^2)(\sum dw^\varepsilon d\bar{w}^\varepsilon) + (\sum \bar{w}^\varepsilon dw^\varepsilon)(\sum w^\varepsilon d\bar{w}^\varepsilon)}{(1 - \sum |w^\varepsilon|^2)^2}.$$

The connection coefficients are

$$\mu_{\beta\beta}^\beta = \frac{2\bar{w}^\beta}{1 - \sum |w^\varepsilon|^2}, \quad \mu_{\gamma\beta}^\gamma = \frac{\bar{w}^\beta}{1 - \sum |w^\varepsilon|^2}, \quad \mu_{\beta\beta}^\alpha = 0, \quad \mu_{\beta\gamma}^\alpha = 0.$$

For complex projective space $\mathbb{C}P^{n+1}$, we have the Fubini–Study metric of constant holomorphic curvature $+1$:

$$ds^2 = 4 \frac{(1 + \sum |w^\varepsilon|^2)(\sum dw^\varepsilon d\bar{w}^\varepsilon) - (\sum \bar{w}^\varepsilon dw^\varepsilon)(\sum w^\varepsilon d\bar{w}^\varepsilon)}{(1 + \sum |w^\varepsilon|^2)^2}.$$

The connection coefficients are

$$\mu_{\beta\beta}^\beta = \frac{-2\bar{w}^\beta}{1 + \sum |w^\varepsilon|^2}, \quad \mu_{\gamma\beta}^\gamma = \frac{-\bar{w}^\beta}{1 + \sum |w^\varepsilon|^2}, \quad \mu_{\beta\beta}^\alpha = 0, \quad \mu_{\beta\gamma}^\alpha = 0.$$

An important observation is that in both cases $\mu_{\beta\beta}^\beta = 2\mu_{\gamma\beta}^\gamma$. In fact, a Kähler manifold of complex dimension ≥ 2 with local coordinates $\{w^\alpha\}$ and satisfying $\mu_{\beta\beta}^\beta = 2\mu_{\gamma\beta}^\gamma$, $\mu_{\beta\beta}^\alpha = 0$, $\mu_{\beta\gamma}^\alpha = 0$ is of constant holomorphic curvature. This fact may be of some independent interest, and a proof is given in the author's paper [2007].

We now use the projections of the vertical coordinate fields from τ to $P\tau$ and the expression for horizontal lifts from the previous section. The following computation is straightforward, though somewhat delicate, but it relies only on the properties $\mu_{\beta\beta}^\beta = 2\mu_{\gamma\beta}^\gamma$, $\mu_{\beta\beta}^\alpha = 0$, $\mu_{\beta\gamma}^\alpha = 0$ of the connection coefficients and is therefore the same for both $\mathbb{C}H^{n+1}$ and $\mathbb{C}P^{n+1}$:

$$\begin{aligned}
& \left(\frac{\partial}{\partial w^0} \right)^H + \sum_{\beta=1}^n \zeta^\beta \left(\frac{\partial}{\partial w^\beta} \right)^H \longrightarrow \\
& \frac{\partial}{\partial w^0} + \mu_{00}^0 \sum_{\gamma=1}^n \zeta^\gamma \frac{\partial}{\partial \zeta^\gamma} - \sum_{\gamma=1}^n \mu_{0\gamma}^\gamma \zeta^\gamma \frac{\partial}{\partial \zeta^\gamma} + \sum_{\gamma,\delta=1}^n \mu_{0\gamma}^0 \zeta^\gamma \zeta^\delta \frac{\partial}{\partial \zeta^\delta} \\
& + \sum_{\beta=1}^n \zeta^\beta \left\{ \frac{\partial}{\partial w^\beta} - \mu_{\beta\beta}^\beta \zeta^\beta \frac{\partial}{\partial \zeta^\beta} - \sum_{\gamma \neq \beta, 0} \mu_{\beta\gamma}^\gamma \zeta^\gamma \frac{\partial}{\partial \zeta^\gamma} \right. \\
& \quad \left. + \mu_{\beta 0}^0 \sum_{\gamma=1}^n \zeta^\gamma \frac{\partial}{\partial \zeta^\gamma} - \sum_{\gamma \neq \beta, 0} \mu_{\beta\gamma}^\beta \zeta^\gamma \frac{\partial}{\partial \zeta^\beta} - \mu_{\beta 0}^\beta \frac{\partial}{\partial \zeta^\beta} \right\} \\
& = \frac{\partial}{\partial w^0} + \frac{1}{2} \mu_{00}^0 \sum_{\gamma=1}^n \zeta^\gamma \frac{\partial}{\partial \zeta^\gamma} + \frac{1}{2} \sum_{\gamma,\delta=1}^n \mu_{\gamma\delta}^\gamma \zeta^\gamma \zeta^\delta \frac{\partial}{\partial \zeta^\delta} \\
& + \sum_{\beta=1}^n \zeta^\beta \frac{\partial}{\partial w^\beta} - \sum_{\beta=1}^n \mu_{\beta\beta}^\beta (\zeta^\beta)^2 \frac{\partial}{\partial \zeta^\beta} - \frac{1}{2} \sum_{\beta=1}^n \sum_{\gamma \neq \beta, 0} \mu_{\beta\gamma}^\beta \zeta^\beta \zeta^\gamma \frac{\partial}{\partial \zeta^\gamma} \\
& + \frac{1}{2} \sum_{\beta,\gamma=1}^n \mu_{\beta\gamma}^\beta \zeta^\beta \zeta^\gamma \frac{\partial}{\partial \zeta^\gamma} - \frac{1}{2} \sum_{\beta=1}^n \sum_{\gamma \neq \beta, 0} \mu_{\beta\gamma}^\gamma \zeta^\beta \zeta^\gamma \frac{\partial}{\partial \zeta^\beta} \\
& - \frac{1}{2} \mu_{00}^0 \sum_{\beta=1}^n \zeta^\beta \frac{\partial}{\partial \zeta^\beta} \\
& = \frac{\partial}{\partial w^0} + \sum_{\beta=1}^n \zeta^\beta \frac{\partial}{\partial w^\beta}.
\end{aligned}$$

For this vector field consider the holomorphic autonomous system

$$\frac{dw^0}{dz} = 1, \quad \frac{dw^\alpha}{dz} = \zeta^\alpha, \quad \frac{d\zeta^\alpha}{dz} = 0.$$

Solving, we have a map ψ_z that maps a point $(w_0^0, w_0^1, \dots, w_0^n, \zeta_0^1, \dots, \zeta_0^n)$ to the point $(w_0^0 + z, w_0^1 + \zeta_0^1 z, \dots, w_0^n + \zeta_0^n z, \zeta_0^1, \dots, \zeta_0^n)$, and the orbits of ψ_z are lifts of the complex geodesics. We call this flow, and the vector field, the *complex geodesic flow*, and we have the following theorem.

Theorem 13.1 *On the projectivized holomorphic tangent bundle of a Kähler manifold of constant holomorphic curvature, the vector field*

$$\Xi = \left(\frac{\partial}{\partial w^0} \right)^H + \sum_{\beta=1}^n \zeta^\beta \left(\frac{\partial}{\partial w^\beta} \right)^H$$

generates a flow whose orbits are horizontal lifts of the complex geodesics of the Kähler manifold.

Before studying this complex geodesic flow further for the cases of the complex hyperbolic plane and complex projective plane, we will prove a converse of Theorem 13.1 for Kähler manifolds. That is, when the base manifold is Kähler we take up the question of the integrability of the subbundle determined by the vector field Ξ .

First recall that when the base manifold M is Kähler, the connection coefficients take the form

$$\mu_{\beta\gamma}^\alpha = \mathbf{G}^{\bar{\delta}\alpha} \frac{\partial \mathbf{G}_{\gamma\bar{\delta}}}{\partial w^\beta},$$

and the curvature of M is given by

$$\frac{\partial \mu_{\alpha\gamma}^\sigma}{\partial \bar{w}^\beta} = -\mathbf{R}_{\alpha\bar{\beta}\gamma\bar{\delta}} \mathbf{G}^{\bar{\delta}\sigma}.$$

Thus the Lie brackets of horizontal lifts of the coordinate fields become

$$\begin{aligned} \left[\left(\frac{\partial}{\partial w^\beta} \right)^H, \left(\frac{\partial}{\partial \bar{w}^\delta} \right)^H \right] &= \hat{\zeta}^\gamma \frac{\partial \mu_{\beta\gamma}^\alpha}{\partial \bar{w}^\delta} \frac{\partial}{\partial \hat{\zeta}^\alpha} - \bar{\hat{\zeta}}^\varepsilon \frac{\partial \bar{\mu}_{\delta\varepsilon}^\alpha}{\partial w^\beta} \frac{\partial}{\partial \hat{\bar{\zeta}}^\alpha} \\ &= -\hat{\zeta}^\gamma \mathbf{R}_{\beta\bar{\delta}\gamma\bar{\varepsilon}} \mathbf{G}^{\bar{\varepsilon}\alpha} \frac{\partial}{\partial \hat{\zeta}^\alpha} + \bar{\hat{\zeta}}^\varepsilon \mathbf{R}_{\beta\bar{\delta}\gamma\bar{\varepsilon}} \mathbf{G}^{\gamma\bar{\alpha}} \frac{\partial}{\partial \hat{\bar{\zeta}}^\alpha}. \end{aligned}$$

To find the integrability condition of the subbundle determined by Ξ we write Ξ as

$$\Xi = \frac{1}{\hat{\zeta}^0} \sum_{\delta=0}^n \hat{\zeta}^\delta \left(\frac{\partial}{\partial w^\delta} \right)^H$$

and compute the Lie bracket $[\Xi, \bar{\Xi}]$. The horizontal part vanishes and the vertical part is imaginary, namely

$$-\frac{2}{|\hat{\zeta}^0|^2} \Im \left(\hat{\zeta}^\delta \overline{\hat{\zeta}^\varepsilon} \hat{\zeta}^\gamma \mathbf{R}_{\delta\bar{\varepsilon}\gamma\bar{\beta}} \mathbf{G}^{\bar{\beta}\alpha} \frac{\partial}{\partial \hat{\zeta}^\alpha} \right),$$

where we have used the symmetry $\overline{\mathbf{R}_{\alpha\bar{\beta}\gamma\bar{\delta}}} = \mathbf{R}_{\beta\bar{\alpha}\delta\bar{\gamma}}$. Therefore the subbundle will be integrable when the inner product of this with $\frac{\partial}{\partial \zeta^\sigma}$ vanishes. Computing this on the bundle τ in terms of the metric \hat{g} and relabeling, we have

$$\hat{g} \left(\left(\frac{\partial}{\partial w^\sigma} \right)^P, \hat{\zeta}^\delta \overline{\hat{\zeta}^\varepsilon} \hat{\zeta}^\beta \mathbf{R}_{\delta\bar{\varepsilon}\gamma\bar{\beta}} \mathbf{G}^{\gamma\bar{\alpha}} \left(\frac{\partial}{\partial w^\alpha} \right)^P \right) = |\hat{\zeta}^0|^2 \hat{\zeta}^\delta \overline{\hat{\zeta}^\varepsilon} \hat{\zeta}^\beta (\mathbf{R}_{\delta\bar{\varepsilon}\sigma\bar{\beta}} - P_\sigma \mathbf{R}_{\delta\bar{\varepsilon}\gamma\bar{\beta}} \hat{\zeta}^\gamma).$$

Now if this vanishes, we interpret it in terms of $(1, 0)$ vectors Z and \bar{W} , giving

$$\mathbf{G}(\mathbf{R}_{Z\bar{Z}} \bar{W}, \bar{Z}) = \mathbf{G}(\bar{W}, \bar{Z}) \mathbf{G}(\mathbf{R}_{Z\bar{Z}} Z, \bar{Z}).$$

Taking conjugates, we have

$$\mathbf{G}(\mathbf{R}_{Z\bar{Z}} Z, \bar{W}) = \lambda \mathbf{G}(Z, \bar{W}),$$

where $\lambda = \mathbf{G}(\mathbf{R}_{Z\bar{Z}} Z, \bar{Z})$. Writing Z as $X - iJX$ and taking the $(1, 0)$ part, we have

$$\mathbf{R}_{(X-iJX)(X+iJX)}(X - iJX) = \lambda(X - iJX),$$

since the left-hand side expands to $2i(\mathbf{R}_{XJX}X - iJ\mathbf{R}_{XJX}X)$. From this we see that the integrability condition of the subbundle is that $\mathbf{R}_{XJX}X$ be proportional to JX , a well-known characterization of constant holomorphic curvature (Kosmanek [1964] for Kähler manifolds and generalized by Tanno [1973] to almost Hermitian manifolds satisfying $\mathbf{G}(\mathbf{R}_{JXJY}JX, JZ) = \mathbf{G}(\mathbf{R}_{XY}X, Z)$). Combining with Theorem 13.1 we have the following theorem.

Theorem 13.2 *On the projectivized holomorphic tangent bundle of a Kähler manifold M , the vector field $\Xi = \left(\frac{\partial}{\partial w^0}\right)^H + \sum_{\beta=1}^n \zeta^\beta \left(\frac{\partial}{\partial w^\beta}\right)^H$ determines an integrable subbundle if and only if M is of constant holomorphic curvature.*

To compare the vector field here with the corresponding one in the real case we refer the reader to Sections 9.2 and 9.5.

For the complex space forms the differential of the complex geodesic flow, ψ_z , is easily computed and for $n = 2$ is simply the matrix

$$\begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & z \\ 0 & 0 & 1 \end{pmatrix}.$$

We now discuss in greater detail the geometry of the complex geodesic flow for the complex hyperbolic plane and the complex projective plane. In particular, we will first analyze the flow ψ_z for the vector field $\frac{\partial}{\partial w^0} + \zeta \frac{\partial}{\partial w^1}$ on the bundle $P\tau$ over $\mathbb{C}H^2$; since $n + 1 = 2$, we denote the fiber coordinate ζ^1 simply by ζ , but for the index on a tensor we use 1^* . The metric g of the previous section on $P\tau$ is given with respect to the coordinates $\{w^0, w^1, \zeta\}$ by

$$\begin{aligned} g_{00} &= \frac{4(1 - |w^1|^2)}{(1 - |w^0|^2 - |w^1|^2)^2} + \frac{|\zeta|^2 |\bar{w}^0 + \zeta \bar{w}^1|^2}{(1 - |w^0|^2 - |w^1|^2)(1 + |\zeta|^2 - |\zeta w^0 - w^1|^2)^2}, \\ g_{0\bar{1}} &= \frac{4\bar{w}^0 w^1}{(1 - |w^0|^2 - |w^1|^2)^2} - \frac{\zeta |\bar{w}^0 + \zeta \bar{w}^1|^2}{(1 - |w^0|^2 - |w^1|^2)(1 + |\zeta|^2 - |\zeta w^0 - w^1|^2)^2}, \\ g_{1\bar{1}} &= \frac{4(1 - |w^0|^2)}{(1 - |w^0|^2 - |w^1|^2)^2} + \frac{|\bar{w}^0 + \zeta \bar{w}^1|^2}{(1 - |w^0|^2 - |w^1|^2)(1 + |\zeta|^2 - |\zeta w^0 - w^1|^2)^2}, \\ g_{0\bar{1}^*} &= -\frac{\zeta(\bar{w}^0 + \zeta \bar{w}^1)}{(1 + |\zeta|^2 - |\zeta w^0 - w^1|^2)^2}, \\ g_{1\bar{1}^*} &= \frac{(\bar{w}^0 + \zeta \bar{w}^1)}{(1 + |\zeta|^2 - |\zeta w^0 - w^1|^2)^2}, \quad g_{1^* \bar{1}^*} = \frac{1 - |w^0|^2 - |w^1|^2}{(1 + |\zeta|^2 - |\zeta w^0 - w^1|^2)^2}. \end{aligned}$$

Now consider the vector fields

$$V^\pm = \begin{pmatrix} 0 \\ w^0 + \bar{\zeta} w^1 \pm \sqrt{1 + |\zeta|^2 - |\zeta w^0 - w^1|^2} \\ 1 + |\zeta|^2 \end{pmatrix},$$

the entries being the components with respect to the basis $\{\frac{\partial}{\partial w^0}, \frac{\partial}{\partial w^1}, \frac{\partial}{\partial \zeta}\}$. Notice that $\zeta w^0 - w^1$ is constant along the orbits of the flow ψ_z . Apply

the differential of the flow to the values of V^\pm at (w_0^0, w_0^1, ζ_0) :

$$\begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & z \\ 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} 0 \\ w_0^0 + \bar{\zeta}_0 w_0^1 \pm \sqrt{1 + |\zeta_0|^2 - |\zeta_0 w_0^0 - w_0^1|^2} \\ 1 + |\zeta_0|^2 \end{pmatrix} \\ = \begin{pmatrix} 0 \\ w_0^0 + \bar{\zeta}_0 w_0^1 + (1 + |\zeta_0|^2)z \pm \sqrt{1 + |\zeta_0|^2 - |\zeta_0 w_0^0 - w_0^1|^2} \\ 1 + |\zeta_0|^2 \end{pmatrix}$$

but this is the value of V^\pm at the image point $(w_0^0 + z, w_0^1 + \zeta_0 z, \zeta_0)$. Therefore V^\pm determine invariant subbundles transverse to orbits of the flow.

Next we compute the square of the length of V^\pm and obtain

$$g(V^\pm, \bar{V}^\pm) = \frac{|w^0 + \bar{\zeta} w^1 \pm \sqrt{1 + |\zeta|^2 - |\zeta w^0 - w^1|^2}|^2}{(1 - |w^0|^2 - |w^1|^2)^2 (1 + |\zeta|^2 - |\zeta w^0 - w^1|^2)} \\ \times (4(1 - |w^0|^2)(1 + |\zeta|^2 - |\zeta w^0 - w^1|^2) + 1 - |w^0|^2 - |w^1|^2).$$

The idea is to compare this at the image point with its value at the initial point; not surprisingly, this is quite complicated. Also the map ψ_z is not an isometry. However, consider the isometry of $\mathbb{C}H^2$ that maps the disk $\zeta w^0 - w^1 = \zeta w_0^0 - w_0^1$, fixed ζ , to the disk $w^1 = 0$ ($\zeta = 0$) and the point (w_0^0, w_0^1) to the origin. Now w^0 is just z on the unit disk. Moreover, the function $w^0 + \bar{\zeta} w^1 \pm \sqrt{1 + |\zeta|^2 - |\zeta w^0 - w^1|^2}$, which is the second component of V^\pm , is mapped to the function $z \pm 1$, which is the value of the second component of V^\pm along $(w^0, 0, 0)$. Now along $(w^0, 0, 0)$ the square of the length of V^\pm as given above is just

$$\frac{|z \pm 1|^2}{1 - |z|^2} \times 5,$$

and we study this function on the unit disk. The function tends to infinity as $|z| \rightarrow 1$ along all radii except the negative real axis in the plus case and the positive real axis in the minus case; in these cases the limit is 0. Restricted to the real axis this is hyperbolic behavior, with the plus case corresponding to the unstable bundle and the minus case to the stable bundle relative to increasing real parameter. Moreover, returning to the complex disk, choose a number R such that $0 < R < 1$. Let $C = 1 - R$ and α such that $R^\alpha = \frac{(1-R)^2}{1+R}$. Then on $|z| \leq R$, we have

$$C|z|^\alpha 5 \leq \frac{|z \pm 1|^2}{1 - |z|^2} 5 \leq \frac{C}{|z|^\alpha} 5$$

in the spirit of Ghys's condition for a holomorphic Anosov flow.

Now in contrast consider $\mathbb{C}P^2$. The metric g on $P\tau$ with respect to the coordinates $\{w^0, w^1, \zeta\}$ is given by

$$\begin{aligned} g_{0\bar{0}} &= \frac{4(1+|w^1|^2)}{(1+|w^0|^2+|w^1|^2)^2} + \frac{|\zeta|^2|\bar{w}^0+\zeta\bar{w}^1|^2}{(1+|w^0|^2+|w^1|^2)(1+|\zeta|^2+|\zeta w^0-w^1|^2)^2}, \\ g_{0\bar{1}} &= -\frac{4\bar{w}^0w^1}{(1+|w^0|^2+|w^1|^2)^2} - \frac{\zeta|\bar{w}^0+\zeta\bar{w}^1|^2}{(1+|w^0|^2+|w^1|^2)(1+|\zeta|^2+|\zeta w^0-w^1|^2)^2}, \\ g_{1\bar{1}} &= \frac{4(1+|w^0|^2)}{(1+|w^0|^2+|w^1|^2)^2} + \frac{|\bar{w}^0+\zeta\bar{w}^1|^2}{(1+|w^0|^2+|w^1|^2)(1+|\zeta|^2+|\zeta w^0-w^1|^2)^2}, \\ g_{0\bar{1}^*} &= \frac{\zeta(\bar{w}^0+\zeta\bar{w}^1)}{(1+|\zeta|^2+|\zeta w^0-w^1|^2)^2}, \\ g_{1\bar{1}^*} &= -\frac{(\bar{w}^0+\zeta\bar{w}^1)}{(1+|\zeta|^2+|\zeta w^0-w^1|^2)^2}, \\ g_{1^*\bar{1}^*} &= \frac{1+|w^0|^2+|w^1|^2}{(1+|\zeta|^2+|\zeta w^0-w^1|^2)^2}. \end{aligned}$$

Now suppose the vector field $V = a\frac{\partial}{\partial w^0} + b\frac{\partial}{\partial w^1} + \frac{\partial}{\partial \zeta}$ is invariant under ψ_{z*} and compute the square of its length along $(w^0, 0, 0)$:

$$g(V, \bar{V}) = |a|^2 \frac{4}{(1+|w^0|^2)^2} + |b|^2 \frac{4+|w^0|^2}{1+|w^0|^2} - b\bar{w}^0 - \bar{b}w^0 + (1+|w^0|^2).$$

Taking the initial point to be $(0, 0, 0)$ and $V = a_0\frac{\partial}{\partial w^0} + b_0\frac{\partial}{\partial w^1} + \frac{\partial}{\partial \zeta}$ at this point we have at $z (= w^0)$,

$$g(V, \bar{V}) = \frac{4|a_0|^2}{(1+|z|^2)^2} + \frac{3|z+b_0|^2}{1+|z|^2} + 1 + |b_0|^2,$$

which has limit $4+|b_0|^2$ as $|z| \rightarrow \infty$. On the other hand, if either $|V(z)| \leq \frac{1}{|z|^\alpha}|V(0)|$ or $|V(z)| \geq |z|^\alpha|V(0)|$, then the limit of $g(V, \bar{V})$ as $|z| \rightarrow \infty$ is either 0 or ∞ . Thus for $\mathbb{C}P^2$, ψ_{z*} on $P\tau$ does not admit holomorphic hyperbolic behavior. Thus, as with the classical geodesic flow, we see some hyperbolic behavior in the case of the negatively curved space $\mathbb{C}H^2$, but not in the positively curved space $\mathbb{C}P^2$.

13.4 Complex almost contact metric structure on $P\tau$

One of the difficulties of Hermitian geometry is the lack of holomorphicity in the metric. In fact, the only time the components of a Hermitian metric are holomorphic functions is when they are constants, and hence the metric is already flat, a much too restrictive situation. Thus, instead of seeking a complex contact metric structure on $P\tau$, which insists on the holomorphicity of the local contact forms, we relax this to require only that the local forms be of type $(1, 0)$. Another difficulty of Hermitian geometry is that using the metric to raise or lower indices reverses type (e.g., lowering the index on a $(1, 0)$ vector field gives a $(0, 1)$ -form). However this difficulty is easily overcome by taking conjugates. Using the metric we define a form by looking at the inner products $g(\frac{\partial}{\partial w^\alpha}, \bar{\Xi})$ giving a $(1, 0)$ -form θ ,

$$\theta = \frac{1}{(\hat{\zeta}^0)} \sum_{\alpha=0}^n P_\alpha dw^\alpha,$$

or equivalently

$$\theta = \sum_{\alpha=0}^n (\mathbf{G}_{\alpha\bar{0}} + p_\alpha) dw^\alpha, \quad p_\alpha = \sum_{\gamma=1}^n \mathbf{G}_{\alpha\bar{\gamma}} \bar{\zeta}^\gamma.$$

We remark that in the overlap with the coordinate patch defined by $\hat{\zeta}^1 \neq 0$ one has $\theta_1 = (\hat{\zeta}^0/\hat{\zeta}^1)\theta_0$, where we have used subscripts to identify the forms on the particular coordinate patches.

Staying within Kähler geometry, we can ask whether the local $(1, 0)$ -forms θ are contact forms in the sense that $\theta \wedge (d\theta)^n \neq 0$, even though they are not necessarily holomorphic forms giving a complex contact structure. Recalling that $\partial \mathbf{G}_{\alpha\bar{\gamma}}/\partial w^\beta$ is symmetric in α and β on a Kähler manifold, the differential of θ is a form of type $(1, 1)$ and given by

$$d\theta = -\left(\frac{\partial \mathbf{G}_{\alpha\bar{0}}}{\partial \bar{w}^\beta} + \sum_{\gamma=1}^n \frac{\partial \mathbf{G}_{\alpha\bar{\gamma}}}{\partial \bar{w}^\beta} \bar{\zeta}^\gamma\right) dw^\alpha \wedge d\bar{w}^\beta - \sum_{\gamma=1}^n \mathbf{G}_{\alpha\bar{\gamma}} dw^\alpha \wedge d\bar{\zeta}^\gamma.$$

To show that $\theta \wedge (d\theta)^n \neq 0$, it is enough to evaluate it on the coordinate vector fields

$$\left\{ \frac{\partial}{\partial w^0}, \frac{\partial}{\partial w^1}, \dots, \frac{\partial}{\partial w^n}, \frac{\partial}{\partial \bar{\zeta}^1}, \dots, \frac{\partial}{\partial \bar{\zeta}^n} \right\}.$$

Let \bigoplus denote the sum on the signed permutations of $\kappa_0, \dots, \kappa_n$. It is sufficient to look at sums of the form

$$\begin{aligned} & \bigoplus \theta \left(\frac{\partial}{\partial w^{\kappa_0}} \right) d\theta \left(\frac{\partial}{\partial w^{\kappa_1}}, \frac{\partial}{\partial \bar{\zeta}^1} \right) \cdots d\theta \left(\frac{\partial}{\partial w^{\kappa_n}}, \frac{\partial}{\partial \bar{\zeta}^n} \right) \\ &= \bigoplus \frac{1}{(\hat{\zeta}^0)} P_{\kappa_0} \left(\frac{-1}{2} \right)^n \mathbf{G}_{\kappa_1 \bar{1}} \cdots \mathbf{G}_{\kappa_n \bar{n}} \\ &= \left(\frac{-1}{2} \right)^n \frac{1}{(\hat{\zeta}^0)} \bigoplus \left(\sum_{\gamma=0}^n \mathbf{G}_{\kappa_0 \bar{\gamma}} \overline{\hat{\zeta}^\gamma} \right) \mathbf{G}_{\kappa_1 \bar{1}} \cdots \mathbf{G}_{\kappa_n \bar{n}}. \end{aligned}$$

The terms in the \sum -sum for $\gamma = 1, \dots, n$ will give zero upon doing the \bigoplus -sum. Therefore we have

$$\left(\frac{-1}{2} \right)^n \bigoplus \mathbf{G}_{\kappa_0 \bar{0}} \mathbf{G}_{\kappa_1 \bar{1}} \cdots \mathbf{G}_{\kappa_n \bar{n}} = \left(\frac{-1}{2} \right)^n \det(\mathbf{G}_{\kappa \bar{\gamma}}) \neq 0.$$

Therefore $\theta \wedge (d\theta)^n \neq 0$, giving us a local contact form of type $(1, 0)$.

Returning to the general setting, we show that even though we cannot expect a complex contact metric structure on the projectivized holomorphic tangent bundle $P\tau$, it does carry a complex almost contact metric structure.

Define local real 1-forms u and v by

$$u - iv = \frac{|\hat{\zeta}^0|}{2} \theta$$

and vector fields U and V by

$$U + iV = 4|\hat{\zeta}^0| \Xi.$$

Since both Ξ and θ are of type $(1, 0)$, we have that $V = -JU$ and $v = u \circ J$. By an easy computation we see that $\theta(\Xi) = \frac{1}{|\hat{\zeta}^0|^2}$ and then

$2u(U + iV) = \frac{|\hat{\zeta}^0|}{2} (\theta + \bar{\theta}) (4|\hat{\zeta}^0| \Xi) = 2$, giving $u(U) = 1$ and $u(V) = 0$. Similarly $v(V) = 1$ and $v(U) = 0$.

Before defining the fields of endomorphisms G and H , we introduce the following vector fields:

$$E_\alpha = \hat{\zeta}^0 \left(\left(\frac{\partial}{\partial w^\alpha} \right)^H - P_\alpha \hat{\zeta}^0 \Xi \right), \quad \alpha = 1, \dots, n.$$

Then one can check directly that the subbundle \mathcal{H} defined by $\theta = 0$ is spanned by $\{E_1, \dots, E_n, \frac{\partial}{\partial \zeta^1}, \dots, \frac{\partial}{\partial \zeta^n}\}$.

We now define G by $GU = GV = 0$ (or $G\Xi = 0$),

$$GE_\alpha = \frac{\partial}{\partial \bar{\zeta}^\alpha}, \quad G \frac{\partial}{\partial \zeta^a} = -\bar{E}_\alpha, \quad G\bar{E}_\alpha = \frac{\partial}{\partial \zeta^\alpha}, \quad G \frac{\partial}{\partial \bar{\zeta}^\alpha} = -E_\alpha.$$

Then it is straightforward to show that $G^2 = -I + u \otimes U + v \otimes V$ and that $GJ + JG = 0$. Setting $H = GJ$, one can easily check the corresponding conditions.

For the metric we first recall that the real metric associated to a Hermitian metric g , also denoted by g , is given by

$$g(X, Y) = \frac{1}{2} \Re g(X - iJX, Y + iJY).$$

Recall also that for the standard contact metric structure on the tangent sphere bundle (Section 9.2), a homothetic change with factor $\frac{1}{4}$ was made in the induced metric from the Sasaki metric on the tangent bundle. Similarly, we take the real Riemannian metric g on $P\tau$ to be $\frac{1}{4}$ of that used in Sections 13.2 and 13.3 now denoted by g' . This ensures that u and v are the covariant forms of U and V . The condition $g(X, GY) = -g(GX, Y)$ is equivalent to

$$\Re g'(X - iJX, G(Y - iJY)) = -\Re g'(G(X + iJX), Y + iJY).$$

Note also that

$$g'(E_\alpha, \bar{E}_\beta) = g'\left(\frac{\partial}{\partial \zeta^\alpha}, \frac{\partial}{\partial \bar{\zeta}^\beta}\right) = |\hat{\zeta}^0|^2 (\mathbf{G}_{\alpha\bar{\beta}} - P_\alpha P_{\bar{\beta}}).$$

It now becomes straightforward to check the skew-symmetry of G . Thus we have the following theorem.

Theorem 13.3 *The projectivized holomorphic tangent bundle of a Hermitian manifold carries a natural complex almost contact metric structure.*

13.4.1 A complex contact structure with nonintegrable vertical subbundle

The motivation to consider the question of a complex geodesic flow in this chapter and in the author's paper [2007] was not just the question

itself, even though that is a natural consideration, but also the question of the complex analogue of the tangent sphere bundle as a canonical example in complex contact geometry. We noted in Section 13.2 that the projectivized holomorphic cotangent bundle of a complex manifold carries a natural complex contact structure and we have just seen that the projectivized holomorphic tangent bundle carries a natural complex almost contact metric structure. In Riemannian geometry one can easily use the metric and its inverse to give a diffeomorphism between the tangent bundle and the cotangent bundle. In particular, we pass from vector fields to their dual 1-forms with ease. As remarked above, doing this in Hermitian geometry reverses type, but using conjugation, one could still construct a diffeomorphism between the holomorphic tangent bundle and the holomorphic cotangent bundle. However, it will not in general be a holomorphic or antiholomorphic map; as we noted above, the only time the components of a Hermitian metric are holomorphic functions is when they are constants. In the case of $\mathbb{C}^{n+1} \times \mathbb{C}P^n(16)$ (Example 12.2.7) the vector field of type $(1, 0)$ corresponding to the vertical subbundle of the complex contact structure is

$$\frac{\partial}{\partial w^0} + \sum_{\alpha=1}^n \bar{\zeta}^\alpha \frac{\partial}{\partial w^\alpha},$$

which is not holomorphic, but the Hermitian metric of the structure and conjugation give the underlying local complex contact form $dw^0 + \sum \zeta^\alpha dw^\alpha$. One might consider reversing our procedure for a given holomorphic 1-form θ and seek a vector field Ξ on $P\tau$ such that $\theta_\alpha = g_{\alpha\bar{B}} \bar{\Xi}^{\bar{B}}$, where B denotes that the sum is over the whole range of coordinates.. One might first try for $\mathbb{C}H^2$ the 1-forms $\theta = dw^0 + \zeta dw^1$, or analogous to the real projectivized tangent bundle of the Beltrami model of the real hyperbolic plane,

$$\theta = (1 - w^1(w^1 - \zeta w^0))dw^0 + (\zeta + w^0(w^1 - \zeta w^0))dw^1;$$

again see Section 9.5. While these lend themselves to higher-dimensional generalizations as holomorphic contact forms, via the metric on $P\tau$ they yield vector fields Ξ that do not give rise to a natural vertical subbundle, \mathcal{V} to play the role of the characteristic vector field in real contact geometry.

Instead for the projectivized holomorphic tangent bundle of $\mathbb{C}H^2$ we make a shift in the role of the coordinates and consider the local

holomorphic contact form

$$\theta = dw^1 - \zeta dw^0 = g_{\alpha\bar{B}} \bar{\Xi}^{\bar{B}}.$$

The resulting vector field via the metric is

$$\begin{aligned} \Xi = & \frac{(1 - |w^0|^2 - |w^1|^2)}{4} \left\{ -(\bar{\zeta}(1 - |w^0|^2) + w^0 \bar{w}^1) \frac{\partial}{\partial w^0} \right. \\ & + (1 - |w^1|^2 + \bar{\zeta} \bar{w}^0 w^1) \frac{\partial}{\partial w^1} \\ & \left. - \frac{\bar{w}^0 + \zeta \bar{w}^1}{1 - |w^0|^2 - |w^1|^2} (1 + |\zeta|^2 - |\zeta w^0 - w^1|^2) \frac{\partial}{\partial \zeta} \right\}. \end{aligned}$$

Note that

$$\sum_{\alpha=0}^1 G_{0\bar{\alpha}} \bar{\Xi}^{\bar{\alpha}} = -\zeta \quad \text{and} \quad \sum_{\alpha=0}^1 G_{1\bar{\alpha}} \bar{\Xi}^{\bar{\alpha}} = 1.$$

This is the analogue of the Liouville form $\sum p_i dq^i$ on the cotangent bundle, where the q^i are the generalized coordinates and p_i the momenta.

Now $\theta(\Xi) = \frac{(1 - |w^0|^2 - |w^1|^2)}{4} (1 + |\zeta|^2 - |\zeta w^0 - w^1|^2)$, and we consider the normalized form

$$u - iv = \frac{1}{\sqrt{1 - |w^0|^2 - |w^1|^2} \sqrt{1 + |\zeta|^2 - |\zeta w^0 - w^1|^2}} \theta.$$

The equation $\theta = 0$ defines the horizontal subbundle \mathcal{H} , which is spanned by

$$\left\{ \frac{\partial}{\partial \zeta} \pm \frac{\partial}{\partial \bar{\zeta}}, \left(\frac{\partial}{\partial w^0} + \zeta \frac{\partial}{\partial w^1} \right) \pm \left(\frac{\partial}{\partial \bar{w}^0} + \bar{\zeta} \frac{\partial}{\partial \bar{w}^1} \right) \right\}.$$

Consider the normalized vector field

$$U + iV = \frac{8}{\sqrt{1 - |w^0|^2 - |w^1|^2} \sqrt{1 + |\zeta|^2 - |\zeta w^0 - w^1|^2}} \Xi.$$

As we saw in Section 12.1 the defining requirement for the corresponding vertical subbundle of a complex contact structure is a vector field U such that $du(U, X) = 0$ for $X \in \mathcal{H}$, $u(U) = 1$ and $v(U) = 0$. Then U and $V = -JU$ span the vertical subbundle \mathcal{V} and one checks that U defined above satisfies these conditions.

Now recall that it is typically assumed that the vertical subbundle \mathcal{V} of a complex contact structure is integrable. The 1-form

$$(1 - |w^1|^2 + \bar{\zeta}\bar{w}^0w^1)dw^0 + (\bar{\zeta}(1 - |w^0|^2) + w^0\bar{w}^1)dw^1$$

applied to any linear combination of U and V is zero. Thus if \mathcal{V} were integrable,

$$((1 - |w^1|^2 + \bar{\zeta}\bar{w}^0w^1)dw^0 + (\bar{\zeta}(1 - |w^0|^2) + w^0\bar{w}^1)dw^1)([U, V])$$

would vanish, but it does not. To our knowledge this complex contact structure on the projectivized holomorphic tangent bundle of $\mathbb{C}H^2$ is the first known example of a complex contact structure for which \mathcal{V} is not integrable.

13.5 Special directions on complex contact manifolds and the Lie group $SL(2, \mathbb{C})$

In this section we first summarize the ideas of holomorphic and real special directions on a 3-dimensional complex contact metric manifold as given by B. Korkmaz and the author in [2009] and analogous to the real case as discussed in Chapter 11. We then discuss a 2-parameter family of complex contact metric structures on the Lie group $SL(2, \mathbb{C})$. Recall from Section 13.1 the notion of a holomorphic Anosov flow, which was introduced by E. Ghys in [1995]. On $SL(2, \mathbb{C})$ we show that for our 2-parameter family of structures, the holomorphic special directions determine subbundles which agree with the stable and unstable subbundles of the corresponding holomorphic Anosov flow. Reducing to a 1-parameter family of complex contact metric structures, we also show that $SL(2, \mathbb{C})$ admits a real vector field generating a partially hyperbolic flow whose central bundle has dimension 2.

Another treatment of complex contact structures on $SL(2, \mathbb{C})$ was given by Foreman [2002a], which we will discuss at the end of this section.

To begin let $k = h_U - h_V$. Then it is straightforward (or see Blair–Korkmaz [2009]) to show that

$$kJ + Jk = 0.$$

Since k anticommutes with J , if it is nonzero it admits a positive eigenvalue, say $\kappa > 0$ with unit eigenvector X . Then X is a horizontal vector

field and $-\kappa$ is also an eigenvector of k with eigenvector JX . Since the real dimension is 6, there is another eigenvector W of k with nonnegative eigenvalue ν . Then again W is a horizontal vector field and $-\nu$ is also an eigenvalue of k with eigenvector JW .

Definition. A vector $Y - iJY$ of type $(1, 0)$ on a complex contact metric manifold M where Y is horizontal is called a *holomorphic special direction* if

$$\nabla_{Y-iJY}(U+iV) = (\gamma+i\delta)(Y-iJY)$$

for some nonzero complex number $\gamma + i\delta$.

We remark that if $h_U = h_V$ ($k = 0$), then there are no holomorphic special directions, and from the author's paper with B. Korkmaz [2009] we have the following result.

Theorem 13.4 *Let M be a complex contact metric manifold of complex dimension 3 with $h_U \neq h_V$. If $\sigma = 0$ and if k is nonsingular on the horizontal subbundle, then there exist holomorphic special directions on M .*

One can also raise the question of real special directions on a complex contact metric manifold.

Definition. A horizontal real vector Y on a complex contact metric manifold M is called a *real special direction* for U (resp. for V) if

$$\nabla_Y U = \gamma Y$$

(resp. $\nabla_Y V = \gamma Y$) for a nonzero number γ .

Again from B. Korkmaz and the author [2009] we have for the vector field U the following result and a similar one for V .

Theorem 13.5 *Let M be a complex contact metric manifold with $\sigma = 0$. If h_U has an eigenvalue $\lambda > 1$, then there are real special directions for U . In particular, if all plane sections generated by U and a horizontal vector field have negative sectional curvatures, real special directions for U exist.*

We now study the Lie group

$$SL(2, \mathbb{C}) = \left\{ \begin{pmatrix} z_1 & z_2 \\ z_3 & z_4 \end{pmatrix} \mid |z_1 z_4 - z_2 z_3| = 1 \right\}.$$

We form a 2-parameter family of complex contact metric structures on $SL(2, \mathbb{C})$ as follows. Take $\lambda > \mu > 0$ and consider the matrices

$$\frac{1}{2}\sqrt{\lambda^2 - \mu^2} \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}, \quad \sqrt{\frac{\lambda + \mu}{2}} \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}, \quad \sqrt{\frac{\lambda - \mu}{2}} \begin{pmatrix} 0 & -1 \\ -1 & 0 \end{pmatrix}$$

in the Lie algebra $\mathfrak{sl}(2, \mathbb{C})$, which we regard as the tangent space of $SL(2, \mathbb{C})$ at the identity. Applying the differential of left translation to these matrices gives the vector fields

$$\begin{aligned} \xi_1 &= \frac{1}{2}\sqrt{\lambda^2 - \mu^2} \left(z_1 \frac{\partial}{\partial z_1} - z_2 \frac{\partial}{\partial z_2} + z_3 \frac{\partial}{\partial z_3} - z_4 \frac{\partial}{\partial z_4} \right), \\ \xi_2 &= \sqrt{\frac{\lambda + \mu}{2}} \left(z_2 \frac{\partial}{\partial z_1} - z_1 \frac{\partial}{\partial z_2} + z_4 \frac{\partial}{\partial z_3} - z_3 \frac{\partial}{\partial z_4} \right), \\ \xi_3 &= -\sqrt{\frac{\lambda - \mu}{2}} \left(z_2 \frac{\partial}{\partial z_1} + z_1 \frac{\partial}{\partial z_2} + z_4 \frac{\partial}{\partial z_3} + z_3 \frac{\partial}{\partial z_4} \right). \end{aligned}$$

The complex contact form on $SL(2, \mathbb{C})$ is

$$\omega = \frac{2}{\sqrt{\lambda^2 - \mu^2}} (z_4 dz_1 - z_2 dz_3) = u - iv.$$

Set $\xi_1 = \frac{1}{2}(U + iV)$, $\xi_2 = \frac{1}{2}(E_2 - iJE_2)$, $\xi_3 = \frac{1}{2}(E_3 - iJE_3)$. An associated metric is determined by left translation of the basis $\{U, V, E_2, JE_2, E_3, JE_3\}$, as an orthonormal basis at the identity, and the structure tensors G and $H = GJ$ are determined by $GE_2 = E_3$ and $HE_2 = -JE_3$.

The basis $\{U, V, E_2, JE_2, E_3, JE_3\}$ is an eigenvector basis of the operators h_U and h_V . In particular,

$$h_U E_2 = \lambda E_2, \quad h_U JE_2 = -\lambda JE_2, \quad h_V E_2 = \mu E_2, \quad h_V JE_2 = -\mu JE_2.$$

Using the anticommutativities $h_U G + Gh_U = 0$ and $h_V H + Hh_V = 0$, we get

$$h_U E_3 = -\lambda E_3, \quad h_U JE_3 = \lambda JE_3, \quad h_V E_3 = \mu E_3, \quad h_V JE_3 = -\mu JE_3.$$

Also, from this Lie algebra of vector fields it is straightforward to compute covariant derivatives and, in particular, to easily show that $\sigma = 0$.

We now turn to the question of holomorphic special directions in $SL(2, \mathbb{C})$. Using the eigenvalues and the eigenvectors of h_U and h_V , we

see that E_2 and JE_3 are eigenvectors of $k = h_U - h_V$ corresponding to the eigenvalues $\lambda - \mu$ and $\lambda + \mu$ respectively. If a and b are two numbers satisfying $a^2 + b^2 = \frac{\lambda - \mu}{2\lambda}$, then

$$Y = aE_2 + bJE_2 + \sqrt{\frac{\lambda + \mu}{\lambda - \mu}}bJE_3 + \sqrt{\frac{\lambda + \mu}{\lambda - \mu}}aE_3$$

gives a holomorphic special direction with

$$\nabla_{Y-iJY}(U+iV) = -\sqrt{\lambda^2 - \mu^2}(Y - iJY),$$

and

$$Z = aE_2 + bJE_2 - \sqrt{\frac{\lambda + \mu}{\lambda - \mu}}bJE_3 - \sqrt{\frac{\lambda + \mu}{\lambda - \mu}}aE_3$$

gives an independent holomorphic special direction with

$$\nabla_{Z-iJZ}(U+iV) = \sqrt{\lambda^2 - \mu^2}(Z - iJZ).$$

Theorem 13.6 *On $SL(2, \mathbb{C})$ the vector field ξ_1 is a holomorphic Anosov flow, and the stable and unstable subbundles, E^s and E^u , agree with the subbundles determined by the special directions corresponding to the vector fields Y and Z respectively.*

Proof. The complex flow associated to the holomorphic vector field

$$2\xi_1 = U + iV = \sqrt{\lambda^2 - \mu^2} \left(z_1 \frac{\partial}{\partial z_1} - z_2 \frac{\partial}{\partial z_2} + z_3 \frac{\partial}{\partial z_3} - z_4 \frac{\partial}{\partial z_4} \right)$$

is

$$\psi_z = \begin{pmatrix} z_1 e^{\sqrt{\lambda^2 - \mu^2} z} & z_2 e^{-\sqrt{\lambda^2 - \mu^2} z} \\ z_3 e^{\sqrt{\lambda^2 - \mu^2} z} & z_4 e^{-\sqrt{\lambda^2 - \mu^2} z} \end{pmatrix},$$

and its differential with respect to $\{\frac{\partial}{\partial z_1}, \frac{\partial}{\partial z_2}, \frac{\partial}{\partial z_3}, \frac{\partial}{\partial z_4}\}$ is given by

$$\psi_{z*} = \begin{pmatrix} e^{\sqrt{\lambda^2 - \mu^2} z} & 0 & 0 & 0 \\ 0 & e^{-\sqrt{\lambda^2 - \mu^2} z} & 0 & 0 \\ 0 & 0 & e^{\sqrt{\lambda^2 - \mu^2} z} & 0 \\ 0 & 0 & 0 & e^{-\sqrt{\lambda^2 - \mu^2} z} \end{pmatrix}.$$

Now for the vector field Y ,

$$\begin{aligned}
 Y - iJY &= (a + ib)(E_2 - iJE_2) + \sqrt{\frac{\lambda + \mu}{\lambda - \mu}}(a + ib)(E_3 - iJE_3) \\
 &= 2(a + ib)\xi_2 + 2\sqrt{\frac{\lambda + \mu}{\lambda - \mu}}(a + ib)\xi_3 \\
 &= -4\sqrt{\frac{\lambda + \mu}{2}}(a + ib)\left(z_1\frac{\partial}{\partial z_2} + z_3\frac{\partial}{\partial z_4}\right).
 \end{aligned}$$

Applying ψ_{z*} to $Y - iJY$ at the point p , we have

$$\psi_{z*}(Y - iJY)_p = e^{-\sqrt{\lambda^2 - \mu^2}z}(Y - iJY)_p = e^{-2\sqrt{\lambda^2 - \mu^2}z}(Y - iJY)_{\psi_z(p)}$$

and

$$\|\psi_{z*}(Y - iJY)_p\| = e^{-\sqrt{\lambda^2 - \mu^2}\Re(z)}\|(Y - iJY)_p\|.$$

Therefore the special direction determined by Y determines the stable subbundle.

Similarly, the vector field Z yields the holomorphic special direction

$$\begin{aligned}
 Z - iJZ &= 2(a + ib)\xi_2 - 2\sqrt{\frac{\lambda + \mu}{\lambda - \mu}}(a + ib)\xi_3 \\
 &= 4\sqrt{\frac{\lambda + \mu}{2}}(a + ib)\left(z_2\frac{\partial}{\partial z_1} + z_4\frac{\partial}{\partial z_3}\right).
 \end{aligned}$$

Therefore

$$\psi_{z*}(Z - iJZ)_p = e^{2\sqrt{\lambda^2 - \mu^2}z}(Z - iJZ)_{\psi_z(p)}$$

and

$$\|\psi_{z*}(Z - iJZ)_p\| = e^{\sqrt{\lambda^2 - \mu^2}\Re(z)}\|(Z - iJZ)_p\|,$$

giving the unstable subbundle. ■

We now study the real special directions associated to the vector field U .

Theorem 13.7 *If $\lambda > 1$, then there exist real special directions associated to the vector field U on $SL(2, \mathbb{C})$. Moreover, when $\mu = 1$, U determines a partially hyperbolic flow with 2-dimensional central subbundle E^c spanned by U and V .*

Proof. The first statement is clear from Theorem 13.5. In terms of the coordinates $z_j = x_j + iy_j$, when $\mu = 1$,

$$U = \frac{1}{2}\sqrt{\lambda^2 - 1} \sum_{j=1}^4 (-1)^{j+1} \left(x_j \frac{\partial}{\partial x_j} + y_j \frac{\partial}{\partial y_j} \right).$$

The corresponding flow ψ_t maps a point $(x_1, y_1, x_2, y_2, x_3, y_3, x_4, y_4)$ to the point

$$e^{\frac{\sqrt{\lambda^2 - 1}}{2}t} (x_1, y_1, 0, 0, x_3, y_3, 0, 0) + e^{-\frac{\sqrt{\lambda^2 - 1}}{2}t} (0, 0, x_2, y_2, 0, 0, x_4, y_4).$$

Consider the vector field

$$Y = aE_2 + bE_3 = -\sqrt{\frac{\lambda^2 - 1}{\lambda}} \left(x_1 \frac{\partial}{\partial x_2} + y_1 \frac{\partial}{\partial y_2} + x_3 \frac{\partial}{\partial x_4} + y_3 \frac{\partial}{\partial y_4} \right).$$

Applying ψ_{t*} at a point p , we have

$$\psi_{t*}Y_p = e^{-\frac{\sqrt{\lambda^2 - 1}}{2}t}Y_p = e^{-\sqrt{\lambda^2 - 1}t}Y_{\psi_t(p)}$$

and

$$\|\psi_{t*}Y_p\| = e^{-\frac{\sqrt{\lambda^2 - 1}}{2}t}\|Y_p\|.$$

Now consider the vector field

$$JY = aJE_2 + bJE_3 = \sqrt{\frac{\lambda^2 - 1}{\lambda}} \left(y_1 \frac{\partial}{\partial x_2} - x_1 \frac{\partial}{\partial y_2} + y_3 \frac{\partial}{\partial x_4} - x_3 \frac{\partial}{\partial y_4} \right).$$

Applying ψ_{t*} to this vector we have

$$\psi_{t*}JY_p = e^{-\frac{\sqrt{\lambda^2 - 1}}{2}t}JY_p = e^{-\sqrt{\lambda^2 - 1}t}JY_{\psi_t(p)}$$

and

$$\|\psi_{t*}JY_p\| = e^{-\frac{\sqrt{\lambda^2 - 1}}{2}t}\|JY_p\|.$$

Thus Y and JY give a subbundle, E^s , which ψ_{t*} leaves invariant and for which ψ_{t*} shortens lengths exponentially. We remark that since JE_3 is also an eigenvector of h_U with eigenvalue λ , $aJE_3 + bJE_2$ is a special direction but the corresponding vector field is not invariant under ψ_{t*} .

Similarly, we have the special direction

$$Z = aE_2 + bE_3 = -\sqrt{\frac{\lambda^2 - 1}{\lambda}} \left(x_2 \frac{\partial}{\partial x_1} + y_2 \frac{\partial}{\partial y_1} + x_4 \frac{\partial}{\partial x_3} + y_4 \frac{\partial}{\partial y_3} \right),$$

which satisfies

$$\psi_{t*} Z_p = e^{\sqrt{\lambda^2 - 1}t} Z_{\psi_t(p)}$$

and together with JZ defines an unstable bundle E^u .

Finally, consider the vector field

$$V = \frac{1}{2} \sqrt{\lambda^2 - 1} \sum_{j=1}^4 (-1)^{j+1} \left(y_j \frac{\partial}{\partial x_j} - x_j \frac{\partial}{\partial y_j} \right).$$

Applying ψ_{t*} , we have

$$\psi_{t*} V_p = V_{\psi_t(p)}.$$

Thus U defines a partially hyperbolic flow whose central bundle is 2-dimensional and spanned by U and V ; E^s and E^u are the stable and unstable bundles. ■

In [2002a] B. Foreman looked at complex contact structures on $SL(2, \mathbb{C})$ from a more algebraic point of view. He considers $PSL(2, \mathbb{C})$ which has a well-known identification with the space \mathcal{M} of Möbius transformations, the isometry group of hyperbolic 3-space H^3 . Every linear fractional transformation $z \mapsto \frac{az+b}{cz+d}$ can be written in such a way that the matrix $\begin{pmatrix} a & b \\ c & d \end{pmatrix}$ has determinant +1; the matrix is unique up to sign, giving $SL(2, \mathbb{C})$ as a 2-fold cover of \mathcal{M} . The boundary ∂H^3 of H^3 is readily identified with $\hat{\mathbb{C}} \cong S^2$. If $g \in \mathcal{M}$ has exactly one fixed point in $\hat{\mathbb{C}}$, then g is said to be *parabolic*. Suppose $g \in \mathcal{M}$ has two fixed points. Then if g has infinitely many fixed points in H^3 , g is *elliptic*; on the other hand, if g has no fixed points in H^3 and preserves an open disk or half-plane in $\hat{\mathbb{C}}$, g is *hyperbolic*. Foreman first proves an Iwasawa decomposition theorem.

Theorem 13.8 *Given two distinct points $q_1, q_2 \in \partial H^3$ and a point $x \in H^3$ lying on the geodesic connecting q_1 and q_2 , there is a unique Iwasawa decomposition of \mathcal{M} given by*

$$\mathcal{M} = K \cdot A \cdot N,$$

where

$$\begin{aligned} K &= \{g \in \mathcal{M} : g \text{ is elliptic, } g(x) = x\} \cup \{\text{id}\}, \\ A &= \{g \in \mathcal{M} : g \text{ is hyperbolic, } g(q_1) = q_1, g(q_2) = q_2\} \cup \{\text{id}\}, \\ N &= \{g \in \mathcal{M} : g \text{ is parabolic, } g(q_2) = q_2\} \cup \{\text{id}\}. \end{aligned}$$

As a real Lie algebra, $\mathfrak{sl}(2, \mathbb{C})$ has a natural complex structure given by

$$J \begin{pmatrix} a & b \\ c & -a \end{pmatrix} = \begin{pmatrix} ia & ib \\ ic & -ia \end{pmatrix}.$$

Also, from the Iwasawa decomposition of \mathcal{M} , N is the set of all parabolic transformations fixing some $q \in \hat{\mathbb{C}}$, and we denote the corresponding space by N_q . Then the corresponding Lie algebra \mathfrak{n}_q is a J -invariant subspace of real dimension 2. Again taking $q \in \hat{\mathbb{C}}$, let $F_q = \{g \in \mathcal{M} : g(q) = q\}$, a Lie subgroup of \mathcal{M} ; its Lie algebra \mathfrak{f}_q is a J -invariant subspace of complex dimension 2.

We are interested in left-invariant complex contact structures on \mathcal{M} , i.e., those that are described by $\ker \theta$ for some $\theta \in \mathfrak{sl}(2, \mathbb{C})^*$. The main result of Forman [2002] is the following.

Theorem 13.9 *Let $\ker \theta$ be a left-invariant 2-dimensional subspace of $\mathfrak{sl}(2, \mathbb{C})$. Then $\ker \theta$ satisfies exactly one of the following statements:*

1. $\ker \theta = \mathfrak{n}_{q_1} + \mathfrak{n}_{q_2}$ for two distinct $q_1, q_2 \in \partial H^3$. In this case $\ker \theta$ is a complex contact structure.
2. $\ker \theta = \mathfrak{f}_q$ for some $q \in \partial H^3$. In this case $\ker \theta$ is a foliation.

In the first case, for any $f \in \mathcal{M} - \{\text{id}\}$ such that $f(q_1) = q_1, f(q_2) = q_2$ there is a unique complex contact structure θ such that the corresponding characteristic vector field $\xi (= \frac{1}{2}(U + iV))$ satisfies $\exp(\xi) = f$.

14

3-Sasakian Manifolds

In this chapter we will give more of a survey of 3-Sasakian manifolds and only a few proofs. A more thorough treatment is given in the book by Boyer and Galicki [2008, Chapter 13].

14.1 3-Sasakian manifolds

If a manifold M^{2m+1} admits three almost contact structures (ϕ_i, ξ_i, η_i) , $i = 1, 2, 3$, satisfying the following for an even permutation (i, j, k) of $(1, 2, 3)$,

$$\begin{aligned}\phi_k &= \phi_i \phi_j - \eta_j \otimes \xi_i = -\phi_j \phi_i + \eta_i \otimes \xi_j, \\ \xi_k &= \phi_i \xi_j = -\phi_j \xi_i, \quad \eta_k = \eta_i \circ \phi_j = -\eta_j \circ \phi_i,\end{aligned}$$

then the manifold is said to have an *almost contact 3-structure*. This notion was introduced by Kuo [1970] and independently under the name *almost coquaternion structure* by Udriste [1969]. Some authors follow different sign conventions, taking $\phi_k = -\phi_i \phi_j + \eta_j \otimes \xi_i$, etc. (see, e.g., the latter part of Section 14.2, the author's paper with Baikoussis [1995], Boyer and Galicki [2008, p. 486]). Note that given two almost contact

metric structures satisfying

$$\begin{aligned}\phi_1\phi_2 - \eta_2 \otimes \xi_1 &= -\phi_2\phi_1 + \eta_1 \otimes \xi_2, \\ \phi_1\xi_2 &= -\phi_2\xi_1, \quad \eta_1 \circ \phi_2 = -\eta_2 \circ \phi_1, \quad \eta_2(\xi_1) = \eta_1(\xi_2) = 0,\end{aligned}$$

then there exists a third almost contact structure defined by

$$\phi_3 = \phi_1\phi_2 - \eta_2 \otimes \xi_1, \quad \xi_3 = \phi_1\xi_2, \quad \eta_3 = -\eta_2 \circ \phi_1$$

giving an almost contact 3-structure.

Now given an almost contact 3-structure (ϕ_i, ξ_i, η_i) , define on $M^{2m+1} \times \mathbb{R}$ three almost complex structures J_i using each of the almost contact structures as in Section 6.1. It is then easy to check that $J_k = J_i J_j = -J_j J_i$. Therefore $M^{2m+1} \times \mathbb{R}$ has an almost quaternionic structure, and hence its dimension is a multiple of 4. Thus the dimension of a manifold with an almost contact 3-structure is of the form $4n + 3$. Tachibana and Yu [1970] used this idea to show that there cannot be a fourth almost contact structure (ϕ_4, ξ_4, η_4) with $\eta_i(\xi_4) = \eta_4(\xi_i) = 0, i = 1, 2, 3$, and satisfying the anticommutativity conditions with the first three structures. To see this, let J_4 be the almost complex structure on $M^{2m+1} \times \mathbb{R}$ constructed using (ϕ_4, ξ_4, η_4) . Then pairing J_4 with each of J_1, J_2, J_3 yields $J_4 J_i = -J_i J_4, i = 1, 2, 3$. This contradicts $J_3 J_4 = J_1 J_2 J_4 = -J_1 J_4 J_2 = J_4 J_1 J_2 = J_4 J_3$.

The normality of these almost contact structures was discussed by Yano, Ishihara and Konishi [1973]. In particular, if two of the almost contact structures are normal, then so is the third.

Kuo [1970] proved that given an almost contact 3-structure, there exists a Riemannian metric compatible with each of them, and hence we can speak of an *almost contact metric 3-structure* $(\phi_i, \xi_i, \eta_i, g), i = 1, 2, 3$. He also showed that the structural group of the tangent bundle is reducible to $Sp(n) \times I_3$. Moreover, the vector fields $\{\xi_1, \xi_2, \xi_3\}$ are orthonormal with respect to the compatible metric.

If the three structures $(\phi_i, \xi_i, \eta_i, g)$ are contact metric structures, we say that M^{4n+3} has a *contact metric 3-structure*. If the three structures are Sasakian, we say that M^{4n+3} has a *3-Sasakian structure*, sometimes called a *Sasakian 3-structure*, and M^{4n+3} is a *3-Sasakian manifold*.

As we remarked at the end of Section 6.5, Boyer and Galicki [2008, p. 477] define a 3-Sasakian manifold by considering the cone $\mathbb{R}_+ \times M$. A Riemannian manifold (M^m, g) is a 3-Sasakian manifold if and only

if the holonomy group of the cone $(\mathbb{R}_+ \times M, dr^2 + r^2g)$ reduces to a subgroup of $Sp(\frac{m+1}{4})$. Again we see that $m = 4n + 3$, $n \geq 1$. Moreover, $(\mathbb{R}_+ \times M, dr^2 + r^2g)$ is a hyper-Kähler manifold, i.e., it has a quaternionic structure consisting of three global almost complex structures that are Kähler with respect to the metric $dr^2 + r^2g$. For a proof of the equivalence of these definitions see Boyer, Galicki, Mann [1994].

One can also have the notion of an almost hyper-Kähler manifold, where the three fundamental 2-forms of the almost Hermitian structures are only required to be closed. However, it is an important result of Hitchin [1987] that an almost hyper-Kähler manifold is in fact hyper-Kähler.

Now the cone over a manifold with a contact metric 3-structure has an immediate almost hyper-Kähler structure and hence by the above mentioned result of Hitchin is a hyper-Kähler manifold. Thus we have the following theorem of Kashiwada [2001]; her proof used a generalization of Hitchin's result that she gave in [1998].

Theorem 14.1 *Every contact metric 3-structure is 3-Sasakian.*

There is also a notion of hypercontact structure that was introduced by Geiges and Thomas [1995]. A *hypercontact structure* is a triple of contact forms $(\alpha_1, \alpha_2, \alpha_3)$ and an almost contact metric 3-structure $(\phi_i, \xi_i, \eta_i, g)$, $i = 1, 2, 3$, such that each $d\alpha_i$ is the covariant form of ϕ_i . If $\eta_i = \alpha_i$, this would be a contact metric 3-structure and hence 3-Sasakian. Perrone [2002] considers hypercontact structures for which the characteristic vector field ζ_i of α_i is the metric dual of α_i and calls such a structure a *hypercontact metric structure*. In their paper Geiges and Thomas prove a connected sum theorem for hypercontact manifolds. In the Perrone paper he proves that a 3-dimensional hypercontact metric manifold is either 3-Sasakian and locally isometric to $S^3(1)$ or is locally isometric to $SL(2, \mathbb{R})$ with a left-invariant hypercontact metric structure that is not a contact metric 3-structure.

Again if M^{4n+3} has two Sasakian structures $(\phi_i, \xi_i, \eta_i, g)$, $i = 1, 2$, with ξ_1 and ξ_2 orthogonal, then the third structure defined by $\xi_3 = \phi_1\xi_2$ and $\phi_3X = -\nabla_X\xi_3$ gives a 3-Sasakian structure. Exploring this idea further, Tachibana and Yu [1970] proved the following theorem.

Theorem 14.2 *Let M be a complete simply connected Riemannian manifold admitting Sasakian structures $(\phi_i, \xi_i, \eta_i, g)$, $i = 1, 2$, such that*

$g(\xi_1, \xi_2)$ is a nonconstant function on M . Then M is isometric to a unit sphere.

The proof of this theorem is to differentiate the function $f = g(\xi_1, \xi_2)$ twice and use the Sasakian conditions to show that f satisfies $\nabla_l \nabla_k f_j + 2f_l g_{kj} + f_k g_{lj} + f_j g_{lk} = 0$, where $f_j = \nabla_j f$. Then by a well-known theorem of Obata [1965] (see also Tanno [1978a]), M is isometric to a unit sphere.

Using $\nabla_X \xi_i = -\phi_i X$ one readily obtains on a manifold with a 3-Sasakian structure that $[\xi_i, \xi_j] = 2\xi_k$. Thus the subbundle spanned by $\{\xi_1, \xi_2, \xi_3\}$ is integrable with totally geodesic leaves that are easily seen to be of constant curvature $+1$. Notice also that from $\phi_i \xi_j = \xi_k$, etc., each leaf of the foliation is itself a 3-Sasakian manifold.

The canonical example of a manifold with a 3-Sasakian structure is the sphere S^{4n+3} . Its structure is readily obtained by taking S^{4n+3} as a hypersurface in \mathbb{H}^{n+1} . Each of the three almost complex structures forming the quaternionic structure of \mathbb{H}^{n+1} applied to the outer normal of the sphere gives a vector field ξ_i , $i = 1, 2, 3$, on S^{4n+3} . These three vector fields are orthogonal and give rise to the standard 3-Sasakian structure on S^{4n+3} . This 3-Sasakian structure on S^{4n+3} also projects under the Hopf fibration to the quaternionic structure on quaternionic projective space $\mathbb{H}P^n$. We will briefly discuss generalizations of this fibration below.

We first, however, prove an early result of Kashiwada [1971] that a 3-Sasakian manifold is Einstein; this can be regarded as analogous to the well-known fact that a quaternionic Kähler manifold is Einstein. Kashiwada's proof is computational relative to an orthonormal basis, but here we give a noncomputational proof utilizing properties of cones and hyper-Kähler manifolds.

Theorem 14.3 *Let M^{4n+3} be a manifold with a 3-Sasakian structure*

$$(\phi_i, \xi_i, \eta_i, g), \quad i = 1, 2, 3.$$

Then M^{4n+3} is an Einstein space with positive scalar curvature.

Proof. First, recall that the cone over a Riemannian manifold M is Ricci flat if and only if M is Einstein with Einstein constant equal to $\dim M - 1$. Second, recall that hyper-Kähler manifolds are Ricci flat. Thus considering the cone over M^{4n+3} , we have the result. ■

As a corollary, or from our previous remarks, we see that a 3-dimensional manifold with a 3-Sasakian structure is of constant curvature $+1$.

Additional curvature properties of 3-Sasakian manifolds include the following. For a vector X orthogonal to $\{\xi_1, \xi_2, \xi_3\}$ ($4n+3 \geq 7$), the sum of the three ϕ -sectional curvatures satisfies

$$\sum_{i=1}^3 K(X, \phi_i X) = 3;$$

Tanno [1971]. If the sectional curvature of plane sections orthogonal to $\{\xi_1, \xi_2, \xi_3\}$ is constant ($4n+3 \geq 7$), the 3-Sasakian manifold is of constant curvature $+1$; Konishi and Funabashi [1976].

All complete 3-dimensional 3-Sasakian manifolds were classified by Sasaki [1972] in the following theorem.

Theorem 14.4 *A complete 3-dimensional 3-Sasakian manifold is a quotient S^3/Γ , where Γ is one of the following finite subgroups of Clifford translations on S^3 :*

1. $\Gamma = \{I\}$,
2. $\Gamma = \{\pm I\}$,
3. Γ is the cyclic group of order $q > 2$ generated by

$$\begin{pmatrix} \cos \frac{2\pi}{q} & -\sin \frac{2\pi}{q} & 0 & 0 \\ \sin \frac{2\pi}{q} & \cos \frac{2\pi}{q} & 0 & 0 \\ 0 & 0 & \cos \frac{2\pi}{q} & -\sin \frac{2\pi}{q} \\ 0 & 0 & \sin \frac{2\pi}{q} & \cos \frac{2\pi}{q} \end{pmatrix},$$

4. Γ is a group of Clifford translations corresponding to a binary dihedral group or the binary polyhedral groups of the regular tetrahedron, octahedron, or icosahedron.

Boyer, Galicki and Mann [1994] classified all 3-Sasakian homogeneous spaces.

Theorem 14.5 *Let M be a 3-Sasakian homogeneous space. Then M is one of the following:*

$$\frac{Sp(n+1)}{Sp(n)} \simeq S^{4n+3}, \quad \frac{Sp(n+1)}{Sp(n) \times \mathbb{Z}_2} \simeq \mathbb{R}P^{4n+3},$$

$$\frac{SU(m)}{S(U(m-2) \times U(1))}, \quad \frac{SO(k)}{SO(k-4) \times Sp(1)},$$

$$\frac{G_2}{Sp(1)}, \quad \frac{F_4}{Sp(3)}, \quad \frac{E_6}{SU(6)}, \quad \frac{E_7}{Spin(12)}, \quad \frac{E_8}{E_7}.$$

For the first two cases when $n = 0$, $Sp(0)$ is the identity group; $m \geq 3$; and $k \geq 7$. Moreover, M fibers over a quaternionic Kähler manifold; the fiber is $Sp(1)$ for S^{4n+3} and $SO(3)$ in the other cases.

There are in addition many inhomogeneous 3-Sasakian spaces, even simply connected ones. Boyer, Galicki and Mann [1994] prove that there are infinitely many homotopically distinct 7-dimensional strongly inhomogeneous compact simply connected 3-Sasakian manifolds.

Let (M, ϕ, ξ, η, g) be a Sasakian manifold with complete Riemannian metric g and denote by $I(M)$ the isometry group of g . Let $A(M)$ be the automorphism group of the Sasakian structure (ϕ, ξ, η, g) , i.e., $A(M)$ is the subgroup of isometries which also preserve ϕ , ξ and η . In [1970] Tanno proved the following theorem.

Theorem 14.6 *Let (M, ϕ, ξ, η, g) be a complete Sasakian manifold that is not of constant curvature. Then either*

$$\dim I(M) = \dim A(M) \text{ or } \dim I(M) = \dim A(M) + 2.$$

Moreover, $\dim I(M) = \dim A(M)$ if and only if M does not admit a 3-Sasakian structure, and $\dim I(M) = \dim A(M) + 2$ if and only if M admits a 3-Sasakian structure.

Turning to fibration questions, Tanno [1971] proves that of the three dimensional 3-Sasakian manifolds in Theorem 14.4, only $S^3(1)$ and $\mathbb{R}P^3(1)$ are regular with respect to any of the three characteristic vector fields. Consequently, we have the following result of Tanno [1971].

Lemma 14.1 *Let $(M^{4n+3}, \phi_i, \xi_i, \eta_i, g)$, $i = 1, 2, 3$, be a complete Riemannian manifold with a 3-Sasakian structure. If ξ_1 is regular then so are ξ_2 and ξ_3 , and the leaves of the foliation induced by $\{\xi_1, \xi_2, \xi_3\}$ are isometric to $S^3(1)$ and $\mathbb{R}P^3(1)$.*

Tanno's proof uses only the K-contact property, but by Kashiwada's result (Theorem 14.1) the structure is 3-Sasakian. Moreover, M^{4n+3} is a fiber bundle over an almost quaternionic Kähler manifold M^{4n} that is both Einstein and quaternionic Kähler. These properties of 3-Sasakian manifolds have been proved by various authors at different points in time. Under the assumptions of regularity, Tanno proved in [1971] that the base manifold of the fibration is Einstein. When $n \geq 2$, Ishihara

[1973] proved that the base manifold is quaternionic Kähler. Conversely, if the induced structure on the base manifold is quaternionic Kähler, then the structure is 3-Sasakian; Konishi [1973]. The reason for the restriction $n \geq 2$ is that the usual definition of a quaternionic Kähler manifold M^{4n} in terms of holonomy breaks down in dimension 4; more precisely, the condition that holonomy group be contained in $Sp(n) \cdot Sp(1)$ ($= Sp(n) \times Sp(1)/\{\pm I\}$) $\subset SO(4n)$ becomes in dimension 4 simply the orientability of the manifold since $Sp(1) \cdot Sp(1) = SO(4)$. Thus in dimension 4, a manifold is said to be a quaternionic Kähler manifold if it is Einstein with nonzero scalar curvature and self-dual (see, e.g., LeBrun [1991]). With this understanding Tanno [1996] extended Ishihara's result to include $n = 1$. We summarize this discussion in the following theorem.

Theorem 14.7 *Let $(M^{4n+3}, \phi_i, \xi_i, \eta_i, g)$, $i = 1, 2, 3$, be a complete Riemannian manifold with a contact metric 3-structure. If one of the ξ_i 's is regular, then M^{4n+3} is an $Sp(1)$ or $SO(3)$ bundle over a quaternionic Kähler manifold M^{4n} .*

The converse question was considered by Konishi in [1975], i.e., starting with a quaternionic Kähler manifold M^{4n} with $n > 1$, she constructs a canonical $SO(3)$ bundle, M^{4n+3} , over M^{4n} . When the scalar curvature of M^{4n} is positive, M^{4n+3} has a 3-Sasakian structure. Tanno [1996] extends this construction to the case $n = 1$ and to the case of negative scalar curvature. We state these results as follows.

Theorem 14.8 *Let M^{4n} be a quaternionic Kähler manifold with nonzero scalar curvature. Then there exists a canonical $SO(3)$ bundle, M^{4n+3} , over M^{4n} . If the scalar curvature of M^{4n} is positive, M^{4n+3} admits a 3-Sasakian structure; if the scalar curvature is negative, M^{4n+3} admits a pseudo-3-Sasakian structure, the signature of the metric g being $(3, 4n)$.*

In dimensions 7 and 11 the complete principal Riemannian fibrations with 3-Sasakian structure of positive scalar curvature were determined by Boyer, Galicki and Mann [1993, p.253]. They are S^7 , $\mathbb{R}P^7$ and $SU(3)/U(1)$ in dimension 7, and S^{11} , $\mathbb{R}P^{11}$, $SU(4)/S(U(2) \times U(1))$ and $G_2/SU(2)$ in dimension 11.

In general, the foliation determined by the vector fields $\{\xi_1, \xi_2, \xi_3\}$ is not a fibration, and Boyer, Galicki, and Mann [1994] prove the following result.

Theorem 14.9 *Let $(M^{4n+3}, \phi_i, \xi_i, \eta_i, g)$, $i = 1, 2, 3$, be a 3-Sasakian manifold such that the vector fields ξ_1, ξ_2, ξ_3 are complete. Then the space of leaves of the foliation determined by $\{\xi_1, \xi_2, \xi_3\}$ is a quaternionic Kähler orbifold of dimension $4n$ with positive scalar curvature $16n(n+2)$.*

In view of our study of twistor spaces as complex contact manifolds over quaternionic Kähler manifolds, we also consider fibrations of a manifold with a contact metric 3-structure by one of the structure vector fields. The following result of Ishihara and Konishi [1979] is not surprising.

Theorem 14.10 *Let M^{4n+3} be a manifold with a contact metric (Sasakian) 3-structure and suppose that one of the structures, say $(\phi_1, \xi_1, \eta_1, g)$, is regular (and Sasakian). Then the orbit space M^{4n+3}/ξ_1 admits a complex contact metric structure. Moreover, the complex contact metric structure on the orbit space is a Kähler Einstein structure of positive scalar curvature.*

Without the regularity, this question was taken up by Boyer and Galicki [1997], who gave an orbifold version and showed that the space $Z = M^{4n+3}/\xi_1$ is the twistor space of a quaternionic Kähler orbifold.

Some important results on the topology of a compact 3-Sasakian manifold are the following due to Galicki and Salamon [1996] (see also Boyer and Galicki [2008, Section 13.5]). In Section 6.8 we noted that the odd Betti numbers of a compact Sasakian manifold are even up to middle dimension. Galicki and Salamon prove a stronger result for 3-Sasakian manifolds.

Theorem 14.11 *Let M^{4n+3} be a compact 3-Sasakian manifold. Then the odd Betti numbers b_{2k+1} are all zero for $0 \leq k \leq n$.*

In the regular case, Galicki and Salamon [1996] relate the Betti numbers of M^{4n+3} to those of the quaternionic Kähler base space M^{4n} and the intermediate space Z obtained as the Boothby–Wang fibration with respect to one of the characteristic vector fields. In particular, the odd Betti numbers of M^{4n} and Z vanish. Furthermore, one has the relations $b_{2k}(M^{4n+3}) = b_{2k}(Z) - b_{2k-2}(Z) = b_{2k}(M^{4n}) - b_{2k-4}(M^{4n})$, $k \leq n$. Galicki and Salamon also prove the following remarkable result.

Theorem 14.12 *Up to isometries, there are only finitely many compact regular 3-Sasakian manifolds in each dimension $4n + 3$, $n \geq 0$, and the*

only compact regular 3-Sasakian manifolds with $b_2 > 0$ are the spaces

$$\frac{U(m)}{U(m-2) \times U(1)}, \quad m \geq 3.$$

In the same paper Galicki and Salomon obtain the following result on sums of Betti numbers.

Theorem 14.13 *The Betti numbers of a compact regular 3-Sasakian manifold of dimension $4n + 3$ satisfy*

$$\sum_{k=1}^n k(n+1-k)(n+1-2k)b_{2k} = 0.$$

Moreover, if $b_4 = 0$ and $n = 3$ or 4 , then the manifold is either a sphere or a real projective space.

14.2 Integral submanifolds

When we have one contact structure on a manifold of dimension $2n + 1$, we have seen that the maximum dimension of an integral submanifold is n . In the present context we have three independent contact structures on a manifold of dimension $4n + 3$, and we begin with the following lemma.

Lemma 14.2 *The maximum dimension of a submanifold which is an integral submanifold of all three contact structures is n .*

Proof. If X_1, \dots, X_r is a local basis tangent to such a submanifold, then the $\phi_i X_a$, $i = 1, 2, 3$, $a = 1, \dots, r$, are normal to the submanifold as well as perpendicular to ξ_i , $i = 1, 2, 3$, since $\phi_i \xi_j = \xi_k$. Moreover, from $\phi_k = \phi_i \phi_j - \eta_j \otimes \xi_i$ these vectors are independent. Thus the codimension is at least $3r + 3$, and hence $r \leq n$. ■

Thus in dimension 7 these would be integral (Legendre) curves, and that they are plentiful can be seen as follows. The characteristic vector fields of the standard 3-Sasakian structure on S^7 are tangent to the fibers S^3 of S^7 viewed as a principal S^3 -bundle over S^4 . Thus the horizontal lift of a curve on S^4 gives a curve in S^7 which is a Legendre curve for each of the three contact structures. Curves in S^7 which are Legendre

curves for all three contact structures and of constant curvature and unit torsion were classified by Baikoussis and the author in [1995].

In contrast to this lemma, it is possible to have submanifolds of dimension up to $2n + 1$ which are integral submanifolds of two of the three contact structures. We will discuss this briefly in the case of S^7 . Note first that if M^3 is a 3-dimensional submanifold of a manifold with 3-Sasakian structure that is an integral submanifold of the first Sasakian structure $(\phi_1, \xi_1, \eta_1, g)$ and an invariant submanifold with respect to the third Sasakian structure $(\phi_3, \xi_3, \eta_3, g)$, then it is an integral submanifold of the second Sasakian structure $(\phi_2, \xi_2, \eta_2, g)$. To see this, let X be tangent to M^3 ; then $\eta_2(X) = g(\xi_2, X) = g(\phi_3 \xi_1, X) = -g(\xi_1, \phi_3 X) = -\eta_1(\phi_3 X) = 0$.

To give a couple of examples, consider Euclidean space E^8 with three complex structures (or quaternionic structure),

$$I = \begin{pmatrix} 0 & -I_4 \\ I_4 & 0 \end{pmatrix}, \quad J = \begin{pmatrix} 0 & 0 & 0 & I_2 \\ 0 & 0 & -I_2 & 0 \\ 0 & I_2 & 0 & 0 \\ -I_2 & 0 & 0 & 0 \end{pmatrix}, \quad K = -IJ,$$

where I_n denotes the $n \times n$ identity matrix. Let \mathbf{x} denote the position vector of the unit sphere in E^8 , and as usual define three vector fields on S^7 by $\xi_1 = -I\mathbf{x}$, $\xi_2 = -J\mathbf{x}$, $\xi_3 = -K\mathbf{x}$. The dual 1-forms η_i are three independent contact structures on S^7 . Now consider the linear subspace L of E^8 defined by $x_5 = x_6 = x_7 = x_8 = 0$. The intersection $S^3 = S^7 \cap L$ is then an integral submanifold for η_1 and η_2 , but ξ_3 is tangent, and this sphere is invariant under the action of K restricted to S^7 . Now consider the torus in the 3-sphere we just constructed defined by $x_1^2 + x_3^2 = \frac{1}{2}$, $x_2^2 + x_4^2 = \frac{1}{2}$. This torus is an integral surface for both η_1 and η_2 , but ξ_3 is tangent. Note also that the image of the tangent space under ϕ_3 is normal.

We now state the following result from Baikoussis and the author [1995].

Theorem 14.14 *Let M^3 be a 3-dimensional submanifold of S^7 isometrically immersed as an integral submanifold of the Sasakian structure $(\phi_1, \xi_1, \eta_1, \bar{g})$ (and $(\phi_2, \xi_2, \eta_2, \bar{g})$) and an invariant submanifold of the third Sasakian structure $(\phi_3, \xi_3, \eta_3, \bar{g})$. Then M^3 is a principal circle bundle over a holomorphic Legendre curve in \mathbb{CP}^3 . Moreover, if M^3 is of*

constant ϕ -sectional curvature then either M^3 is totally geodesic or a principal circle bundle over the holomorphic (Calabi) curve $\mathbb{C}P^1(\frac{4}{3})$.

As a corollary, we note that a holomorphic Legendre curve of constant Gaussian curvature in $\mathbb{C}P^3$ cannot lie in a totally geodesic subspace $\mathbb{C}P^2$. Similarly, if M^3 is a 3-dimensional submanifold of S^7 isometrically immersed as an integral submanifold of one of the Sasakian structures and as an invariant submanifold of one of the other Sasakian structures and if M^3 is of constant ϕ -sectional curvature and not totally geodesic, then M^3 cannot lie in any totally geodesic Sasakian submanifold (S^5) of S^7 .

We now turn to the case of a 2-dimensional submanifold of S^7 which is an integral submanifold of two of the Sasakian structures. Even though it cannot be an integral submanifold for the third structure, we still can have that ϕ_3 maps the tangent bundle into the normal bundle as in the above example of a torus. In this regard we have the following result of Baikoussis and the author [1995].

Theorem 14.15 *Let M^2 be a surface isometrically immersed in S^7 as an integral submanifold of the Sasakian structures $(\phi_1, \xi_1, \eta_1, \bar{g})$ and $(\phi_2, \xi_2, \eta_2, \bar{g})$ and suppose that $\phi_3(TM) \subset T^\perp M$. Then M^2 is flat and ξ_3 is tangent to M^2 .*

Bibliography

- Abbas, C., Cieliebak, K. and Hofer, H.
[2005] The Weinstein conjecture for planar contact structures in dimension three, *Comment. Math. Helv.*, **80**, 771–793.
- Abbena, E.
[1984] An example of an almost Kähler manifold which is not Kählerian, *Bollettino U.M.I.*, (6) **3-A**, 383–392.
- Abe, K.
[1976] Some examples of non-regular almost contact structures on exotic spheres, *Tôhoku Math. J.*, **28**, 429–435.
[1977] On a generalization of the Hopf fibration, I, *Tôhoku Math. J.*, **29**, 335–374.
- Abe, K. and Erbacher, J.
[1975] Non-regular contact structures on Brieskorn manifolds, *Bull. Amer. Math. Soc.*, **81**, 407–409.
- Albers, P. and Hofer, H.
[2009] On the Weinstein conjecture in higher dimensions, *Comment. Math. Helv.*, **84**, 429–436.
- Alegre, P., Blair, D. E. and Carriazo, A.
[2004] Generalized Sasakian space-forms, *Israel J. Math.*, **14**, 157–183.
- Alegre, P. and Carriazo, A.
[2008] Structures on generalized Sasakian space-forms, *Differential Geom. Appl.*, **26**, 656–666.
- Alekseevskii, D. V.
[1968] Riemannian spaces with exceptional holonomy groups, *Funktsional Anal. i Prilozhen*, **2**, 1–10; English translation: *Functional Anal. Appl.*, **2**, 97–105.

Altschuler, S. and Wu, L.

[2000] On deforming confoliations, *J. Differential Geometry*, **54**, 75–97.

Anosov, D. V.

[1967] Geodesic Flows on Closed Riemann Manifolds with Negative Curvature, *Proc. Steklov Inst. Math.*, **90**; English translation: *Amer. Math. Soc.*, (1969).

Apostolov, V. and Draghici, T.

[1999] Hermitian conformal classes and almost Kähler structures on 4-manifolds, *Differential Geom. Appl.*, **11**, 179–195.

Apostolov, V., Draghici, T. and Moroianu, A.

[2001] A splitting theorem for Kähler manifolds whose Ricci tensors have constant eigenvalues, *International J. Math.*, **12**, 769–789.

[2006] The odd-dimensional Goldberg conjecture, *Math. Nachr.*, **279**, 948–952.

Armstrong, J.

[1998] Almost Kähler geometry, Ph.D. thesis, Oxford.

Auslander, L., Green, L. and Hahn, F.

[1963] Flows on Homogeneous Spaces, *Annals of Math. Studies*, **53**, Princeton.

Baikoussis, C. and Blair, D. E.

[1992] Integral surfaces of Sasakian space forms, *J. Geom.*, **43**, 30–40.

[1994] On Legendre curves in contact 3-manifolds, *Geometriae Dedicata*, **49**, 135–142.

[1995] On the geometry of the 7-sphere, *Results in Math.*, **27**, 5–16.

Baikoussis, C., Blair, D. E. and Gouli-Andreou, F.

[1998] Holomorphic Legendre curves in the complex Heisenberg group, *Bull. Inst. Math. Acad. Sinica*, **26**, 179–194.

Baikoussis, C., Blair, D. E. and Koufogiorgos, Th.

[1995] Integral submanifolds of Sasakian space forms $\bar{M}^7(k)$, *Results in Math.*, **27**, 207–226.

Bang, K.

[1994] Riemannian Geometry of Vector Bundles, thesis, Michigan State University.

Bang, K. and Blair, D. E.

[2008] The Schouten tensor and conformally flat manifolds, *Topics in Differential Geometry*, Editura Academiei Române, Bucharest, 1–28.

Banyaga, A.

[1990] A note on Weinstein's conjecture, *Proc. Amer. Math. Soc.*, **109**, 855–858.

Banyaga, A. and Rukimbira, P.

[1994] Weak stability of almost regular contact foliations, *J. Geom* **50**, 16–27.

Barletta, E. and Dragomir, S.

[2001] Differential equations on contact Riemannian manifolds, *Ann. Scuola Norm. Sup. Pisa Cl. Sci.*, **30**, 63–95.

Bejancu, A.

[1978] CR-submanifolds of a Kaehler manifold I, *Proc. Amer. Math. Soc.*, **69**, 135–142.

Belgun, F. A.

[2000] On the metric structure of non-Kähler complex surfaces, *Math. Ann.*, **317**, 1–40.

Bennequin, D.

[1983] Entrelacements et équations de Pfaff, *Astérisque*, **107–108**, 87–161.

Benson, C. and Gordon, C. S.

[1988] Kähler and symplectic structures on nilmanifolds, *Topology*, **27**, 513–518.

[1990] Kähler structures on compact solvmanifolds, *Proc. Amer. Math. Soc.*, **108**, 971–908.

Berger, M.

[1966] Remarques sur le groupe d'holonomie des variétés Riemanniennes, *C.R. Acad. Sc. Paris*, **262**, 1316–1318.

[1970] Quelques formules de variation pour une structure riemannienne, *Ann. Sc. École Norm. Sup.*, **3**, 285–294.

Berger, M. and Ebin, D.

[1969] Some decompositions of the space of symmetric tensors on a Riemannian manifold, *J. Differential Geometry*, **3**, 379–392.

Berndt, J.

[1997] Riemannian geometry of complex two-plane Grassmannians, *Rend. Sem. Mat.*, Univ. Pol. Torino, **55**, 19–83.

Berndt, J. and Vanhecke, L.

[1999] ϕ -symmetric spaces and weak symmetry, *Bollettino U.M.I.*, (8) **2-B**, 389–392.

Besse, A. L.

[1987] *Einstein Manifolds*, Springer, Berlin.

Bishop, R. L. and Crittenden, R. J.

[1964] *Geometry of Manifolds*, Academic Press, New York.

Blair, D. E.

[1967] The theory of quasi-Sasakian structures, *J. Differential Geometry*, **1**, 331–345.

- [1971] Almost contact manifolds with Killing structure tensors, *Pacific J. Math.*, **39**, 285–292.
 - [1976] On the non-existence of flat contact metric structures, *Tôhoku Math. J.*, **28**, 373–379.
 - [1977] Two remarks on contact metric structures, *Tôhoku Math. J.*, **29**, 319–324.
 - [1978] On the non-regularity of tangent sphere bundles, *Proc. Royal Soc. Edinburgh*, **82A**, 13–17.
 - [1983] On the set of metrics associated to a symplectic or contact form, *Bull. Inst. Math. Acad. Sinica*, **11**, 297–308.
 - [1984] Critical associated metrics on contact manifolds, *J. Austral. Math. Soc.*, Series A, **37**, 82–88.
 - [1989] When is the tangent sphere bundle locally symmetric? *Geometry and Topology*, World Scientific, Singapore, 15–30.
 - [1991a] The “total scalar curvature” as a symplectic invariant and related results, *Proc. 3rd Congress of Geometry*, Thessaloniki, 79–83.
 - [1991b] Critical associated metrics on contact manifolds III, *J. Austral. Math. Soc.*, Series A, **50**, 189–196.
 - [1996] On the class of contact metric manifolds with a 3- τ -structure, *Note di Mat.*, **16**, 99–104.
 - [1998] Special directions on contact metric manifolds of negative ξ -sectional curvature, *Ann. Fac. Sc. Toulouse VII*, 365–378.
 - [2000] Curvature and conformally Anosov flows in contact geometry, Congresso Internazionale in onore di Pasquale Calapso, *Rend. Sem. Mat. Messina*, 41–50.
 - [2007] A complex geodesic flow, *Proc. Conference RIGA*, 29–42.
- Blair, D. E., and Carriazo, A.
- [2000/2001] The contact Whitney sphere, *Note di Matematica*, **20**, 125–133.
- Blair, D. E., and Chen, B.-Y.
- [1979] On CR-submanifolds of Hermitian manifolds, *Israel J. Math.*, **34**, 353–363.
- Blair, D. E., and Chen, H.
- [1992] A classification of 3-dimensional contact metric manifolds with $Q\phi = \phi Q$, II, *Bull. Inst. Math. Acad. Sinica*, **20**, 379–383.
- Blair, D. E., Dillen, F., Verstraelen, L. and Vrancken, L.
- [1996] Calabi curves as holomorphic Legendre curves and Chen’s inequality, *Kyungpook Math. J.*, **35**, 407–416.
- Blair, D. E. and Draghici, T.
- [2009] Remarks on Weyl curvature in almost Kähler geometry, *Bull. Math. Soc.Sci. Math. Roumanie*, **52**, 241–249.

- Blair, D. E. and Goldberg, S. I.
 [1967] Topology of almost contact manifolds, *J. Differential Geometry*, **1**, 347–354.
- Blair, D. E. and Ianus, S.
 [1986] Critical associated metrics on symplectic manifolds, *Contemp. Math.*, **51**, 23–29.
- Blair, D. E., Ishihara, S. and Ludden, G. D.
 [1978] Projectable almost complex contact structures, *Kōdai Math. J.*, **1**, 75–84.
- Blair, D. E., Kim, J.-S. and Tripathi, M. M.
 [2005] On the concircular curvature tensor of a contact metric manifold, *J. Korean Math. Soc.*, **42**, 883–892.
- Blair, D. E. and Korkmaz, B.
 [2009] Special directions in complex contact manifolds, *Beiträge Algebra Geom.*, **50**, 309–325.
- Blair, D. E., Korkmaz, B. and Vrancken, L.
 [2000] The Calabi (Veronese) imbeddings as integral submanifolds of $\mathbb{C}P^{2n+1}$, *Glasgow Math. J.*, **42**, 183–193.
- Blair, D. E. and Koufogiorgos, Th.
 [1994] When is the tangent sphere bundle conformally flat?, *J. Geom.*, **49**, 55–66.
- Blair, D. E., Koufogiorgos, Th. and Papantoniou, B.J.
 [1995] Contact metric manifolds satisfying a nullity condition, *Israel J. Math.*, **91**, 189–214.
- Blair, D. E., Koufogiorgos, Th. and Sharma, R.
 [1990] A classification of 3-dimensional contact metric manifolds with $Q\phi = \phi Q$, *Kōdai Math. J.*, **13**, 391–401.
- Blair, D. E. and Ledger, A. J.
 [1986] Critical associated metrics on contact manifolds II, *J. Austral. Math. Soc.*, (Series A) **41**, 404–410.
- Blair, D. E. and Oubina, J. A.
 [1990] Conformal and related changes of metric on the product of two almost contact metric manifolds, *Publicacions Matemàtiques*, **34**, 199–207.
- Blair, D. E. and Perrone, D.
 [1992] A variational characterization of contact metric manifolds with vanishing torsion, *Canad. Math. Bull.*, **35**, 455–462.
 [1995] Second variation of the “total scalar curvature” on contact manifolds, *Canad. Math. Bull.*, **38**, 16–22.

- [1998] Conformally Anosov flows in contact metric geometry, *Balkan J. Geom. and App.*, **3**, 33–46.
- Blair, D. E. and Sharma, R.
- [1990a] Generalization of Myers' theorem on a contact manifold, *Ill. J. Math.*, **34**, 837–844.
- [1990b] Three dimensional locally symmetric contact metric manifolds, *Bollettino U.M.I.*, (7) **4-A**, 385–390.
- Blair, D. E., Showers, D. K. and Yano, K.
- [1976] Nearly Sasakian structures, *Kōdai Math. Sem. Rep.*, **27**, 175–180.
- Blair, D. E. and Vanhecke, L.
- [1987a] New characterizations of ϕ -symmetric spaces, *Kōdai Math. J.*, **10** (1987), 102–107.
- [1987b] Symmetries and ϕ -symmetric spaces, *Tōhoku Math. J.*, **39** (1987), 373–383.
- [1987c] Volume-preserving ϕ -geodesic symmetries, *C. R. Math. Rep. Acad. Sci. Canada*, **IX**, 31–36.
- [1987d] Geodesic spheres and Jacobi vector fields on Sasakian space forms, *Proc. Royal Soc. Edinburgh*, **105A**, 17–22.
- Boeckx, E.
- [1999] A class of locally ϕ -symmetric contact metric spaces, *Arch. Math.*, **72**, 466–472.
- [2000] A full classification of contact metric (k, μ) -spaces, *Ill. J. Math.*, **44**, 212–219.
- [2003] When are the tangent sphere bundles of a Riemannian manifold reducible? *Trans. Amer. Math. Soc.*, **355**, 2885–2903.
- [2005] A case for curvature: the unit tangent bundle, *Complex, Contact and Symmetric Manifolds*, Birkhäuser, Boston, 15–26.
- [2006] Contact-homogeneous locally φ -symmetric manifolds, *Glasgow Math. J.*, **48**, 93–109.
- Boeckx, E., Bueken, P. and Vanhecke, L.
- [1999] ϕ -symmetric contact metric spaces, *Glasgow Math. J.*, **41**, 409–416.
- Boeckx, E. and Cho J. T.
- [2006] Locally symmetric contact metric manifolds, *Monatsh. Math.*, **148**, 269–281.
- Boeckx, E., Perrone, D. and Vanhecke, L.
- [1998] Unit tangent sphere bundles and two-point homogeneous spaces, *Period. Math. Hungar.*, **36**, 79–95.
- Boeckx, E. and Vanhecke, L.
- [1997] Characteristic reflections on unit tangent sphere bundles, *Houston J. Math.*, **23**, 427–448.

Boothby, W. M.

- [1961] Homogeneous complex contact manifolds, *Proc. Symp. Pure Math.* III, *Amer. Math. Soc.*, 144–154.
- [1962] A note on homogeneous complex contact manifolds, *Proc. Amer. Math. Soc.*, **13**, 276–280.
- [1969] Transitivity of the automorphisms of certain geometric structures, *Trans. Amer. Math. Soc.*, **137**, 93–100.

Boothby, W. M. and Wang, H. C.

- [1958] On contact manifolds, *Ann. of Math.*, **68**, 721–734.

Borisenko, A. A. and Yampol'skii, A. L.

- [1987a] The sectional curvature of the Sasaki metric of T_1M^n , *Ukr. Geom. Sb.*, **30**, 10–17; English translation: *J. Sov. Math.*, **51** (1990), 2503–2508.
- [1987b] On the Sasaki metric of the normal bundle of a submanifold in a Riemannian space, *Mat. Sb.*, **134**(**176**); English translation: *Math. USSR Sb.*, **62** (1989), 157–175.

Borrelli, V. Chen, B.-Y. and Morvan, J.-M.

- [1995] Une caractérisation géométrique de la sphère de Whitney, *C. R. Acad. Sci.*, Paris **321**, 1485–1490.

Bouche, T.

- [1990] La cohomologie coeffective d'une variété symplectique, *Bull. Sc. Math.*, **114**, 115–122.

Bourgeois, F.

- [2002] Odd dimensional tori are contact manifolds, *International Mathematics Research Notices*, no. 30, 1571–1574.

Boyer, C. P. and Galicki, K.

- [1997] The twistor space of a 3-Sasakian manifold, *Internat. J. Math.*, **8**, 31–60.
- [2000] On Sasakian-Einstein geometry, *Internat. J. Math.*, **11**, 873–909.
- [2001] Einstein manifolds and contact geometry, *Proc. Amer. Math. Soc.*, **129**, 2419–2430.
- [2008] *Sasakian Geometry*, Oxford University Press, Oxford.

Boyer, C. P., Galicki, K. and Kollár, J.

- [2005] Einstein metrics on spheres, *Ann. of Math.*, **162**, 557–580.

Boyer, C. P., Galicki, K. and Mann, B. M.

- [1993] Quaternionic reduction and Einstein manifolds, *Communications in Analysis and Geometry*, **1**, 229–279.
- [1994] The geometry and topology of 3-Sasakian manifolds, *J. Reine Angew. Math.* **455**, 183–220.

Boyer, C. P., Galicki, K. and Matzeu, P.

- [2006] On eta-Einstein Sasakian geometry, *Comm. Math. Phys.*, **262**, 177–208.

Boyer, C. P., Galicki, K. and Nakamaye, M.

- [2003a] Sasakian geometry, homotopy spheres and positive Ricci curvature, *Topology*, **42**, 981–1002.
 [2003b] On the geometry of Sasakian-Einstein 5-manifolds, *Math. Ann.*, **325**, 485–524.

Boyer, C. P., Galicki, K. and Simanca, S.

- [2008] Canonical Sasakian metrics, *Comm. Math. Phys.*, **279**, 705–733.
 [2009] The Sasaki cone and extremal Sasakian metrics, *Riemannian Topology and Geometric Structures on Manifolds*, Birkhäuser, Boston, 263–290.

Brieskorn, E. and Van de Ven, A.

- [1968] Some complex structures on products of homotopy spheres, *Topology*, **7**, 389–393.

Bryant, R. L.

- [1982] Conformal and minimal immersions of compact surfaces into the 4-sphere, *J. Differential Geometry*, **17**, 455–473.

Bueken, P. and Vanhecke, L.

- [1988a] Geometry and symmetry on Sasakian manifolds, *Tsukuba J. Math.*, **12**, 403–422.
 [1988b] Curvature characterizations in contact geometry, *Riv. Math. Univ. Parma.*, **14**, 303–313.
 [1989] Reflections in K-contact geometry, *Math. Rep. Toyama Univ.*, **12**, 41–49.
 [1993] Reflections with respect to submanifolds in contact geometry, *Archivum Math.*, **29**, 43–57.

Calabi, E.

- [1953a] Metric Riemann surfaces, *Contributions to the Theory of Riemann Surfaces*, Princeton University Press, Princeton, 77–85.
 [1953b] Isometric imbeddings of complex manifolds, *Ann. of Math.*, **58**, 1–23.
 [1954] The space of Kähler metrics, *Proc. Internat. Congress Math.*, Amsterdam, 206–207.
 [1982] Extremal Kähler metrics, *Seminar on Differential Geometry*, Annals of Math. Studies **102**, Princeton University Press, Princeton, 259–290.
 [1985] Extremal Kähler metrics II, *Differential Geometry and Complex Analysis*, Springer, Berlin, 95–114.

Calabi, E. and Eckmann, B.

- [1953] A class of compact complex manifolds which are not algebraic, *Ann. of Math.*, **58**, 494–500.

Calvaruso, G.

- [2000] Einstein-like and conformally flat contact metric three-manifolds, *Balkan J. Geom. and Appl.*, **5**, 17–36.
- [2005] Contact metric geometry of the unit tangent sphere bundle, *Complex, Contact and Symmetric Manifolds*, Birkhäuser, Boston, 41–57.

Calvaruso, G. and Perrone, D.

- [2000] Torsion and homogeneity on contact metric three-manifolds, *Ann. Mat. Pura Appl.*, (4) **178**, 271–285.

Calvaruso, G., Perrone, D. and Vanhecke, L.

- [1999] Homogeneity on three-dimensional contact metric manifolds, *Israel J. Math.*, **114**, 301–321.

Cappelletti Montano, B.

- [to appear] The foliated structure of contact metric (κ, μ) -spaces.

Cappelletti Montano, B. and Di Terlizzi, L.

- [2008] Contact metric (κ, μ) -spaces as bi-Legendrian manifolds, *Bull. Austral. Math. Soc.*, **77**, 373–386.

Capursi, M.

- [1984] Some remarks on the product of two almost contact manifolds, *An. Sti. Univ. “Al. I. Cuza”, XXX*, 75–79.

Cartan, E.

- [1983] *Geometry of Riemannian spaces*, Math. Sci. Press, Brookline, MA.

Chen, B.-Y.

- [1997a] Interaction of Legendre curves and Lagrangian submanifolds, *Israel J. Math.*, **99**, 69–108.
- [1997b] Complex extensors and Lagrangian submanifolds in complex Euclidean spaces, *Tôhoku Math. J.*, **49**, 277–297.
- [1998] Intrinsic and extrinsic structures of Lagrangian surfaces in complex space forms, *Tsukuba J. Math.*, **22**, 657–680.
- [2000] Riemannian submanifolds, *Handbook of Differential Geometry*, Vol. 1, 153–418.

Chen, B.-Y. and Houh, C.-S.

- [1979] Totally real submanifolds of a quaternion projective space, *Ann. Mat. Pura Appl.*, **120**, 185–199.

Chen, B.-Y. and Ogiue, K.

- [1973] Some characterizations of complex space forms, *Duke Math. J.*, **40**, 797–799.
- [1974a] On totally real submanifolds, *Trans. Amer. Math. Soc.*, **193**, 257–266.
- [1974b] Two theorems on Kaehler manifolds, *Michigan Math. J.*, **21**, 225–229.

Chen, B.-Y. and Vanhecke, L.

- [1989] Isometric, holomorphic and symplectic reflections, *Geometriae Dedicata*, **29**, 259–277.

Chern, S. S.

- [1953] Pseudo-groupes continus infinis, colloques Internationaux du C. N. R. S., Strasbourg, 119–136.

Chern, S. S., Cowen, M. J. and Vitter III, A. L.

- [1974] Frenet frames along holomorphic curves, *Value Distribution Theory*, part A, Dekker, New York, 191–203.

Chern, S. S., do Carmo, M. P. and Kobayashi, S.

- [1970] Minimal submanifolds of a sphere with second fundamental form of constant length, *Functional Analysis and Related Fields*, Springer, Berlin, 59–75.

Chern, S. S. and Hamilton, R. S.

- [1985] On Riemannian metrics adapted to three-dimensional contact manifolds, *Lecture Notes in Mathematics*, Vol. 1111, Springer, Berlin, 279–308.

Chevalley, C.

- [1946] *Theory of Lie Groups*, Princeton University Press, Princeton.

Chinea, D.

- [1985] Invariant submanifolds of a quasi-K-Sasakian manifold, *Riv. Mat. Univ. Parma*, **11**, 25–29.

Chinea, D., de Leon, M. and Marrero, J. C.

- [1993] Topology of cosymplectic manifolds, *J. Math. Pures Appl.*, **72**, 567–591.
 [1995] Coeffective cohomology on almost cosymplectic manifolds, *Bull. Sci. Math.*, **119**, 3–20.
 [1997] Spectral sequences on Sasakian and cosymplectic manifolds, *Houston J. Math.*, **23**, 631–649.

Cho, J. T.

- [1999] A new class of contact Riemannian manifolds, *Israel J. Math.*, **109**, 299–318.
 [2003] A class of contact Riemannian manifolds whose associated CR-structures are integrable, *Publ. Math. Debrecen*, **63**, 193–211.

Cordero, L.A., Fernández, M. and Gray, A.

- [1986] Symplectic manifolds with no Kähler structure, *Topology*, **25**, 375–380.

Cordero, L.A., Fernández, M. and de Leon, M.

- [1985] Examples of compact non-Kähler almost Kähler manifolds, *Proc. Amer. Math. Soc.*, **95**, 280–286.

Davidov, J. and Muškarov, O.

- [1990] Twistor spaces with Hermitian Ricci tensor, *Proc. Amer. Math. Soc.*, **109**, 1115–1120.

Deng, S.

- [1991] The second variation of the Dirichlet energy on contact manifolds, *Kōdai Math. J.*, **14**, 470–476.

Dillen, F. and Vrancken, L.

- [1989] C-totally real submanifolds of $S^7(1)$ with non-negative sectional curvature, *Math. J. Okayama Univ.*, **31**, 227–242.
- [1990] C-totally real submanifolds of Sasakian space forms, *J. Math. Pures Appl.*, **69**, 85–93.

Dolbeault, S.

- [1977] Moving frames in Hermitian geometry, *Proc. Symp. Pure Math.* XXX, *Amer. Math. Soc.*, 103–106.

Dombrowski, P.

- [1962] On the geometry of the tangent bundle, *J. reine und angew. Math.*, **210**, 73–88.

Douady, A.

- [1982/83] Noeuds et structure de contact en dimension 3, *Astérisque*, **105–106**, 129–148.

Draghici, T.

- [1994] On the almost Kähler manifolds with Hermitian Ricci tensor, *Houston J. Math.*, **20**, 293–298.
- [1999] Almost Kähler 4-manifolds with J-invariant Ricci tensor, *Houston J. Math.*, **25**, 133–145.

Dragomir, S.

- [1995] On pseudohermitian immersions between strictly pseudoconvex CR manifolds, *Amer. J. Math.*, **117**, 169–202.

Ebin, D.

- [1970] The manifold of Riemannian metrics, *Proc. Symposia Pure Math.* XV, *Amer. Math. Soc.*, 11–40.

Ejiri, N.

- [1982] Totally real minimal immersions of n -dimensional real space forms into n -dimensional complex space forms, *Proc. Amer. Math. Soc.*, **84**, 243–246.

Eliashberg, Y.

- [1989] Classification of overtwisted contact structures on 3-manifolds, *Invent. Math.*, **98**, 623–637.

Elisahberg, Y. and Thurston, W.

[1998] Confoliations, University Lecture Series **13**, *Amer. Math. Soc.*

Endo, H.

[1985] Invariant submanifolds in a contact Riemannian manifold, *Tensor* (N.S.), **42**, 86–89.

[1986] Invariant submanifolds in contact metric manifolds, *Tensor* (N.S.), **43**, 83–86.

Fernández M. and Gray, A.

[1986] The Iwasawa manifold, *Lecture Notes in Mathematics*, Vol. 1209, Springer, Berlin, 157–159.

[1990] Compact symplectic solvmanifolds not admitting complex structures, *Geometria Dedicata*, **34**, 295–299.

Fernández M., Ibáñez, R. and de Leon, M.

[1997] A Nomizu's theorem for the coeffective cohomology, *Math. Z.*, **226**, 11–23.

[1998] Coeffective and de Rham cohomologies on almost contact manifolds, *Differential Geom. Appl.*, **8**, 285–303.

Fernández M., de Leon, M. and Saralegi, M.

[1996] A six-dimensional compact symplectic solvmanifold without Kähler structures, *Osaka J. Math.*, **33**, 1–19.

Fetcu, D. and Oniciuc, C.

[to appear] A note on integral C -parallel submanifolds in $S^7(c)$.

Foreman, B.

[1996] Variational Problems on Complex Contact Manifolds with Applications to Twistor Space Theory, thesis, Michigan State University.

[1999] Three-dimensional complex homogeneous complex contact manifolds, *Balkan J. Geom. and Appl.*, **4**, 53–67.

[2000a] Boothby-Wang fibrations on complex contact manifolds, *Differential Geom. Appl.* **13**, 179–196.

[2000b] Complex contact manifolds and hyperkähler geometry, *Kōdai Math. J.*, **23**, 12–26.

[2002a] Decompositions and the complex contact structures of $Sl(2, \mathbb{C})$, *Kyungpook Math. J.*, **42**, 351–367.

[2002b] Curvature characterizations of twistor spaces over four-dimensional Riemannian manifolds, *Kōdai Math. J.*, **25**, 167–190.

[2006] Complex contact Lie groups and generalized complex Heisenberg groups, *Differential Geom. Appl.*, **24**, 443–446.

[2010] Discrete groups and the complex contact geometry of $Sl(2, \mathbb{C})$, *Trans. Amer. Math. Soc.*, **362**, 4191–4200.

Fujitani, T.

- [1966] Complex-valued differential forms on normal contact Riemannian manifolds, *Tôhoku Math. J.*, **18**, 349–361.

Galicki, K. and Salamon, S.

- [1996] Betti numbers of 3-Sasakian manifolds, *Geometriae Dedicata*, **63**, 45–68.

Gauntlett, J.P., Martelli, D., Sparks, J. and Waldram, D.

- [2004] Sasaki-Einstein metrics $S^2 \times S^3$, *Adv. Theor. Math. Phys.*, **8**, 711–734.

Geiges, H.

- [1997a] Constructions of contact manifolds, *Proc. Camb. Phil. Soc.*, **121**, 455–464.
 [1997b] Normal contact structures on 3-manifolds, *Tôhoku Math. J.*, **49**, 415–422.
 [2006] Contact geometry, *Handbook of Differential Geometry II*, North-Holland, Amsterdam, 315–382.
 [2008] *An Introduction to Contact Topology*, Cambridge University Press, Cambridge.

Geiges, H. and Gonzalo, J.

- [1995] Contact geometry and complex surfaces, *Invent. Math.*, **121**, 147–209.
 [1997] Contact circles on 3-manifolds, *J. Differential Geometry*, **46**, 236–286.

Geiges, H. and Thomas, C. B.

- [1995] Hypercontact manifolds, *J. London Math. Soc.*, **51**, 342–352.

Ghosh, A., Koufogiorgos, Th. and Sharma, R.

- [2001] Conformally flat contact metric manifolds, *J. Geom.*, **70**, 66–76.

Ghosh, A. and Sharma, R.

- [1999] On contact strongly pseudo-convex integrable *CR* manifolds, *J. Geom.*, **66**, 116–122.

Ghys É.

- [1987] Flots d'Anosov dont les feuilletages stables sont différentiables, *Ann. Sc. École Norm. Sup.*, **20**, 251–270.
 [1995] Holomorphic Anosov systems, *Invent. Math.*, **119**, 585–614.

Ginzburg, V.

- [1995] An embedding $S^{2n-1} \rightarrow R^{2n}$, $2n - 1 \geq 7$, whose Hamiltonian flow has no periodic trajectories, *International Math. Res. Notices*, 83–98.

Goldberg, S. I.

- [1962] *Curvature and Homology*, Academic Press, New York.
 [1967] Rigidity of positively curved contact manifolds, *J. London Math. Soc.*, **42**, 257–263.
 [1968a] Totally geodesic hypersurfaces of Kaehler manifolds, *Pacific J. Math.*, **27**, 275–281.

- [1968b] On the topology of compact contact manifolds, *Tôhoku Math. J.*, **20**, 106–110.
 - [1969] Integrability of almost Kähler manifolds, *Proc. Amer. Math. Soc.*, **21**, 96–100.
 - [1986] Nonnegatively curved contact manifolds, *Proc. Amer. Math. Soc.*, **96**, 651–656.
- Goldberg, S. I. and Moskal, E.
- [1976] The axiom of spheres in Kaehler geometry, *Kōdai Math. Sem. Rep.*, **27**, 188–192.
- Gompf, R.
- [1994] Some new symplectic 4-manifolds, *Turkish J. Math.*, **18**, 7–15.
 - [1995] A new construction of symplectic manifolds, *Ann. of Math.*, **142**, 527–595.
- González-Dávila, J. C., González-Dávila, M. C. and Vanhecke, L.
- [1995] Reflections and isometric flows, *Kyungpook Math. J.*, **35**, 113–144.
- Gonzalo, J.
- [1987] Branched covers and contact structures, *Proc. Amer. Math. Soc.*, **101**, 347–352.
- Gouli-Andreou, F. and Sharma, R.
- [2003] A class of conformally flat contact metric 3-manifolds, *Results Math.*, **43**, 114–120.
- Gouli-Andreou, F. and Tsolakidou, N.
- [2004] Conformally flat contact metric manifolds with $Q\xi = \rho\xi$, *Beiträge Algebra Geom.* **45**, 103–115.
- Gouli-Andreou, F. and Xenos, Ph. J.
- [1998] On 3-dimensional contact metric manifolds with $\nabla_{\xi}\tau = 0$, *J. Geom.*, **62**, 154–165.
 - [1999] Two classes of conformally flat contact metric 3-manifolds, *J. Geom.*, **64**, 80–88.
- Gray, A. and Hervella, L. M.
- [1980] The sixteen classes of almost Hermitian manifolds and their linear invariants, *Ann. Mat. Pura Appl.*, **123**, 35–58.
- Gray, J.
- [1959] Some global properties of contact structures, *Ann. of Math.*, **69**, 421–450.
- Greenfield, S.
- [1968] Cauchy-Riemann equations in several variable, *Ann. Scuola Norm. Sup. Pisa*, **22**, 275–314.

Gromov, M.

- [1971] A topological technique for the construction of solutions of differential equations and inequalities, *Internat. Congr. Math. (Nice 1970)* **2**, Gauthier-Villars, Paris, 221–225.
- [1985] Pseudo holomorphic curves in symplectic manifolds, *Invent. Math.*, **82**, 307–347.

Guan, D.

- [1994] Examples of compact holomorphic symplectic manifolds which admit no Kähler structure, *Geometry and Analysis on Complex Manifolds*, World Scientific, Singapore, 63–74.

Hadjar, A.

- [1998] Sur les structures de contact du tore T^5 , *L'Enseign. Math.*, **44**, 91–93.

Haefliger, A.

- [1971] Lectures on the theorem of Gromov, *Lecture Notes in Mathematics*, Vol. 209 Springer, Berlin, 128–141.

Han, D.-S. and Yim, J.-W.

- [1998] Unit vector fields on spheres which are harmonic maps, *Math. Z.*, **227**, 83–92.

Harada, M.

- [1973a] On Sasakian submanifolds, *Tôhoku Math. J.*, **25**, 103–109.
- [1973b] On Sasakian submanifolds II, *Bull. Tokyo Gakugei Univ.*, **25**, 19–23.
- [1974a] On Sasakian manifolds satisfying the axiom of ϕ -holomorphic 2-spheres, *Bull. Tokyo Gakugei Univ.*, **26**, 1–3.
- [1974b] On Sasakian manifolds satisfying the axiom of anti- ϕ -holomorphic planes or 2-spheres, *Bull. Tokyo Gakugei Univ.*, **26**, 22–25.

Hasegawa, I. and Seino, M.

- [1981] Some remarks on Sasakian geometry, *J. Hokkaido Univ. Education*, **32**, 1–7.

Hatakeyama, Y.

- [1962] On the existence of Riemann metrics associated with a 2-form of rank $2r$, *Tôhoku Math. J.*, **14**, 162–166.
- [1963] Some notes on differentiable manifolds with almost contact structures, *Tôhoku Math. J.*, **15**, 176–181.
- [1966] Some notes on the group of automorphisms of contact and symplectic structures, *Tôhoku Math. J.*, **18**, 338–347.

Hatakeyama, Y., Ogawa, Y. and Tanno, S.

- [1963] Some properties of manifolds with contact metric structures, *Tôhoku Math. J.*, **15**, 42–48.

Hilbert, D.

- [1915] Die Grundlagen der Physik, *Nachr. Ges. Wiss. Göttingen*, 395–407.

Hille, E.

- [1976] *Ordinary Differential Equations in the Complex Domain*, Wiley, New York.

Hitchin, N. J.

- [1987] The self-duality equations on a Riemann surface, *Proc. London Math. Soc.* **55**, 59–126.

Hofer, H.

- [1993] Pseudoholomorphic curves in symplectizations with applications to the Weinstein conjecture in dimension three, *Invent. Math.*, **114**, 515–563.

Hofer, H. and Viterbo, C.

- [1988] The Weinstein conjecture in cotangent bundles and related results, *Ann. Sc. Norm. Sup. Pisa*, **15**, 411–445.

Hofer, H. and Zehnder, E.

- [1987] Periodic solutions on hypersurfaces and a result by C. Viterbo, *Invent. Math.*, **90**, 1–9.

Holubowicz, R. and Mozgawa, W.

- [1998] On compact non-Kählerian manifolds admitting an almost Kähler structure, *Rend. Cir. Mat. Palermo, Serie II, Suppl.* **54**, 53–57.

Houh, C.-S.

- [1973] Some totally real minimal surfaces in CP^2 , *Proc. Amer. Math. Soc.*, **40**, 240–244.
 [1976] On the holonomy group of a normal complex almost contact manifold, *Kōdai Math. Sem. Rep.*, **28**, 72–77.

Hsu, C. J.

- [1960] On some structures which are similar to the quaternion structure, *Tôhoku Math. J.*, **12**, 403–428.

Huygens, C.

- [1690] *Traité de la Lumière*, Vander Aa, Leiden.

Ianus, S.

- [1972] Sulle varietà di Cauchy-Riemann, *Rend. dell'Accademia di Scienze Fisiche e Matematiche*, Napoli, XXXIX, 191–195.

Ishihara, S.

- [1973] Quaternion Kähler manifolds and fibered Riemannian spaces with Sasakian 3-structure, *Kōdai Math. Sem. Rep.*, **25**, 321–329.
 [1974] Quaternion Kählerian manifolds, *J. Differential Geometry*, **9**, 483–500.

Ishihara, S and Konishi M.

- [1979] Real contact 3-structure and complex contact structure, *Southeast Asian Bulletin of Math.*, **3**, 151–161.
 [1980] Complex almost contact manifolds, *Kōdai Math. J.*, **3**, 385–396.

- [1982] Complex almost contact structures in a complex contact manifold, *Kōdai Math. J.*, **5**, 30–37.
- Itoh, M.
 [1997] Odd dimensional tori and contact structure, *Proc. Japan Acad.*, **72**, 58–59.
- Janssens, D. and Vanhecke, L.
 [1981] Almost contact structures and curvature tensors, *Kōdai Math. J.*, **4**, 1–27.
- Jayne, N.
 [1992] Legendre Foliations on Contact Metric Manifolds, thesis, Massey University.
 [1994] A note on the sectional curvature of Legendre foliations, *Yokohama Math. J.*, **41**, 153–161.
 [1998] Contact metric structures and Legendre foliations, *New Zealand J. Math.*, **27**, 49–65.
- Jelonek, W.
 [1996] Some simple examples of almost Kähler non-Kähler structures, *Math. Ann.*, **305**, 639–649.
- Jerison, D. and Lee, J. M.
 [1984] A subelliptic, nonlinear eigenvalue problem and scalar curvature on CR manifolds, *Contemp. Math.*, **27**, 57–63.
- Jiménez, J. A. and Kowalski, O.
 [1993] The classification of ϕ -symmetric Sasakian manifolds, *Mh. Math.*, **115**, 83–98.
- Kashiwada, T.
 [1971] A note on a Riemannian space with Sasakian 3-structure, *Natural Sci. Rep. Ochanomizu Univ.*, **22**, 1–2.
 [1998] A note on Hitchin's lemma, *Tensor (N.S.)*, **60**, 323–326.
 [2001] On a contact 3-structure, *Math. Z.*, **238**, 829–832.
- Kassabov, O.
 [1982] On the axiom of spheres in Kaehler geometry, *C. R. Acad. Bulgare Sci.*, **35**, 303–306.
- Kenmotsu, K.
 [1969] Invariant submanifolds in a Sasakian manifold, *Tôhoku Math. J.*, **21**, 495–500.
 [1972] A class of almost contact Riemannian manifolds, *Tôhoku Math. J.*, **24**, 93–103.
- Kirchberg, K.-D.
 [2004] Some integrability conditions for almost Kähler manifolds, *J. Geom. Phys.*, **49**, 101–115.

Kirichenko, V. F.

- [2001] On the geometry of Lagrangian submanifolds, *Mathematical Notes*, **69**, 32–45; *Matematicheskie Zametki*, **69**, 36–51.

Kirichenko, V. F. and Borisovski, I. P.

- [1998] Integral manifolds of contact distributions, *Sbornik: Mathematics*, **189**, 1855–1870; *Matematicheskiĭ Sbornik*, **189**, 119–134.

Kobayashi, S.

- [1956] Principal fiber bundles with the 1-dimensional toroidal group, *Tôhoku Math. J.*, **8**, 29–45.
 [1959] Remarks on complex contact manifolds, *Proc. Amer. Math. Soc.*, **10**, 164–167.
 [1963] Topology of positively pinched Kaehler manifolds, *Tôhoku Math. J.*, **15**, 121–139.

Kobayashi, S. and Nomizu, K.

- [1963–69] *Foundations of Differential Geometry*, I, II, Wiley-Interscience, New York.

Kobayashi, S. and Wu H.

- [1983] *Complex Differential Geometry*, Birkhäuser, Basel.

Kodaira, K.

- [1964] On the structure of compact complex analytic surfaces, I, *Amer. J. Math.*, **86**, 751–798.

Kollár, J.

- [2007] Einstein metrics on connected sums of $S^2 \times S^3$, *J. Differential Geometry*, **75**, 259–272.
 [2009] Positive Sasakian structures on 5-manifolds, *Riemannian Topology and Geometric Structures on Manifolds*, Birkhäuser, Boston, 93–117.

Kon, M.

- [1976] Invariant submanifolds in Sasakian manifolds, *Math. Ann.*, **219**, 277–290.

Konishi, M.

- [1973] Note on regular K-contact 3-structures, *Natural Sci. Rep. Ochanomizu Univ.*, **22**, 1–5.
 [1975] On manifolds with Sasakian 3-structure over quaternion Kaehler manifolds, *Kōdai Math. Sem. Rep.*, **26**, 194–200.

Konishi, M. and Funabashi, S.

- [1976] On Riemannian manifolds with Sasakian 3-structure of constant horizontal sectional curvature, *Kōdai Math. Sem. Rep.*, **27**, 362–366.

Korkmaz, B.

- [1998] A curvature property of complex contact metric structures, *Kyungpook Math. J.*, **38**, 473–488.

- [2000] Normality of complex contact manifolds, *Rocky Mountain J. Math.*, **30**, 1343–1380.
 - [2003] A nullity condition for complex contact metric manifolds, *J. Geom.*, **77**, 108–128.
- Kosmanek, E.
- [1964] Une propriété caractéristique des variétés Kähleriennes à courbure holomorphe constante, *C. R. Acad. Sc. Paris*, **259**, 705–708.
- Koufogiorgos, Th.
- [1997a] Contact Riemannian manifolds with constant ϕ -sectional curvature, *Tokyo J. Math.*, **20**, 13–22.
 - [1997b] Contact strongly pseudo-convex integrable CR metrics as critical points, *J. Geom.*, **59**, 94–102.
- Koufogiorgos, Th., Markellos, M. and Papantoniou, B. J.
- [2008] The harmonicity of the Reeb vector field on contact metric 3-manifolds, *Pac. Math. J.*, **234**, 325–344.
- Koufogiorgos, Th. and Tsihlias, C.
- [2000] On the existence of a new class of contact metric manifolds, *Canad. Math. Bull.*, **43**, 440–447.
 - [2003] Generalized (κ, μ) -contact metric manifolds with $\|\text{grad } \kappa\| = \text{constant}$, *J. Geom.* **78**, 83–91.
- Kowalski, O.
- [1971] Curvature of the induced Riemannian metric on the tangent bundle of a Riemannian manifold, *J. Reine Angew. Math.*, **250**, 124–129.
- Kowalski, O. and Vanhecke, L.
- [1984] A generalization of a theorem on naturally reductive homogeneous spaces, *Proc. Amer. Math. Soc.*, **91**, 433–435.
- Kowalski, O. and Wegrzynowski, S.
- [1987] A classification of five-dimensional ϕ -symmetric spaces, *Tensor (N.S.)*, **46**, 379–386.
- Kriegel, A. and Michor P. W.
- [1997] *The Convenient Setting of Global Analysis*, Math. Surveys and Monog. **53**, Amer. Math. Soc.
- Kuo, Y.-Y.
- [1970] On almost contact 3-structure, *Tôhoku Math. J.*, **22**, 325–332.
- Lawson, H. B.
- [1970] The Riemannian geometry of holomorphic curves, Proc. Carolina Conf. on Holomorphic Mappings and Minimal Surfaces, 86–107.
- LeBrun, C.
- [1991] On complete quaternionic-Kähler manifolds, *Duke Math. J.*, **63**, 723–743.

- [1995] Fano manifolds, contact structures and quaternionic geometry, *International J. Math.*, **6** (1995), 419–437.
- Libermann, P.
- [1959] Sur les automorphismes infinitésimaux des structures symplectiques et de structures de contact, Coll. Géom. Diff. Globale (Bruxelles 1958), Gauthier-Villars, Paris, 37–59.
- [1991] Legendre foliations on contact manifolds, *Differential Geom. Appl.*, **1**, 57–76.
- Libermann, P. and Marle, C.-M.
- [1987] *Symplectic Geometry and Analytical Mechanics*, D. Reidel, Dordrecht.
- Lie, S.
- [1890] *Theorie der Transformationgruppen*, Vol. 2, Leipzig, Teubner.
- Ludden, G. D., Okumura, M. and Yano, K.
- [1975] A totally real surface in CP^2 that is not totally geodesic, *Proc. Amer. Math. Soc.*, **53**, 186–190.
- Lutz, R.
- [1979] Sur la géométrie des structures de contact invariantes, *Ann. Inst. Fourier*, Grenoble **29**, 283–306.
- Lutz, R. and Meckert, C.
- [1976] Structures de contact sur certaines sphères exotiques, *C. R. Acad. Sc. Paris*, **282**, 591–593.
- Lychagin, V.
- [1977] On sufficient orbits of a group of contact diffeomorphisms, *Math. USSR Sb.*, **33** 223–242.
- Mac Lane, S.
- [1968] Geometrical Mechanics II, Lecture notes, University of Chicago.
- Marrero, J. C.
- [1992] The local structure of trans-Sasakian manifolds, *Ann. Mat. Pura Appl.*, **162**, 77–86.
- Marrero, J. C. and Padron, E.
- [1998] New examples of compact cosymplectic solvmanifolds, *Arch. Math.*, **34**, 337–345.
- Martinet, J.
- [1970] Sur les singularités des formes différentielles, *Ann. Inst. Fourier*, **20**, 95–178.
- [1971] Formes de contact sur les variétés de dimension 3, *Lecture Notes in Mathematics*, Vol. 209 Springer, Berlin, 142–163.
- McCarthy, J. D. and Wolfson, J. G.
- [1994] Symplectic normal connect sum, *Topology*, **33**, 729–764.

- McDuff, D.
 [1984] Examples of simply-connected symplectic non-Kählerian manifolds, *J. Differential Geom.*, **20**, 267–277.
- Milnor, J.
 [1976] Curvature of left invariant metrics on Lie groups, *Advances in Math.*, **21**, 293–329.
- Mitric, G.
 [1991] C. R.-structures on the unit sphere bundle in the tangent bundle of a Riemannian manifold, *Semin. Mec. Univ. Timis. Fac. Mat.*, **32**.
- Mitsumatsu, Y.
 [1995] Anosov flows and non-Stein symplectic manifolds, *Ann. Inst. Fourier*, **45**, 1407–1421.
- Monna, G.
 [1983] \mathfrak{h} -platitude des structures de contact, *Mh. Math.*, **95**, 221–227.
- Morimoto, A.
 [1963] On normal almost contact structures, *J. Math. Soc. Japan*, **15**, 420–436.
 [1964] On normal almost contact structure with a regularity, *Tôhoku Math. J.*, **16**, 90–104.
- Morimoto, S.
 [1992] Almost complex foliations and its application to contact geometry, *Natural Sci. Rep. Ochanomizu Univ.*, **43**, 11–23.
- Moroianu, A. and Semmelmann U.
 [1994] Kählerian Killing spinors, complex contact structures and twistor spaces, *Quaternionic Structures in Mathematics and Physics*, SISSA, Trieste, 225–231.
- Morrow, J. and Kodaira, K.
 [1971] *Complex Manifolds*, Holt, Rinehart and Winston, New York.
- Morvan, J.-M.
 [1983] On Lagrangian immersions, *Res. Notes in Math.*, Vol. 80, Pitman, Boston, 167–171.
- Moser, J.
 [1965] On the volume element on a manifold, *Trans. Amer. Math. Soc.*, **120**, 286–294.
- Moskal, E.
 [1966] Contact Manifolds of Positive Curvature, thesis, University of Illinois.
 [1977] On the tridegree of forms on f -manifolds, *Kōdai Math. Sem. Rep.*, **28** (1977), 115–128.

Musso, E. and Tricerri, F.

- [1988] Riemannian metrics on tangent bundles, *Ann. Mat. Pura Appl.*, **150**, 1–20.

Munteanu, G.

- [2004] *Complex Spaces in Finsler, Lagrange and Hamilton Geometries*, Kluwer, Dordrecht.

Muto, Y.

- [1974a] On Einstein metrics, *J. Differential Geometry*, **9**, 521–530.
 [1974b] Curvature and critical Riemannian metric, *J. Math. Soc. Japan*, **26**, 686–697.
 [1975] Critical Riemannian metrics, *Tensor* (N. S.), **29**, 125–133.

Myers, S. B.

- [1941] Riemannian manifolds with positive mean curvature, *Duke Math. J.*, **8**, 401–404.

Nagano, T.

- [1967] A problem on the existence of an Einstein metric, *J. Math. Soc. Japan*, **19**, 30–31.

Nakagawa, H. and Ogiue, K.

- [1976] Complex space forms immersed in complex space forms, *Trans. Amer. Math. Soc.*, **219**, 289–297.

Newlander, A. and Nirenberg, L.

- [1957] Complex analytic coordinates in almost complex manifolds, *Ann. of Math.*, **65**, 391–404.

Niederkrüger, K.

- [2006] The plastikstufe - a generalization of the overtwisted disk to higher dimensions, *Algebr. Geom. Topol.*, **6**, 2473–2508.

Nurowski, P. and Przanowski, M.

- [1999] A four dimensional example of Ricci flat metric admitting almost-Kähler non-Kähler structure, *Class. Quantum Grav.*, **16**, L9–L13.

Obata, M.

- [1965] Riemannian manifolds admitting a solution of a certain system of differential equations, *Proc. U.S.-Japan Seminar in Differential Geometry*, 101–114.

Ogiue, K.

- [1964] On almost contact manifolds admitting axiom of planes or axiom of free mobility, *Kōdai Math. Sem. Rep.*, **16**, 115–128.
 [1965] On fiberings of almost contact manifolds, *Kōdai Math. Sem. Rep.*, **17**, 53–62.
 [1972] Positively curved complex submanifolds immersed in a complex projective space, *J. Differential Geometry*, **7**, 603–606.

- [1974] Differential geometry of Kaehler submanifolds, *Advances in Math.*, **13** (1974), 73–114.
- [1976a] Positively curved complex submanifolds immersed in a complex projective space III, *J. Differential Geometry*, **11**, 613–615.
- [1976b] Positively curved totally real minimal submanifolds immersed in a complex projective space, *Proc. Amer. Math. Soc.*, **56**, 264–266.
- Okumura, M.
 - [1962a] Some remarks on space with a certain contact structure, *Tôhoku Math. J.*, **14**, 135–145.
 - [1962b] On infinitesimal conformal and projective transformations of normal contact spaces, *Tôhoku Math. J.*, **14**, 398–412.
 - [1964a] Certain almost contact hypersurfaces in Euclidean spaces, *Kôdai Math. Sem. Rep.*, **16**, 44–54.
 - [1964b] Certain almost contact hypersurfaces in Kaehlerian manifolds of constant holomorphic sectional curvatures, *Tôhoku Math. J.*, **16**, 270–284.
 - [1965] Cosymplectic hypersurfaces in Kaehlerian manifold of constant holomorphic sectional curvature, *Kôdai Math. Sem. Rep.*, **17**, 63–73.
 - [1966] Contact hypersurfaces in certain Kaehlerian manifolds, *Tôhoku Math. J.*, **18**, 74–102.
- Olszak, Z.
 - [1979] On contact metric manifolds, *Tôhoku Math. J.*, **31**, 247–253.
- O'Neill, B.
 - [1966] The fundamental equations of a submersion, *Michigan Math. J.*, **13**, 459–469.
- Oubina, J. A.
 - [1985] New classes of almost contact metric structures, *Publicationes Mathematicae Debrecen*, **32**, 187–193.
- Palais, R. S.
 - [1957] A Global Formulation of the Lie Theory of Transformation Groups, *Mem. Amer. Math. Soc.*, #22.
- Pang, M.-Y.
 - [1990] The structure of Legendre foliations, *Trans. Amer. Math. Soc.*, **320**, 417–455.
- Park, J.-S. and Oh, W. T.,
 - [1996] The Abbena-Thurston manifold as a critical point, *Canad. Math. Bull.*, **39**, 352–359.
- Pastore, A. M.
 - [1998] Classification of locally symmetric contact metric manifolds, *Balkan J. Geom. and Appl.*, **3**, 89–96.

Perrone, D.

- [1989] 5-dimensional contact manifolds with second Betti number $b_2 = 0$, *Tôhoku Math. J.*, **41**, 163–170.
- [1990] Torsion and critical metrics on contact three-manifolds, *Kôdai Math. J.*, **13**, 88–100.
- [1992a] Contact Riemannian manifolds satisfying $R(X, \xi) \cdot R = 0$, *Yokohama Math. J.*, **39**, 141–149.
- [1992b] Torsion tensor and critical metrics on contact $(2n+1)$ -manifolds, *Mh. Math.*, **114**, 245–259.
- [1994] Tangent sphere bundles satisfying $\nabla_\xi \tau = 0$, *J. Geom.*, **49**, 178–188.
- [1998] Homogeneous contact Riemannian three-manifolds, *Ill. J. Math.*, **42**, 243–256.
- [2000] Special directions on contact metric three-manifolds, *J. Geom.*, **69**, 180–191.
- [2002] Hypercontact metric three-manifolds, *C. R. Acad. Sci. Canada*, **24**, 97–101.
- [2003] Harmonic characteristic vector fields on contact metric three-manifolds, *Bull. Austral. Math. Soc.*, **67**, 305–315.
- [2004] Contact metric manifolds whose characteristic vector field is a harmonic vector field, *Differential Geom. Appl.*, **20**, 367–378.

Perrone, D. and Vanhecke, L.,

- [1991] Five-dimensional homogeneous contact and related problems, *Tôhoku Math. J.*, **43**, 243–248.

Pesin, Y.

- [2004] *Lectures on Partial Hyperbolicity and Stable Ergodicity*, European Mathematical Society.

Petkov, V. and Popov, G.

- [1995] On the Lebesgue measure of the periodic points of a contact manifold, *Math. Z.*, **218**, 91–102.

Priest, E. R.

- [1982] *Solar Magnetohydrodynamics*, D. Reidel, Dordrecht.

Rabinowitz, P. H.

- [1978] Periodic solutions of Hamiltonian systems, *Comm. Pure Appl. Math.*, **XXXI**, 157–184.

Reckziegel, H.

- [1979] On the eigenvalues of the shape operator of an isometric immersion into a space of constant curvature, *Math. Ann.*, **243**, 71–82.

Reeb, G.

- [1952] Sur certaines propriétés topologiques des trajectoires des systèmes dynamiques, *Mémoires de l'Acad. Roy. de Belgique, Sci. Ser. 2*, **27**, 1–62.

Ros, A.

- [1985a] Positively curved Kaehler submanifolds, *Proc. Amer. Math. Soc.*, **93**, 329–331.
- [1985b] A characterization of seven compact Kaehler submanifolds by holomorphic pinching, *Ann. of Math.*, **121**, 377–382.

Ros, A. and Verstraelen, L.

- [1984] On a conjecture of K. Ogiue, *J. Differential Geometry*, **19**, 561–566.

Ros, A. and Urbano, F.

- [1998] Lagrangian submanifolds of C^n with conformal Maslov form and the Whitney sphere, *J. Math. Soc. Japan*, **50**, 203–226.

Rukimbira, P.

- [1993] Some remarks on R-contact flows, *Ann. Global Anal. Geom.*, **11**, 165–171.
- [1995a] Topology and closed characteristics of K-contact manifolds, *Bull. Belg. Math. Soc.*, 349–356.
- [1995b] Vertical sectional curvature and K-contactness, *J. Geom.*, **53**, 163–166.
- [1995c] Chern-Hamilton's conjecture and K-contactness, *Houston J. Math.*, **21**, 709–718.
- [1998] A characterization of flat contact metric geometry, *Houston J. Math.*, **24**, 409–414.
- [2000] Spherical rigidity via contact dynamics, *Bull. Belg. Math. Soc.*, **7**, 563–569; Correction to: Spherical rigidity via contact dynamics, *Bull. Belg. Math. Soc.* **8** (2001), 147–153.
- [2009] The 1-nullity of Sasakian manifolds, *Riemannian Topology and Geometric Structures on Manifolds*, Birkhäuser, Boston, 153–159.

Salamon, S.

- [1982] Quaternionic Kähler manifolds, *Invent. Math.*, **67**, 143–171.

Sasaki, S.

- [1958] On the differential geometry of tangent bundles of Riemannian manifolds, *Tôhoku Math. J.*, **10**, 338–354.
- [1962] On the differential geometry of tangent bundles of Riemannian manifolds II, *Tôhoku Math. J.*, **14**, 146–155.
- [1964] A characterization of contact transformations, *Tôhoku Math. J.*, **16**, 285–290.
- [1965] Almost Contact Manifolds, Lecture Notes, Mathematical Institute, Tôhoku University, Vol. 1.
- [1972] Spherical space forms with normal contact metric 3-structure, *J. Differential Geometry*, **6** (1972), 307–315.

- [1985] Contact structures on Brieskorn manifolds (lecture Japan Mathematical Society 1975), *Shigeo Sasaki Selected Papers*, Kinokuniya, Tokyo, 349–363.
- Sasaki, S. and Hatakeyama, Y.
 [1961] On differentiable manifolds with certain structures which are closely related to almost contact structures II, *Tôhoku Math. J.*, **13**, 281–294.
 [1962] On differentiable manifolds with contact metric structures, *J. Math. Soc. Japan*, **14**, 249–271.
- Sasaki, S. and Hsu, C. J.
 [1976] On a property of Brieskorn manifolds, *Tôhoku Math. J.*, **28**, 67–78.
- Sasaki, S. and Takahashi, T.
 [1976] Almost contact structures on Brieskorn manifolds, *Tôhoku Math. J.*, **28**, 619–624.
- Sato, H.
 [1977] Remarks concerning contact manifolds, *Tôhoku Math. J.*, **29**, 577–584.
- Sekigawa, K.
 [1987] On some compact Einstein almost Kähler manifolds, *J. Math. Soc. Japan*, **39**, 677–684.
- Sharma, R.
 [1995] On the curvature of contact metric manifolds, *J. Geom.*, **53**, 179–190.
- Sharma, R. and Koufogiorgos, T.
 [1991] Locally symmetric and Ricci-symmetric contact metric manifolds, *Ann. Global Anal. Geom.*, **9**, 177–182.
- Shibuya, Y.
 [1978] On the existence of a complex almost contact structure, *Kôdai Math. J.*, **1**, 197–204.
 [1982] Some isospectral problems, *Kôdai Math. J.*, **5**, 1–12.
- Sikorav, J.-C.
 [1986] Non-existence de sous-variété lagrangienne exacte dans \mathbb{C}^n (d’après Gromov), *Aspects Dynamiques et Topologiques des Groupes Infinités de Transformation de la Mécanique*, Travaux en Cours **25**, Hermann, Paris, 95–110.
- Simons, J.
 [1968] Minimal varieties in Riemannian manifolds, *Ann. of Math.*, **88**, 62–105.
- Smolentsev, N. K.
 [1995] Critical associated metrics on a symplectic manifold, *Sibirsk. Mat. Zh.*, **36**, 409–418; English translation: *Siberian Math. J.*, **36**, 359–367.
- Sparks, J.
 [2009] New results in Sasaki-Einstein geometry, *Riemannian Topology and Geometric Structures on Manifolds*, Birkhäuser, Boston, 161–184.

Steenrod, N.

[1951] *Topology of Fiber Bundles*, Princeton University Press, Princeton.

Sternberg, S.

[1983] *Lectures on Differential Geometry*, Chelsea, New York.

Stong, R. E.

[1974] Contact manifolds, *J. Differential Geometry*, **9** (1974), 219–238.

Tachibana, S.

[1965] On harmonic tensors in compact Sasakian spaces, *Tôhoku Math. J.*, **17**, 271–284.

Tachibana, S. and Ogawa, Y.

[1966] On the second Betti number of a compact Sasakian space, *Natural Sci. Rep. Ochanomizu Univ.*, **17**, 27–32.

Tachibana, S. and Okumura, M.

[1962] On the almost complex structure of tangent bundles of Riemannian spaces, *Tôhoku Math. J.*, **14**, 156–161.

Tachibana, S. and Yu, W. N.

[1970] On a Riemannian space admitting more than one Sasakian structure, *Tôhoku Math. J.*, **22**, 536–540.

Takahashi, T.

[1977] Sasakian ϕ -symmetric spaces, *Tôhoku Math. J.*, **29**, 91–113.

Tanaka N.

[1976] On non-degenerate real hypersurfaces, graded Lie algebras and Cartan connections, *Japan J. Math.*, **2**, 131–190.

Tanno S.

[1963] Some transformations on manifolds with almost contact and contact metric structures II, *Tôhoku Math. J.*, **15**, 322–331.

[1965] A theorem on regular vector fields and its applications to almost contact structures, *Tôhoku Math. J.*, **17**, 235–238.

[1967a] Harmonic forms and Betti numbers of certain contact Riemannian manifolds, *J. Math. Soc. Japan*, **19**, 308–316.

[1967b] Locally symmetric K-contact Riemannian manifolds, *Proc. Japan Acad.*, **43**, 581–583.

[1968] The topology of contact Riemannian manifolds, *Illinois J. Math.*, **12**, 700–717.

[1969] Sasakian manifolds with constant φ -holomorphic sectional curvature, *Tôhoku Math. J.*, **21**, 501–507.

[1970] On the isometry group of Sasakian manifolds, *J. Math. Soc. Japan*, **22**, 579–590.

[1971] Killing vectors on contact Riemannian manifolds and fiberings related to the Hopf fibrations, *Tôhoku Math. J.*, **23**, 313–333.

- [1973] Constancy of holomorphic sectional curvature in almost Hermitian manifolds, *Tôhoku Math. J.*, **25**, 190–201.
 - [1978a] Some differential equations on Riemannian manifolds, *J. Math. Soc. Japan*, **30**, 509–531.
 - [1978b] Geodesic flows on C_L -manifolds and Einstein metrics on $S^3 \times S^2$, *Minimal Submanifolds and Geodesics*, Kaigai, Tokyo, 283–292.
 - [1988] Ricci curvatures of contact Riemannian manifolds, *Tôhoku Math. J.*, **40**, 441–448.
 - [1989] Variational problems on contact Riemannian manifolds, *Trans. Amer. Math. Soc.*, **314**, 349–379.
 - [1991] Pseudo-conformal invariants of type $(1, 3)$ of CR manifolds, *Hokkaido Math. J.*, **20**, 195–204.
 - [1992] The standard CR structure on the unit tangent bundle, *Tôhoku Math. J.*, **44**, 535–543.
 - [1996] Remarks on a triple of K-contact structures, *Tôhoku Math. J.*, **48**, 519–531.
- Tashiro, Y.
- [1963] On contact structures of hypersurfaces in complex manifolds I, *Tôhoku Math. J.*, **15**, 62–78.
 - [1969] On contact structures of tangent sphere bundles, *Tôhoku Math. J.*, **21**, 117–143.
- Taubes, C. H.
- [2007] The Seiberg–Witten equations and the Weinstein conjecture, *Geom. Topol.*, **11**, 2117–2202.
- Thomas, C. B.
- [1976] Almost regular contact structures, *J. Differential Geometry*, **11**, 521–533.
- Thurston, W.
- [1976] Some simple examples of symplectic manifolds, *Proc. Amer. Math. Soc.*, **55**, 467–468.
- Tischler, D.
- [1970] On fibering certain foliated manifolds over S^1 , *Topology*, **9**, 153–154.
- Tralle, A. and Oprea, J.
- [1997] Symplectic Manifolds with No Kähler Structure, *Lecture Notes in Mathematics*, Vol. 1661, Springer, Berlin.
- Udriste, C.
- [1969] Structures presque coquaternioniennes, *Bull. Math. Soc. Sci. Math. R. S. Roumanie*, **13**, 487–507.

Urbano, F.

- [1985] Totally real minimal submanifolds of a complex projective space, *Proc. Amer. Math. Soc.*, **93**, 332–334.

Vaisman, I.

- [1978] On the Sasaki-Hsu contact structure of the Brieskorn manifolds, *Tôhoku Math. J.*, **30**, 553–560.

van Coevering, C.

- [2009] Some examples of toric Sasaki-Einstein manifolds, *Riemannian Topology and Geometric Structures on Manifolds*, Birkhäuser, Boston, 185–232.

Vanhecke, L.

- [1988] Geometry in normal and tubular neighborhoods, *Rend. Sem. Fac. Sci. Univ. Cagliari*, LVIII (Supplement), 73–176.

Van Lindt D., Verheyen, P. and Verstraelen, L.

- [1986] Minimal submanifolds in Sasakian space forms, *J. Geom.*, **27**, 180–187.

Vernon, M.

- [1987] Contact hypersurfaces of a complex hyperbolic space, *Tôhoku Math. J.*, **39**, 215–222.

Verstraelen, L. and Vrancken, L.

- [1988] Pinching theorems for C-totally real submanifolds of Sasakian space forms, *J. Geom.*, **33**, 172–184.

Viterbo, C.

- [1987] A proof of Weinstein's conjecture in \mathbf{R}^{2n} , *Ann. Inst. Henri Poincaré*, **4**, 337–356.

Walters, P.

- [1975] Ergodic Theory-Introductory Lectures, *Lecture Notes in Mathematics*, Vol. 458, Springer, Berlin.

Watanabe, Y.

- [1980] Geodesic symmetries in Sasakian locally ϕ -symmetric spaces, *Kôdai Math. J.*, **3**, 48–55.

Watanabe, Y. and Fujita, H.

- [1988] A family of homogeneous Sasakian structures on $S^2 \times S^3$, *C. R. Math. Rep. Acad. Sci. Canada*, X, 57–61.

Watson, B.

- [1983] New examples of strictly almost Kähler manifolds, *Proc. Amer. Math. Soc.*, **88**, 541–544.

Weinstein, A.

- [1971] Symplectic manifolds and their Lagrangian submanifolds, *Advances in Math.*, **6**, 329–346.

- [1977] Lectures on Symplectic Manifolds, CBMS regional conference series in mathematics, **29**, Amer. Math. Soc., Providence.
 - [1978] Periodic orbits for convex hamiltonian systems, *Ann. of Math.*, **108**, 507–518.
 - [1979] On the hypothesis of Rabinowitz's periodic orbit theorems. *J. Diff. Eq.*, **33**, 353–358.
- Weyl H.
- [1939] *The Classical Groups*, Princeton Univ. Press, Princeton.
- Wolf, J. A.
- [1965] Complex homogeneous contact manifolds and quaternionic symmetric spaces, *J. Math. and Mech.*, **14**, 1033–1047.
 - [1968] A contact structure for odd dimensional spherical space forms, *Proc. Amer. Math. Soc.*, **19**, 196.
- Wood, C. M.
- [1995] Harmonic almost-complex structures, *Compositio Math.*, **99**, 183–212.
 - [1997] On the energy of a unit vector field, *Geometria Dedicata*, **64**, 319–330.
- Woodhouse, N.
- [1980] *Geometric Quantization*, Oxford Mathematical Monographs, Clarendon Press, Oxford.
- Yamada, T.
- [2005] A construction of compact pseudo-Kähler solvmanifolds with no Kähler structures, *Tsukuba, J. Math.*, **29**, 79–109.
- Yamaguchi, S. and Chūman, G.
- [1983] Critical Riemannian metrics on Sasakian manifolds, *Kōdai Math. J.*, **6**, 1–13.
- Yamaguchi, S., Kon, M. and Ikawa, T.
- [1976] C-totally real submanifolds, *J. Differential Geometry*, **11**, 59–64.
- Yamaguchi, S., Kon, M. and Miyahara, Y.
- [1976] A theorem on C-totally real minimal surface, *Proc. Amer. Math. Soc.*, **54**, 276–280.
- Yamato, K.
- [1990] Examples of non-Kähler symplectic manifolds, *Osaka, J. Math.*, **17**, 431–439.
- Yampol'skii, A. L.
- [1985] The curvature of the Sasaki metric of tangent sphere bundles, *Ukr. Geom. Sb.*, **28**, 132–145; English translation: *J. Sov. Math.*, **48** (1990), 108–117.

- Yano, K.
 [1965] *Differential Geometry on Complex and Almost Complex Spaces*, Pergamon, New York.
- Yano, K., Ishihara, S. and Konishi, M.
 [1973] Normality of almost contact 3-structure, *Tôhoku Math. J.*, **25**, 167–175.
- Yano, K. and Kon, M.
 [1983] *CR-submanifolds of Kählerian and Sasakian Manifolds*, Birkhäuser, Boston.
- Yano, K. and Mogi, I.
 [1955] On real representations of Kählerian manifolds, *Ann. of Math.*, **61**, 170–189.
- Yau, S.-T.
 [1974] Submanifolds with constant mean curvature I., *Amer. J. Math.*, **96**, 346–366.
- Ye, Y.-G.
 [1994] A note on complex projective threefolds admitting holomorphic contact structures, *Invent. Math.*, **115**, 311–314.
- Zeghib, A.
 [1995] Subsystems of Anosov systems, *Amer. J. Math.*, **117**, 1431–1448.
- Zheng, F.
 [2000] *Complex Differential Geometry*, Amer. Math. Soc., Providence.

Subject Index

(κ, μ) -manifold, 124
 (κ, μ, ν) -manifold, 212
 (ϕ, ξ, η) -structure, 43
*-Ricci form, 204
*-Ricci tensor, 118, 201
*-scalar curvature, 118, 201
 GH - sectional curvature, 253
 J -basis, 42
 J -invariant Ricci tensor, 201
 α -Sasakian, 99
 β -Kenmotsu, 99
 \mathcal{D} -homothetic deformation, 125, 141
 η -Einstein, 128, 131
 κ -nullity distribution, 128
 ϕ -basis, 44
 ϕ -geodesic, 143
 ϕ -geodesic symmetry, 143
 ϕ -sectional curvature, 137
 \mathcal{H} -homothetic deformation, 254
3- τ -manifold, 226
3-Sasakian manifold, 292

 ξ -sectional curvature, 219

Abbena metric, 5, 217
almost complex structure, 41
almost contact 3-structure, 291
almost contact metric 3-structure, 292
almost contact metric structure, 44
almost contact structure, 43
almost coquaternion structure, 291
almost Hermitian structure, 4, 42
almost hyper-Kähler, 293
almost Kähler, 4, 42
almost regular, 40
Anosov directions, 222
Anosov flow, 221
Anosov-like, 226
anti-invariant submanifold, 77
associated metric on a complex contact manifold, 236
associated metric on a contact manifold, 47
associated metric on a symplectic manifold, 3, 47

- axiom of ϕ -holomorphic 2-spheres, 155
- axiom of ϕ -holomorphic planes, 155
- axiom of anti- ϕ -holomorphic planes, 167
- axiom of antiholomorphic k -planes, 166
- axiom of holomorphic $2k$ -spheres, 155
- axiom of holomorphic planes, 155
- axiom of totally real k -planes, 166

- Boothby–Wang fibration, 36, 64
- Brieskorn manifold, 104

- C-parallel, 164
- C-totally real submanifold, 69, 159
- Calabi curves, 261
- Calabi embeddings, 154, 261
- characteristic class of S^1 -bundle, 21
- characteristic subbundle, 235
- characteristic vector field, 23
- characteristic vector field being Anosov, 222
- Codazzi equation, 152
- coeffective p -form, 109
- coisotropic, 8
- compatible metric, 44
- complex almost contact metric manifold, 236
- complex contact Lie group, 247
- complex contact manifold, 234
- complex contact metric manifold, 237
- complex contact space form, 253
- complex contact subbundle, 234
- complex geodesic flow, 273
- complex Sasakian manifold, 252
- complex symplectic manifold, 6, 245
- conformally Anosov flow, 227
- connection form, 20
- connection map, 170
- connection on a principal bundle, 19
- contact circle, 35
- contact distribution, 24, 69
- contact homogeneous, 146
- contact manifold, 23
- contact metric 3-structure, 292
- contact metric structure, 47
- contact Riemannian space form, 141
- contact Riemannian structure, 47
- contact structure in the wider sense, 25
- contact subbundle, 24, 69
- contact transformation, 25, 71
- contact type, 29
- contact Whitney sphere, 76
- cosymplectic, 95
- CR-dimension, 90
- CR-manifold, 90, 94
- CR-submanifold, 90
- curvature form, 20

- Darboux coordinate system, 24
- Douady portrait, 33

- effective p -form, 108
- extremal metric, 196
- extremal Sasakian structure, 196

- first normal space, 258
- Frenet curve, 164
- fundamental 2-form, 4, 42, 44

- gauge transformation, 216
- Gauss equation, 151
- generalized (κ, μ) -manifold, 127
- generalized Chern form, 203
- generalized Sasakian space form, 141
- generalized Tanaka connection, 212
- generalized Tanaka–Webster scalar curvature, 213
- generating function, 6
- geodesic flow, 177
- geodesic integrability, 167
- geodesic plane field, 120
- Goldberg conjecture, 202

- H-contact manifold, 211
- Hamiltonian vector field, 3

- harmonic vector field, 211
- Heisenberg group, 61
- Hermitian manifold, 42
- Hermitian metric, 3, 41
- Hermitian Ricci tensor, 201
- holomorphic Anosov flow, 267
- holomorphic Frenet curve, 258
- holomorphic integral submanifold, 237
- holomorphic Legendre curve, 237
- holomorphic special direction, 284
- holomorphic symplectic manifold, 6
- holomorphically planar curve, 102
- homogeneous contact metric manifolds, 146
- horizontal, 19
- horizontal lift, 19, 170, 184
- horizontal subbundle, 234
- hyper-Kähler manifold, 49, 293
- hypercontact metric structure, 293
- hypercontact structure, 293
- infinitesimal contact transformation, 72
- integral submanifold, 69, 237
- invariant submanifold, 152
- isotropic, 8
- Iwasawa manifold, 240
- K-contact structure, 65, 83, 87
- Kähler, 4
- Kenmotsu manifold, 98
- Kodaira–Thurston manifold, 4
- Lagrangian H-umbilical submanifold, 159
- Lagrangian submanifold, 6
- Legendre curve, 24, 75
- Legendre foliation, 77
- Levi form, 90
- Liouville form, 3, 171, 269
- locally \mathcal{D} -symmetric space, 143
- locally Killing-transversally-symmetric space, 149
- nearly cosymplectic structure, 103
- nearly Kähler, 64, 102
- nearly Sasakian structure, 104
- normal almost contact structure, 80
- normal complex contact metric structure, 251
- normalized complex contact structure, 235
- overtwisted, 33
- Pang invariant, 78
- partially hyperbolic, 265
- periodic point, 38
- planar contact structure, 39
- Plastikstufe, 34
- principal G -bundle, 15
- principal circle bundle, 16
- projective lift, 191, 270
- projectively Anosov flow, 228
- projectivized holomorphic tangent bundle, 269
- PS-overtwisted, 34
- pseudo-Hermitian manifold, 90, 94
- quasi-Kähler, 4, 203
- quasi-regular, 40
- quaternionic Kähler manifold, 242, 297
- R-contact manifold, 40
- real special direction, 284
- Reeb vector field, 23, 247
- regular contact structure, 24
- Ricci form, 204
- Ricci–Kühne, equation of, 152
- Salamon, Bérard-Bergery metric, 243
- Sasaki metric, 171, 184
- Sasakian globally ϕ -symmetric space, 144
- Sasakian locally ϕ -symmetric space, 143
- Sasakian manifold, 86, 97

- Sasakian space form, 140
- second normal space, 258
- semi-Kähler, 257
- special directions, 221
- strict complex contact manifold, 235
- strict contact transformation, 25, 71
- strict infinitesimal contact transformation, 72
- strongly locally ϕ -symmetric space, 146
- strongly pseudoconvex CR-manifold, 90, 94
- symplectic manifold, 1
- symplectic vector field, 11
- symplectomorphism, 1

- tangential lift, 176
- Tanno's equation, 210
- taut contact circle, 35
- Thurston manifold, 4, 217

- tight contact structure, 35
- total scalar curvature, 202, 257
- totally real submanifold, 8, 156
- trans-Sasakian structure, 99
- tridegree, 109
- twistor space, 242

- Veronese embeddings, 154, 261
- vertical, 20
- vertical lift, 170, 184
- vertical subbundle, 235

- weakly locally ϕ -symmetric space, 145
- Webster metric, 90
- Webster scalar curvature, 146, 213, 214

Author Index

- Abbas, C., 39
Abbena, E., 5, 217
Abe, K., 106
Albers, P., 39
Alegre, P., 141
Alekseevskii, D. V., 242
Altschuler, S., 32
Anosov, D. V., 221, 222, 226
Apostolov, V., 58, 130, 131, 202
Armstrong, J., 202
Auslander, L., 221

Bérard-Bergery, L., 243
Baikoussis, C., 164, 166, 258–260,
291, 300, 301
Bang, K., 132–134, 174, 183, 186,
189
Banyaga, A., 40
Barletta, E., 210
Bejancu, A., 90
Belgun, F. A., 89
Bennequin, D., 32, 33
Benson, C., 6
Berger, M., 59, 196, 242

Berndt, J., 147, 148
Besse, A. L., 196, 243
Bishop, R. L., 16, 20, 21
Blair, D. E. (=author), 51, 54, 55,
77, 90, 95, 99, 102–104, 107,
108, 112, 120–122, 124, 127–
129, 132, 134, 141, 143, 145,
148, 149, 164, 166, 175, 179–
181, 199–206, 208, 214, 219,
223, 226–228, 230, 231, 236,
258–262, 272, 280, 283, 284,
291, 300, 301
Boeckx, E., 124–126, 133, 145–147,
176, 180, 181, 183
Boothby, W. M., 13, 36, 74, 175,
233, 234
Borisenko, A. A., 176, 186
Borisovski, I. P., 167
Borrelli, V., 76, 77, 159
Bouche, T., 109
Bourgeois, F., 32
Boyer, C. P., 97, 106, 107, 130–132,
196, 291–293, 295–298

- Brieskorn, E., 105
 Bryant, R. L., 260
 Bueken, P., 141, 145–147, 149

 Calabi, E., 98, 196, 259, 260
 Calvaruso, G., 85, 134, 136, 146, 183
 Cappelletti Montano, B., 78, 127, 128
 Capursi, M., 98
 Carriazo, A., 77, 141
 Cartan, E., 181, 209
 Chūman, G., 196
 Chen, B.-Y., 8, 76, 77, 90, 146, 152, 156, 159, 161, 166
 Chen, H., 128
 Chern, S. S., 43, 94, 154, 206, 213, 257, 258
 Chevalley, C., 45
 Chinae, D., 95, 108, 109, 152
 Cho, J. T., 133, 140, 148
 Cieliebak, K., 39
 Cordero, L. A., 6
 Cowen, M. J., 257, 258
 Crittenden, R. J., 16, 20, 21

 Davidov, J., 201
 de Leon, M., 6, 95, 108, 109
 Deng, S., 209
 Di Terlizzi, L., 78, 127
 Dillen, F., 163, 164, 261
 do Carmo, M. P., 154
 Dolbeault, S., 257, 258
 Dombrowski, P., 88, 171
 Douady, A., 32, 33
 Draghici, T., 58, 130, 131, 202, 204
 Dragomir, S., 90, 210

 Ebin, D., 49, 50, 58, 59, 210
 Eckmann, B., 98
 Ejiri, N., 156
 Eliashberg, Y., 31, 33, 35, 227, 228
 Endo, H., 152, 153
 Erbacher, J., 106

 Fernández, M., 6, 109, 240
 Fetcu, D., 165
 Foreman, B., 235, 237, 243–247, 251, 252, 255, 256, 283, 289
 Fujita, H., 149
 Fujitani, T., 107, 109
 Funabashi, S., 295

 Galicki, K., 97, 106, 107, 130–132, 196, 291–293, 295–298
 Gauntlett, J. P., 131
 Geiges, H., 27, 32, 35, 83, 85, 293
 Ghosh, A., 133
 Ghys, É., 223, 265, 266, 283
 Ginzburg, V., 39
 Goldberg, S. I., 80, 102, 107, 108, 142, 155, 202, 234
 Gompf, R., 6
 González-Dávila, J. C., 149
 González-Dávila, M. C., 149
 Gonzalo, J., 27, 35, 85, 229
 Gordon, C. S., 6
 Gouli-Andreou, F., 133, 134, 226, 258–260
 Gray, A., 6, 41, 99, 203, 240
 Gray, J., 25, 28, 43
 Green, L., 221
 Greenfield, S., 90
 Gromov, M., 4, 8
 Guan, D., 6

 Hadjar, A., 31
 Haefliger, A., 4
 Hahn, F., 221
 Hamilton, R. S., 94, 206, 213
 Han, D.-S., 211
 Harada, M., 153, 155, 167
 Hasegawa, I., 120
 Hatakeyama, Y., 12, 45, 72, 80, 81, 100, 112, 113, 175, 236
 Hervella, L. M., 41, 99, 203
 Hilbert, D., 195
 Hille, E., 267
 Hitchin, N., 293

- Hofer, H., 39
 Holubowicz, R., 6
 Houh, C.-S., 156, 251
 Hsu, C. J., 106, 171
 Huygens, C., 26

 Ianus, S., 91, 200, 201
 Ibáñez, R., 109
 Ikawa, T., 161
 Ishihara, S., 234–237, 241, 242, 251,
 292, 296, 298
 Itoh, M., 66

 Janssens, D., 99
 Jayne, N., 78, 128, 239, 250
 Jelonek, W., 6
 Jerison, D., 216
 Jiménez, J. A., 148

 Kashiwada, T., 293, 294
 Kassabov, O., 155
 Kenmotsu, K., 98, 154
 Kim, J.-S., 129
 Kirchberg, K.-D., 202
 Kirichenko, V. F., 166, 167
 Kobayashi, S., 16, 20, 21, 38, 41, 42,
 80, 154, 233, 234, 236
 Kodaira, K., 4, 80
 Kollár, J., 130, 131
 Kon, M., 154, 155, 161, 189
 Konishi, M., 234, 235, 237, 241, 251,
 292, 295, 297, 298
 Korkmaz, B., 248, 249, 251–254, 262,
 283, 284
 Kosmanek, E., 274
 Koufogiorgos, Th., 124, 127, 128, 133,
 134, 140, 149, 164, 166, 181,
 183, 212, 214, 227
 Kowalski, O., 148, 149, 172, 173
 Kriegl, A., 13, 71, 74
 Kuo, Y.-Y., 291, 292

 Lawson, H. B., 259
 LeBrun, C., 233, 235, 242, 244, 297

 Ledger, A. J., 199, 204, 205
 Lee, J. M., 216
 Libermann, P., 9, 11, 78, 95, 128
 Lie, S., 26
 Ludden, G. D., 156, 236
 Lutz, R., 31, 106
 Lychagin, V., 71

 Mac Lane, S., 26
 Mann, B. M., 97, 293, 295–297
 Markellos, M., 212
 Marle, C.-M., 9, 11
 Marrero, J. C., 95, 99, 108, 109
 Martelli, D., 131
 Martinet, J., 9, 27
 Matzeu, P., 132
 McCarthy, J. D., 6
 McDuff, D., 6
 Meckert, C., 106
 Michor, P. W., 13, 71, 74
 Milnor, J., 146, 223
 Mitric, G., 140, 182
 Mitsumatsu, Y., 227, 228
 Miyahara, Y., 161
 Mogi, I., 155, 271
 Monna, G., 26
 Morimoto, A., 98, 101
 Morimoto, S., 131
 Moroianu, A., 130, 131, 202, 233
 Morrow, J., 80
 Morvan, J.-M., 8, 76, 77, 159
 Moser, J., 9
 Moskal, E., 107, 109, 137, 155
 Mozgawa, W., 6
 Muškarov, O., 202
 Munteanu, G., 270
 Musso, E., 180
 Muto, Y., 196, 198, 199
 Myers, S. B., 120, 130

 Nagano, T., 195
 Nakagawa, H., 261
 Nakamaye, M., 106, 131
 Newlander, A., 79

- Niederkrüger, K., 34
 Nirenberg, L., 79
 Nomizu, K., 16, 20, 21, 41, 80
 Nurowski, P., 202

 O'Neill, B., 143
 Obata, M., 294
 Ogawa, Y., 107, 112, 113
 Ogiue, K., 8, 76, 139, 143, 154–156,
 161, 166, 261
 Oh, W. T., 217
 Okumura, M., 63, 88, 102, 104, 132,
 133, 142, 145, 156, 171
 Olszak, Z., 116, 118, 119
 Oniciuc, C., 165
 Oprea, J., 6
 Oubina, J. A., 96, 99

 Padron, E., 95
 Palais, R. S., 24, 37
 Pang, M.-Y., 78
 Papantoniou, B. J., 124, 127, 212
 Park, J.-S., 217
 Pastore, A. M., 132
 Perrone, D., 85, 108, 132–134, 146,
 148, 149, 180, 181, 203–206,
 210–214, 226, 228, 230, 231,
 293
 Pesin, Y., 265
 Petkov, V., 38
 Popov, G., 38
 Priest, E. R., 136
 Przanowski, M., 202

 Rabinowitz, P. H., 39
 Reckziegel, H., 186
 Reeb, G., 23, 29
 Ros, A., 76, 77, 154, 159
 Rukimbira, P., 40, 66, 107, 114, 120,
 121, 129, 149, 210

 Salamon, S., 243
 Saralegi, M., 6

 Sasaki, S., 25, 29, 70, 80, 81, 105,
 106, 171, 175, 216, 295
 Sato, H., 83, 106
 Seino, M., 120
 Sekigawa, K., 130, 202
 Semmelmann, U., 233
 Sharma, R., 121, 128, 129, 132–134,
 149, 227
 Shibuya, Y., 143, 148, 236
 Showers, D. K., 104
 Sikorav J.-C., 8
 Simanca, S., 196
 Simons, J., 154
 Smolentsev, N. K., 59
 Sparks, J., 130, 131
 Steenrod, N., 75
 Sternberg, S., 10
 Stong, R. E., 25, 27

 Tachibana, S., 88, 107, 171, 292, 293
 Takahashi, T., 105, 143–145
 Tanaka, N., 212
 Tanno, S., 37, 65, 92, 101, 106–108,
 112, 113, 125, 129, 131–133,
 140, 142, 176, 179, 182, 206,
 212, 213, 216, 274, 294–297
 Tashiro, Y., 61, 96, 104, 175, 178
 Taubes, C. H., 39
 Thomas, C. B., 40, 210, 293
 Thurston, W., 4, 31, 35, 227, 228
 Tischler, D., 7
 Tralle, A., 6
 Tricerri, F., 180
 Tripathi, M.M., 129
 Tsihlias, C., 127, 212
 Tsolakidou, N., 134

 Udriste, C., 291
 Urbano, F., 76, 77, 155, 156, 159

 Vaisman, I., 106
 van Coevering, C., 130
 Van de Ven, A., 105

- Vanhecke, L., 99, 134, 141, 143,
 145–149, 176, 180, 181
 VanLindt, D., 155, 163
 Verheyen, P., 155, 163
 Vernon, M., 104
 Verstraelen, L., 154, 155, 163, 261
 Viterbo, C., 39
 Vitter III, A. L., 257, 258
 Vrancken, L., 163, 164, 261, 262

 Waldram, D., 131
 Walters, P., 222
 Wang, H. C., 36, 175
 Watanabe, Y., 148, 149
 Watson, B., 6
 Wegrzynowski, S., 148
 Weinstein, A., 7–9, 38, 39, 76
 Weyl, H., 2
 Wolf, J. A., 29, 233
 Wolfson, J. G., 6
 Wood, C. M., 211, 217
 Woodhouse, N., 9
 Wu, H., 41, 42, 80
 Wu, L., 32

 Xenos, Ph. J., 133, 226

 Yamada, T., 6
 Yamaguchi, S., 161, 196
 Yamato, K., 6
 Yampol'skii, A. L., 176, 186
 Yano, K., 41, 80, 104, 155, 156, 189,
 271, 292
 Yau, S.-T., 8, 156
 Ye, Y.-G., 234
 Yim, J.-W., 211
 Yu, W. N., 292, 293

 Zeghib, A., 120
 Zehnder, E., 39
 Zheng, F., 80